

## **PATIENT INFORMATION LEAFLET**

### **ABTERON**

#### **(Abiraterone Acetate Tablets USP 250 mg)**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Abiraterone Acetate Tablet USP 250 mg is and what it is used for
2. What you need to know before you take Abiraterone Acetate Tablet USP 250 mg
3. How to take Abiraterone Acetate Tablet 250 mg
4. Possible side effects
5. How to store Abiraterone Acetate Tablet USP 250 mg
6. Contents of the pack and other information

#### **1. What Abiraterone Acetate Tablet USP 250 mg is and what it is used for**

Abiraterone Acetate Tablet USP 250 mg contains a medicine called abiraterone acetate. It is used to treat prostate cancer in adult men that has spread to other parts of the body. Abiraterone Acetate Tablet USP 250 mg stops your body from making testosterone; this can slow the growth of prostate cancer.

When Abiraterone Acetate Tablet USP 250 mg is prescribed for the early stage of disease where it is still responding to hormone therapy, it is used with a treatment that lowers testosterone (androgen deprivation therapy).

When you take this medicine your doctor will also prescribe another medicine called prednisone or prednisolone. This is to lower your chances of getting high blood pressure, having too much water in your body (fluid retention), or having reduced levels of a chemical known as potassium in your blood.

## **2. What you need to know before you take Abiraterone Acetate Tablet USP 250 mg.**

Do not take Abiraterone Acetate Tablet USP 250 mg

- if you are allergic to abiraterone acetate or any of the other ingredients of this medicine (listed in section 6).
- if you are a woman, especially if pregnant. Abiraterone Acetate Tablet USP 250 mg is for use in male patients only.
- if you have severe liver damage.

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking this medicine.

### **Warnings and precautions:**

Talk to your doctor or pharmacist before taking this medicine:

- If you have liver problems
- if you have been told you have high blood pressure or heart failure or low blood potassium (low blood potassium may increase the risk of heart rhythm problems)
- if you have had other heart or blood vessel problems
- if you have an irregular or rapid heart rate
- if you have shortness of breath
- if you have gained weight rapidly
- if you have swelling in the feet, ankles, or legs
- if you have taken a medicine known as ketoconazole in the past for prostate cancer
- about the need to take this medicine with prednisone or prednisolone
- about possible effects on your bones
- if you have high blood sugar

Tell your doctor if you have been told you have any heart or blood vessel conditions, including heart rhythm problems (arrhythmia), or are being treated with medicines for these conditions.

Tell your doctor if you have yellowing of the skin or eyes, darkening of the urine, or severe nausea or vomiting, as these could be signs or symptoms of liver problems. Rarely, failure of the liver to function (called acute liver failure) may occur, which can lead to death.

Decrease in red blood cells, reduced sex drive (libido), muscle weakness and/or muscle pain may occur.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking this medicine.

**Blood monitoring:**

Abiraterone Acetate Tablet USP 250 mg may affect your liver, and you may not have any symptoms. When you are taking this medicine, your doctor will check your blood periodically to look for any effects on your liver.

**Children and adolescents:**

This medicine is not for use in children and adolescents. If Abiraterone Acetate Tablet USP 250 mg is accidentally ingested by a child or adolescent, go to the hospital immediately and take the package leaflet with you to show to the emergency doctor.

**Other medicines and Abiraterone Acetate Tablet USP 250 mg**

Ask your doctor or pharmacist for advice before taking any medicine.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is important because Abiraterone Acetate Tablet USP 250 mg may increase the effects of a number of medicines including heart medicines, tranquilisers, herbal medicines (e.g., St John's wort) and others. Your doctor may want to change the dose of these medicines. Also, some medicines may increase or decrease the effects of Abiraterone Acetate Tablet USP 250 mg. This may lead to side effects or to Abiraterone Acetate Tablet USP 250 mg not working as well as it should.

Androgen deprivation treatment may increase the risk of heart rhythm problems. Tell your doctor if you are receiving medicine

- used to treat heart rhythm problems (e.g. quinidine, procainamide, amiodarone and sotalol);
- known to increase the risk of heart rhythm problems [e.g. methadone (used for pain relief and part of drug addiction detoxification), moxifloxacin (an antibiotic), antipsychotics (used for serious mental illnesses)].

Tell your doctor if you are taking any of the medicines listed above.

**Abiraterone Acetate Tablet USP 250 mg with food**

- This medicine must not be taken with food (see section 3, “Taking this medicine”).
- Taking Abiraterone Acetate Tablet USP 250 mg with food may cause side effects.

**Pregnancy and breast-feeding:**

Abiraterone Acetate Tablet USP 250 mg is not for use in women.

- This medicine may cause harm to the unborn child if it is taken by women who are pregnant.
- If you are having sex with a woman who can become pregnant, use a condom and another effective birth control method.
- If you are having sex with a pregnant woman, use a condom to protect the unborn child.

**Driving and using machines:**

This medicine is not likely to affect your being able to drive and use any tools or machines.

**Abiraterone Acetate Tablet USP 250 mg contains lactose and sodium:**

- Abiraterone Acetate Tablet USP 250 mg contains lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.
- This medicine also contains sodium. To be taken into consideration by patients on a controlled sodium diet.

**3. How to take Abiraterone Acetate Tablet USP 250 mg:**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

**How much to take:**

The recommended dose is 1,000 mg once a day.

**Taking this medicine:**

- Take this medicine by mouth.
- Do not take Abiraterone Acetate Tablet USP 250 mg with food.

- Take Abiraterone Acetate Tablet USP 250 mg at least two hours after eating and do not eat anything for at least one hour after taking Abiraterone Acetate Tablet USP 250 mg (see section 2, “Abiraterone Acetate Tablet USP 250 mg with food”).

- Swallow the tablets whole with water.

- Do not break the tablets.

- Abiraterone Acetate Tablet USP 250 mg is taken with a medicine called prednisone or prednisolone. Take the prednisone or prednisolone exactly as your doctor has told you.

- You need to take prednisone or prednisolone every day while you are taking Abiraterone Acetate Tablet USP 250 mg.

- The amount of prednisone or prednisolone you take may need to change if you have a medical emergency. Your doctor will tell you if you need to change the amount of prednisone or prednisolone you take. Do not stop taking prednisone or prednisolone unless your doctor tells you to.

Your doctor may also prescribe other medicines while you are taking Abiraterone Acetate Tablet USP 250 mg and prednisone or prednisolone.

**If you take more Abiraterone Acetate Tablet USP 250 mg than you should**

If you take more than you should, talk to your doctor or go to a hospital immediately.

**If you forget to take Abiraterone Acetate Tablet USP 250 mg:**

- If you forget to take Abiraterone Acetate Tablet USP 250 mg or prednisone or prednisolone, take your usual dose the following day.

- If you forget to take Abiraterone Acetate Tablet USP 250 mg or prednisone or prednisolone for more than one day, talk to your doctor without delay.

**If you stop taking Abiraterone Acetate Tablet USP 250 mg:**

Do not stop taking Abiraterone Acetate Tablet USP 250 mg or prednisone or prednisolone unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. Possible side effects:**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Abiraterone Acetate Tablet USP 250 mg and see a doctor immediately if you notice any of the following:

- Muscle weakness, muscle twitches or a pounding heart beat (palpitations). These may be signs that the level of potassium in your blood is low.

#### **Other side effects include:**

##### **Very common (may affect more than 1 in 10 people):**

Fluid in your legs or feet, low blood potassium, high blood pressure, urinary tract infection, diarrhoea.

##### **Common (may affect up to 1 in 10 people):**

High fat levels in your blood, liver function test increases, chest pain, irregular heart beat (atrial fibrillation), heart failure, rapid heart rate, severe infections called sepsis, bone fractures, indigestion, blood in urine, rash.

##### **Uncommon (may affect up to 1 in 100 people):**

Adrenal gland problems (related to salt and water problems), abnormal heart rhythm (arrhythmia), muscle weakness and/or muscle pain.

##### **Rare (may affect up to 1 in 1,000 people):**

Lung irritation (also called allergic alveolitis).

Failure of the liver to function (also called acute liver failure).

##### **Not known (frequency cannot be estimated from the available data):**

Heart attack, changes in ECG - electrocardiogram (QT prolongation).

Bone loss may occur in men treated for prostate cancer. Abiraterone Acetate Tablet USP 250 mg in combination with prednisone or prednisolone may increase bone loss.

## **5. How to store Abiraterone Acetate Tablet USP 250 mg:**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton, cardboard wallet, and the blister. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information:**

What Abiraterone Acetate Tablet USP 250 mg contains:

- The active substance is abiraterone acetate. Each uncoated tablet contains, Abiraterone Acetate USP .....250mg.
- The other ingredients are Lactose Monohydrate, Microcrystalline cellulose pH-101, Cross Carmellose cellulose, Sodium Lauryl Sulphate, PVP K-30, Purified Water, Colloidal silicone Dioxide, Magnesium Stearate.

### **What Abiraterone Acetate Tablet USP 250 mg looks like and contents of the pack:**

- Abiraterone Acetate Tablet USP 250 mg tablets are an uncoated white to off white colour caplet shape tablet with break line on one side.

### **Packing material:**

60 tablets is packed in 75CC HDPE bottle of 38 mm child resistance cap with silica gel sachet 1 g & nonabsorbent cotton.

### **Marketing Authorization Holder and Manufacturer:**

BLISS GVS PHARMA LTD  
102, Hyde Park, Saki Vihar road,  
Andheri East.  
Mumbai 400072