

## **Patient Information Leaflet**

### **G-DERM Cream**

(Betamethasone Dipropionate, Gentamicin, Tolnaftate & Idochlorhydroxyquinoline Cream)

#### **Read all this leaflet carefully before you start taking this medicine**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist or nurse.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **In this leaflet:**

1. What G-DERM Cream are and what they are used for
2. Before you use G-DERM Cream
3. How to use G-DERM Cream
4. Possible side effects
5. How to store G-DERM Cream
6. Further information

### **1. WHAT G-DERM CREAM IS AND WHAT IT IS USED FOR**

G-DERM Cream is used for relief from itching and inflammation associated with a wide variety of skin conditions like allergic dermatitis, atopic dermatitis, contact dermatitis and plaque psoriasis. G-DERM is even indicated in topical treatment of folliculitis, furunculosis, paronychia and other bacterial skin infections (including types infected insect bites, minor burns, eczematoid dermatitis, seborrheic dermatitis, excoriation, lacerations, skin abscesses and cysts, skin ulcers, stasis ulcers, stings, bacterial infections of minor fungal and viral infections, sycosis barbae and minor surgical wounds) caused by staphylococci, streptococci, proteus vulgaris, Escherichia coli, Pseudomonas aeruginosa and Enterobacter aerogenes.

G-DERM stops the growth of fungi that causes skin infections, including athlete's foot, jock itch, ring worms, infections in nails, scalp, palms and soles. It helps in oxidative damage in Parkinson, Alzheimer, amyotrophic lateral sclerosis and other neurogenerative diseases.

## 2. BEFORE YOU USE G-DERM CREAM

Combining G-DERM with topical anthralin may increase psoriasis symptoms. Therefore, G-DERM should be discontinued 1 week before starting anthralin's. Other skin medications may affect the absorption or effectiveness of G-DERM cream. Avoid using other topical preparations at the same time except under the direction of the physician.

G-DERM is contraindicated in patients with hypersensitivity or allergy to any of its content.

### How to use this medicine

Follows instructions given in to below Table.

For topical use only, which means it is applied directly to the affected area of skin.

Do not use more than the stated dose shown in the Insert.

## 3. HOW TO USE G-DERM CREAM

Wash the infected area and dry it well. As many skin conditions are contagious, you should keep a towel and flannel for your own use and not share it so that you do not infect anyone else.

Apply the cream onto the infected area and surrounding skin.

Unless the infected skin is on your hands, wash your hands carefully after applying the cream to avoid spreading the infection to other parts of the body or to other people. Similarly, clothing which comes into contact with the infected areas, such as socks, should be washed and changed frequently.

### Adults and children aged 10 years and over

---

Age	Dose
Adults and children aged 10 years and over	Put a small amount of the cream on the affected area twice daily. Rub it in gently until the cream has disappeared.

---

Do not use for more than 7 days.  
If symptoms persist talk to your doctor.

---

## **If anyone has swallowed this product**

If anyone accidentally swallows G-DERM Cream, contact a doctor or your nearest Accident and Emergency department (Casualty), taking this leaflet and pack with you.

## **If you forget to use the medicine**

If you forget to use a dose, use the next dose when needed. Do not use a double dose.

## **4. POSSIBLE SIDE EFFECTS**

The most noted side effects of G-DERM are burning at the site of application, itching, irritation (erythema and pruritus) and dryness that did not usually require discontinuation of treatment. Possible photo sensitization has been reported in several patients. Stop using G-DERM if unusual or severe blistering, itching peeling or redness occurs.

If you experience any of the above mentioned symptoms, stop using the medicine and talk to your doctor:

## **5. HOW TO STORE G-DERM CREAM**

Store in cool & dry place, below 30°C.

Do not freeze.

## **6. FURTHER INFORMATION**

What **G-DERM CREAM** contain

Each gram contains:

Betamethasone Dipropionate USP 0.643 mg

Gentamicin Sulphate B.P. equivalent to Gentamicin base 1 mg

Tolnaftate USP 10 mg

Iodochlorhydroxyquinoline 10 mg

Chlorocresol B.P. (Preservative) 1 mg

### **What G-DERM CREAM look like and contents of the pack**

Printed Laminated tube with sealed nozzle fitted with white colour stand up cap.

#### **Presentation:**

15 gm /30 gm tube packed in a mono carton with a leaflet.

#### **Marketing Authorization Holder and Manufacturer**

Bliss GVS Pharma Ltd.

102, Hyde Park, Saki-Vihar Road,

Andheri (East), Mumbai – 400072.

**This leaflet was last approved on September 2020**