

## PACK INSERT

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory.

### MULTIVITAMIN SYRUP

#### **COMPOSITION:**

Each 5 ml Contains:

Vitamin A (Palmitate) BP.....	3600 IU
Thiamine Hydrochloride BP.....	2.0 mg
Riboflavin BP.....	1.2mg
Pyridoxine Hydrochloride BP.....	0.2 mg
Cyanocobalamin BP.....	2.0 mg
Nicotinamide BP.....	10 mg
Ascorbic Acid BP.....	50 mg
Vitamin D3 (Cholecalciferol) BP.....	1200 IU
Appropriate Overages added to compensate for loss on Storage	

#### **PHARMACEUTICAL GROUP WITH ATC Code:**

Pharmacotherapeutic Group: Multivitamins supplement

ATC No.: A11A

#### **INDICATIONS:**

**Multivitamin Syrup** is indicated for:

Loss of appetite and poor eating habits. Anorexia after a brief illness. Patients experiencing loss of appetite during convalescence following surgery or after any major illness.

#### **DOSAGE & ADMINISTRATION:**

1 Year to 6 Year: 1/2 teaspoonful (2.5 ml),

7 Years to 12 Years - 1 teaspoonful (5 ml)

These dosage may be repeated 3 to 4 times daily as required.

#### **Method of Administration**

For oral administration.

#### **CONTRAINDICATIONS:**

Contraindicated in patients known to be hypersensitive to any of its components and in patients with hypervitaminosis.

#### **WARNING & PRECAUTIONS:**

Multivitamins are not recommended for the treatment of severe specific deficiencies of vitamins and minerals. In such cases, the underlying cause should be determined and corrected if possible. They are not intended for the treatment of pernicious anaemia, since neurologic involvement may develop or progress despite temporary remission of anaemia in patients with vitamin B<sub>12</sub> deficiency who receive supplemental folic acid and who are inadequately treated with vitamin B<sub>12</sub>.

While taking the multivitamins, both protein and energy are also required to provide complete nutrition in the daily diet. No other vitamins, minerals or supplements with or without vitamin A should be taken with this preparation except under medical supervision.

Do not take on an empty stomach. Do not exceed the stated dose. Keep out of the reach of children. If symptoms persist, consult your doctor.

**PREGNANCY AND LACTATION:**

Vitamins are generally considered to be safe for use during pregnancy and breast-feeding. Special prenatal vitamins are available and are important for the healthy development of a baby. However, large doses of some vitamins can be harmful. Talk to your doctor about taking Multivitamin Syrup during pregnancy or breast-feeding.

**UNDESIRABLE EFFECTS:**

Some people may have a feeling of lightheadedness, sleepy, blurred eyesight or a change in thinking clearly. Avoid tasks or actions that call for being alert or have clear eyesight until you see how this drug works. Stomach pain. dry mouth may be reported. Unexpected excitement may rarely happen.

**OVERDOSE**

No information available.

**STORAGE:**

Store below 300C, protect from light.

**PRESENTATION:**

100 ml Pet Bottle in a Mono Carton along with Measuring Cup and pack insert

**LEGAL STATUS:**

**Prescription Drug**

**KEEP OUT OF REACH OF CHILDREN.**

**Carefully read the accompanying instructions before use.**

**MANUFACTURED FOR:**

**NEW GLOBAL PHARMACEUTICALS LTD.**

Plot No.1, Light Industrial Area.

ACCRA-GHANA.