

**PATIENT INFORMATION LEAFLET FOR
RECTOSKY OINTMENT**

**(HYDROCORTISONE ACETATE, LIDOCAINE, ZINC
OXIDE & ALLANTOIN OINTMENT)**

What you need to know about RECTOSKY OINTMENT

Your doctor has decided that you need this medicine to help treat your condition.

Read all this leaflet carefully before you start to use the medication. It contains important information.

Keep the leaflet in a safe place because you may want to read it again.

If you have any other questions, or if there is something you don't understand, please ask your doctor or pharmacist.

This medicine has been prescribed for you. Never give it to someone else. It may not be the right medicine for them even if their symptoms seem to be the same as yours.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What **RECTOSKY ointment** is, and what they are used for
2. Things to consider before you start to use **RECTOSKY ointment**
3. How to use **RECTOSKY ointment**
4. Possible side effects
5. How to store **RECTOSKY ointment**
6. Further information

1. What RECTOSKY OINTMENT is and what they are used for

RECTOSKY OINTMENT is indicated in managing inflammatory conditions of anorectal region e.g.

1. Bleeding or non-bleeding haemorrhoids
2. Internal or external haemorrhoids
3. Anal fissure
4. Proctitis
5. Post haemorrhoidectomy pain and discomfort
6. Further information

2. Things to consider before you start to use RECTOSKY OINTMENT

The safety of use of corticosteroids in pregnancy is not established. Topical application of corticosteroids to pregnant animal can cause abnormalities in foetal developments e.g. cleft plate and intra uterine growth retardation. It is advisable therefore to limit use of RECTOSKY OINTMENT during pregnancy, only if in the physician opinion benefits overweight the possible risks. In the absence of improvement duration should be limited to not more than two weeks. In the event of developing irritation the use of RECTOSKY OINTMENT should be discontinued.

3. How to use:

Apply RECTOSKY OINTMENT twice or thrice a day before or after defecation OR as directed by the physician.

If you use more RECTOSKY OINTMENT than you should:

Wash off or wipe out the excess or all ointment applied on the affected area, if still uncomfortable feeling persists contact physician and seek for advice.

If you forget to use RECTOSKY OINTMENT

Look for the next dose schedule, if closer than do not apply, missed dose.

If you stop using RECTOSKY OINTMENT

Consult your doctor or health advisor for discontinuation of treatment.

4. Possible side effects

A medicine is only made for rectal use.

During, immediately, or following application of product, there may be transient stinging or burning from open areas of skin, or transient blanching (lightening), or erythema (redness) of the skin. Please discontinue the use and consult your physician.

5. How to store

Store at cool and dry place, below 30°C.

Do not freeze.

6. Further information

RECTOSKY OINTMENT is contraindicated in persons with history of allergy to any of the constituents. RECTOSKY OINTMENT should not be used in the presence of acute infection of viral or bacterial nature unless a proper antimicrobial cover is provided.

Product contains below excipients along with Active Hydrocortisone acetate and Lidocaine

1. Zinc Oxide
2. Allantoin
3. Microcrystalline wax
4. Light Liquid Paraffin
5. White Soft Paraffin

Product supplied in 30 gm lami tube in a monocarton along with leaflet.

Manufacturer & Marketing Authorization Holder

Kremoint Pharma Pvt. Ltd.

B-8, Additional Ambernath M.I.D.C.,
Ambernath (E), Thane-421506.