

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
1	PRAISE	Phase II/III	1. Oral FT-4202 Pyruvate Kinase Activator 2. Placebo	2nd June 2022	Dr Prince Agyapong	1. Kintampo Health Research Center 2. Ghana Institute of Clinical Genetics, KBTH	Forma Therapeutics, Inc.	Application Approved, 43 Months	Objectives of the study are: 1. To assess the efficacy of FT-4202 in adolescents and adults with SCD as compared to placebo as measured by improvement in hemoglobin (Hb) 2. To assess the efficacy of FT-4202 as compared to placebo on the annualized vaso-occlusive crisis (VOC) rate 3. To measure the effects of FT-4202 on clinical measures and sequelae of hemolysis 4. To evaluate the effects of FT-4202 on the sequelae of VOC 5. To assess changes in fatigue of sickle cell patients taking FT-4202
2	GBT-2104-132	Phase III	1. Inclacumab 2. Placebo	5th July, 2021	Professor Alex Osei-Akoto	Komfo Anokye Teaching Hospital (KATH)	Global Blood Therapeutics, Inc.	Application Approved 2 years	The primary objective of this study is to evaluate the safety and efficacy of a single dose of inclacumab compared to placebo to reduce the incidence of re admission to a healthcare facility for a vaso-occlusive crisis (VOC) after an admission for an index VOC in participants with sickle cell disease (SCD). Additional objectives of the study are to evaluate the pharmacokinetics (PK) and pharmacodynamics (PD) of inclacumab, the presence of anti-drug antibodies (ADAs), and changes in quality of life (QOL).
3	FORTIFIED BULLION CUBES		Shrimp Flavour Stock Cubes	13th December 2021	Prof. Seth Adu-Afarwuah	University of Ghana	Helen Keller International (Through a grant from the Bill & Melinda Gates Foundation)	Application Approved, 9 months	This study aims to assess the impacts of household use of multiple micronutrient-fortified bouillon cubes (containing vitamin A, folic acid, vitamin B12, iron, and zinc in addition to iodine), compared to control bouillon cubes fortified with iodine only, on: a) Micronutrient status among women 15-49 years of age and children 2-5 years of age after 9 months of intervention b) Haemoglobin concentrations among women 15-49 years of age and children 2-5 years of age after 9 months of intervention. c) Breast milk micronutrient among lactating women 4-8 months postpartum after 3 months of intervention.
4	ANTIPSYCHOTIC STUDY	Phase IV	Omega-3 Fatty Acids	15th December 2021	Debrah Akosua Bema	Accra Psychiatric Hospital	Dr. Sammy Ohene. P. O. Box KB 77 Korle-Bu	Application Approved, 29 Weeks	The primary objective of this study is to determine the use of once daily dose of 1000mg omega 3 fish oil as a clinically effective and safe intervention for reducing the burden associated with antipsychotic induced movement disorders. Secondary: To determine the demographic and clinical characteristics of psychiatric patients with antipsychotic induced movement disorder. To determine the efficacy of omega 3 supplementation in relieving the symptoms of AIM disorders To evaluate the impact of omega 3 supplementation on the clinical outcomes of psychosis, cognitive function and quality of life/ adherence of participants. To determine the correlations between the demographic and clinical parameters and the outcomes of therapy To understand the experiences of patients who have used other complementary and alternative medicines aside omega 3 fish oil as adjunct to conventional therapy, in an attempt to be free from their symptoms

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
5	PROBIOTIC		1.Synbiotic (Nutraflora and Maltrin M100 P-95 and L. plantarum (Lp) 2.Placebo	27th July, 2021	Dr Seyram Kaali	Kintampo Municipal Hospital	Dr. Kwaku Poku Asante	Application Approved 6 months	Primary A pilot trial to evaluate the administration of probiotic supplementation among pregnant women in the third trimester and effective colonization of the gut microbiome of their infants one-month post-partum. Secondary 1. To assess compliance of administering a synbiotic product (L. plantarum with Fructooligosaccharide) among pregnant women. 2. To assess birth outcomes among participants who receive synbiotic products compared to those on placebo. 3. To assess if maternal stool microbiome profoundly changes from immediately after childbirth to one-month post-partum. 4. To characterize the diversity of vaginal microbiomes among pregnant women in the study area. 5. To determine the safety of the probiotic supplementation among pregnant women from 5 to 6 months until up to two weeks post partum.
6	GBT 2104-131	Phase III	1. Inclacumab 2.Placebo	5th July, 2021	Professor Alex Osei-Akoto	Komfo Anokye Teaching Hospital (KATH)	Global Blood Therapeutics, Inc.	Application Approved 2 years	The primary objective of this study is to evaluate the safety and efficacy of treatment every 12 weeks with inclacumab to reduce the incidence of VOCs in participants with SCD. Additional objectives of the study are to evaluate the pharmacokinetics (PK) and pharmacodynamics (PD) of inclacumab, the presence of anti-drug antibodies (ADAs), and changes in quality of life (QOL).
7	COVID 19 CHO-CELL	Phase II/III	1.Recombinant two-component COVID-19 vaccine (CHO cell) 2. ReCOV Placebo	16th November 2021	Dr. Patrick Ansah	1. Dodowa Health Research Centre 2. Navorongo Health Research Centre.	Jiangsu Recbio Technology Co., Ltd.	Application Approved, 13 months	1. To evaluate the safety and reactogenicity of the recombinant two-component COVID-19 vaccine (CHO cell) (ReCOV for short) in adults aged 18 years and older. 2. To evaluate SARS-CoV-2 neutralizing antibody of ReCOV on Day 14 after 2 doses vaccination in adults aged 18 years and older. 3. To evaluate the efficacy of ReCOV in preventing RT-PCR confirmed symptomatic COVID-19 in adults aged 18 years and older. 4. To evaluate the safety and reactogenicity of ReCOV in adults aged 18 years and older.
8	EBSI-LSV	Phase I	1.EBSI-LSV 2. Placebo	1st September 2021	1.Dr Seyram Kaali 2.Dr.Patrick Ansah	1.Kintampo Health Research Centre 2.Navrorongo Health Research Centre	Emergent BioSolutions (EBS)	Application Approved 2 years	1. To evaluate the safety and tolerability of increasing dose levels of EBS-LASV vaccine administered as a single dose or two-dose series. 2. To evaluate the humoral immune response to EBS-LASV vaccine at various dose levels and dosing schedules for the purpose of selecting two regimens (dose and schedule) for further evaluation in a Phase 2 study.
9	ASAAP	Phase III	1. Artemether Lumefantrine 2.Atovaquone-Proguanil 3. Placebo of Atovaquone-Proguanil	4th October 2021	1. John Humphrey, AMUASI 2. Dr Oumou Maiga Ascofare	St. Francis Xavier Hospital	Kumasi Centre for Collaborative Research (KCCR), Kumasi, Ghana	Application Approved 21 months	The overall aim of this phase III clinical trial(main study = study II) is to develop a readily deployable highly efficacious, safe and well tolerated antimalarial triple combination therapy for young children. This is achieved by evaluating the efficacy, safety and tolerability of artemether-lumefantrine (AL) + atovaquone-proguanil (AP) tri-therapy (AL+AP) compared to standard AL therapy (+placebo) for the treatment of uncomplicated Plasmodium falciparum malaria in African children aged 6 to 59 months

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
10	POLYPHENOL-RICH COCOA POWDER TRIAL		Polyphenol-rich natural cocoa powder	10th January 2022	Prof. George Obeng Adjei	Ga East Municipal Hospital, Ghana Infectious Disease Centre	Ghana Cocoa Board	Application Approved, 4 Months	General objective is to evaluate effects of polyphenol-rich cocoa as adjuvant therapy in COVID 19 patients. Specific objectives: 1. to determine the effects of natural polyphenol-rich natural cocoa powder (5 % v/w) (as adjuvant therapy) on symptom resolution and illness duration in COVID-19 patients 2. to determine the effects of natural polyphenol-rich natural cocoa powder (5 % v/w) on selected markers of coagulopathy in COVID-19 patients 3. to determine the effects of natural polyphenol-rich natural cocoa powder (5 % v/w) on virologic clearance COVID-19 patients 4. to determine the effects of natural polyphenol-rich natural cocoa powder (5% v/w) on disease prognosis COVID-19 patients
11	PIVOT STUDY	Phase II	1.Hydroxyurea 2.Placebo	18th June 2021	Dr. Yvonne A. Dei-Adomakoh	Korle-Bu Teaching Hospital	Cincinnati Children's Hospital Medical Center	Application Approved 5 years	To measure the toxicities of hydroxyurea treatment on laboratory parameters. To assess the effects of hydroxyurea treatment on a variety of sickle-related clinical and laboratory parameters in a large cohort of children and adults with HbSC disease. To identify which study endpoints are suitable for a future Phase III trial of patients with HbSC disease receiving hydroxyurea therapy.
12	RECOVERY	Phase III	1.Dexamethasone 2.Empagliflozin	21st May, 2021	Dr. John H. Amuasi	Komfo Anokye Teaching Hospital Ghana Infectious Disease Centre	University of Oxford Clinical Trials and Research Governance.	Application Approved 2 years	For each pairwise comparison with the 'no additional treatment' arm, the primary objective is to provide reliable estimates of the effect of study treatments on all-cause mortality at 28 days after randomisation (with subsidiary analyses of cause of death and of death at various timepoints following discharge). The secondary objectives are to assess the effects of study treatments on duration of hospital stay; and, among patients not on invasive mechanical ventilation at baseline, the composite endpoint of death or need for invasive mechanical ventilation or ECMO.
13	VR-AD-1005 STUDY	Phase II	VR-AD-1005	1st July 2021	Dr. Ernest Kenu	Pentecost Hospital, Madina, Madina Polyclinic –	Vanessa Research Holdings, Inc.,	Application Approved. Study not yet commenced 1 year 2 months	To assess the efficacy and safety of VR-AD-1005 for the treatment of acute diarrhea in cholera in combination with standard rehydration treatment with or without antibiotics (as indicated by WHO or other applicable guidelines) versus standard treatment alone. Efficacy is measured as reduction in stool output and/or duration of diarrhea between the start of treatment until final diarrheal stool before recovery or end of study treatment (treatment duration 120 hours).
14	HOPE KIDS 2	Phase III	1.Voxelotor 2.Placebo	16th December 2020	Dr. Catherine Segbefia	*Korlebu Teaching Hospital Department of Child Health *Sickle cell office Directorate Child(KATH)	Global Blood Therapeutics, inc	Application Approved. Study not yet commenced 38 Months	The purpose is to evaluate the effect of voxelotor compared to placebo on the transcranial Doppler(TCD) time-averaged mean of the maximum velocity(TAMMV) arterial cerebral blood flow at 24 weeks in SCD participants >2 to < 15 years of age with conditional (170 to <200cm/sec) TCD flow velocity.
15	VAT00008	Phase III	1.SARS-CoV2 prefusion Spike delta TM with AS03 adjuvant, monovalent 2.SARS-CoV2 prefusion Spike delta TM with AS03 adjuvant, bivalent 3.Matching placebo	26th May, 2021	Dr. Kwaku Poku Asante	*Navrongo Health Research Centre *Kintampo Health Research Centre *Kwame Nkrumah University of Science and Technology (KNUST)	SANOFI	Application Approved. Actively Enrolling at KCCR and Navrongo while Kintampo closed enrolment 18 months	To assess, in participants who are SARS-CoV-2 naïve, the clinical efficacy of the CoV2 preS dTM-AS03 vaccines for the prevention of symptomatic COVID-19 occurring ≥ 14 days after the second injection. To assess the safety of the CoV2 preS dTM-AS03 vaccines compared to placebo throughout the study.
16	BURULIRIFDAC	Phase III	1.Rifampicin 2.Clarithromycin 3.Dialkylcarbamoyl chloride (DACC) Dressing	12th December 2020	Prof. Richard Phillips	*KCCR *Ga East municipal hospital *Pakro Health Centre *Wassa Amenfi East Hospital	London school of Hygiene and Tropical Medicine	Application Approved. Study not yet commenced 2 Years 6 Months	Compare the time to clearance of viable Mycobacterium from wounds of patients treated with high-dose rifampicin and DACC dressings (HR-DACC) to those receiving standard dose rifampicin and DACC dressings

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
17	EMODEPSIDE	Phase II	Emodepside (5mg)	5th November, 2020	Dr. Nicholas Opoku	<ul style="list-style-type: none"> •School of Public Health Research Centre, (UHAS). •Municipal Hospital, Hohoe, Volta Region, Ghana •Kpassa, Nkwanta-North District, Oti Region, Ghana 	DNDi (Drugs for Neglected Diseases initiative)	Application Approved. Study commenced 67 months	<p>The purpose of this study is to</p> <ul style="list-style-type: none"> •Ensure the safety and tolerability of emodepside after single oral doses administered as solution (liquid service formulation, LSF) or immediate release (IR) tablets in healthy male subjects <p>•Plasma PK of emodepside (solution and tablets), the effect of food on the bioavailability of emodepside</p>
18	BURULINOX	Phase III	<ol style="list-style-type: none"> 1. Nitric Oxide generating dressing (EDX110TM) 2. Vaseline Gauze dressing materials 	24th September 2018	Prof. Richard Odame Phillips	<ol style="list-style-type: none"> 1. Kumasi Centre for Collaborative Research in Tropical Medicine 2. Agogo Presbyterian Hospital 3. Tapa Government Hospital 4. Dunkwa Government Hospital 	Kumasi Center For Collaborative Research (KCCR)	Application Approved. Study yet to commence 36 MONTHS	<p>Buruli ulcer is a neglected disease caused by infection with Mycobacterium ulcerans (Mu), which manifests as large, disfiguring skin ulcers mainly in children aged 5 to 15 years. Access to treatment in rural areas can be challenging and late presentation is typical, due to fear, stigma, suspicion about conventional medicine and economic consequences for poor families. The current recommended regimen of oral rifampicin together with intramuscular streptomycin or clarithromycin for 8 weeks is far from ideal, particularly given the increasing global threat of antimicrobial resistance. Although the disease can be cured in most patients who adhere to this regimen, healing rates are highly variable even in patients with seemingly similar lesions.</p> <p>The purpose of the study is to compare the healing measured by the percentage area reduction of EDX110 dressing with oral rifampicin and clarithromycin (EDX-RC) versus 'Usual Care' with routine Vaseline gauze dressing and oral rifampicin and clarithromycin (VG-RC).</p>
19	TyVEGHA	Phase IV	<ol style="list-style-type: none"> 1. Typbar TCV (Vi polysaccharide-tetanus toxoid conjugate vaccine) 2. Meningococcal Group A conjugate vaccine (MCV-A 5) 	3rd March 2021	Prof. Ellis Owusu-Dabo	Agogo Trial Center/ KNUST- International Vaccine Institute (IVI) Collaborating Center	International Vaccine Institute	Application Approved. Study commenced 3 Years 5 months	<p>The purpose of the study is to</p> <ul style="list-style-type: none"> •To determine the total protection conferred by single-dose vaccination with Vi-TT against blood culture-confirmed symptomatic S. Typhi infection in the intervention vaccine clusters, compared with the control vaccine clusters • To investigate the safety outcomes associated with Vi-TT vaccination in the intervention vaccine recipients compared with the comparator vaccine recipients • To determine the overall protection of Vi-TT vaccination against blood culture-confirmed symptomatic infection caused by S. Typhi in intervention clusters compared with control clusters • To determine the total protection of Vi-TT vaccination against severe TF in the intervention vaccine recipients compared with the comparator vaccine recipients • To determine the overall protection of Vi-TT vaccination against severe TF caused by S. Typhi in intervention clusters compared with control clusters • To investigate the total protection of Vi-TT vaccination against clinical TF (defined below in "Trial Outcome Measures") in the intervention vaccine recipients compared with the comparator vaccine recipients • To investigate the overall protection of Vi-TT vaccination against clinical TF in intervention clusters compared with control clusters • To measure the indirect protection conferred by single-dose vaccination with Vi-TT against blood culture-confirmed symptomatic S. Typhi infection in the intervention vaccine clusters, compared with the control vaccine clusters • To investigate the immunogenicity profile in a subset of Vi-TT recipients compared with the comparator vaccine recipients.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
20	SPUTNIK LIGHT	Phase III	1.Sputnik Light Vector Vaccine 2.Placebo	5th March 2021	1. Dr. Nana Akosua Ansa 2. Dr. Alberta Amu	1. Navrogo Health Research 2. Centre Dodowa Health Research Centre Ghana	Human Vaccine LLC	Application Approved Enrolment closed participants are in follow up 8 months	The purpose of the study is to <ul style="list-style-type: none"> Assess efficacy of the Sputnik-Light vector vaccine against the SARS-CoV-2-induced coronavirus infection compared to placebo Assess tolerability and safety of the Sputnik-Light vector vaccine against the SARS-CoV-2-induced coronavirus infection compared to placebo Assess humoral immunogenicity of the Sputnik-Light vector vaccine against the SARS-CoV-2-induced coronavirus infection compared to placebo on Subset A . Assess protective properties of the SputnikLight vector vaccine against the SARSCoV-2-induced coronavirus infection compared to placebo for prevention of serologically confirmed SARS-CoV-2 infection Assess efficacy of the Sputnik-Light vector vaccine against the SARS-CoV-2-induced coronavirus infection compared to placebo based on severity of COVID-19 disease
21	SHEA LIDO	Phase III	1.Optilube Active Sterile Lubricating Jelly 2.Shealube	10th September 2020	Dr. Kekeli Kodjo Adanu	Ho Teaching Hospital	University of Health and Allied Sciences	Application Approved Study commenced 12 months	This study is a randomized controlled trial which compares the effectiveness, complications and ease of use of shea butter as a surgical lubricant to lidocaine gel. The purpose is to: <ul style="list-style-type: none"> To determine the ease of use of shea butter by clinicians as compared to lidocaine gel as a lubricant for rectal examination. To determine the complication rate related to the use of shea butter as a lubricant for rectal examination. To ascertain the complication rate associated with the use of lidocaine gel as a lubricant for rectal examination To compare the complication rate related to the use of shea butter to that of lidocaine gel.
22	CECOLIN	Phase III	1.Cecolin® 2.Gardasil®	1st September 2020	Prof. Tsiri Agbenyega	•Agogo Asante Akim North District	PATH	Application Approved 30 months	The purpose of this study is to demonstrate the non-inferiority of Cecolin® administered on 0, 6-month; 0, 12-month; and 0, 24-month two-dose regimens, to Gardasil® using a 0, 6-month two-dose regimen, based on HPV Immunoglobulin G (IgG) antibody levels measured one month after the last dose for HPV types 16 and 18.
23	ASTAWOL	Phase II	1.Rifampicin 2.Albendazole	25th June 2020	Prof. Alexander Yaw Debrah	•Bawku west •Buisa South •Nabdam Fumbisi •Garu-Tempane •Kayoro	Kumasi Centre for Collaborative Research (KCCR), Kumasi, Ghana	Application Approved Actively Enrolling 24 months	The purpose of this study is to <ul style="list-style-type: none"> To show efficacy (Depletion of Wolbachia) of the combination of Rifampicin plus Albendazole against lymphatic filariasis using PCR compared to treatment with albendazole and "no treatment" (other than ivermectin) - Lymphatic Filariasis (LF) trial To show efficacy (depletion of Wolbachia and interruption of embryogenesis in female adult worms) of the combination of Rifampicin plus Albendazole, using PCR and immunohistology compared to treatment with albendazole and "no treatment" (other than ivermectin) – Onchocerciasis trial
24	CROWN CORONATION	Phase III	1.Measles Rubella Vaccine 2.Matching Placebo 3.AstraZeneca vaccine	7th September 2020	Prof. Kwadwo Koram	•Ga East Municipal Hospital •Korle-Bu Teaching Hospital •UGMC •Effia-Nkwanta Hospital •Pentecost Treatment Center	Each country serves as its own sponsor but will receive funding from the Covid 19 Therapeutics Accelerator and Gates Foundation through Washington University in St. Louis.	Application Approved Enrolment closed, Participants are receiving treatment 8 Months .	The purpose of this study is to determine that MR vaccine increases the likelihood of making the specific AstraZeneca COVID-19 vaccine more effective in people with prior exposure to the MR vaccine. This study has two different groups: one group will receive the active MR vaccine and one will receive a placebo. Thirty and sixty days later, participants in each group will receive the AstraZeneca COVID-19 vaccine.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
25	MDGH-MOX	Phase I	Moxidectin tablet (2mg)	February 2020	Dr. Nicholas Opoku	School of Public Health Research Centre, University of Health and Allied Health Sciences, Ho.	Medicines Development for Global Health	Application Approved Actively Enrolling 12 months	To characterize the pharmacokinetics and safety of moxidectin in children (aged 4 to 11 years) and adolescents (aged 12 to 17 years) and to enable determination of an optimal dose for treatment of children 4 to 11 years
26	INOVIO	1b	1.INO-4500 2.CELLECTRA™ 3.SSC-0001	30th September 2019	Prof. Kwadwo Ansah Koram	Noguchi Memorial Institute for Medical Research University of Ghana, Legon	Inovio Pharmaceuticals, Inc	Application Approved Actively Enrolling 20 Months	The LASV DNA vaccine expressing the glycoprotein precursor (LASV GPC, Josiah strain matched) paired with intradermal EP is a promising vaccine platform that has been shown to elicit protective immunity and completely protect guinea pigs and non-human primates (NHP) against viremia, illness (acute and chronic), and death after Lassa virus exposure [26, 27] and protect NHPs from hearing loss [unpublished data]. This LASV DNA vaccine, INO-4500, targets GPC because it represents the most conserved region in this genetically diverse virus. In the case of Lassa virus infection, the generation of a robust T cell response appears to be the key to protection from infection. As such, the DNA-EP platform is highly amenable to this disease target. The purpose of this study is to evaluate the tolerability and safety of INO-4500 administered by ID injection followed by EP in healthy adult volunteers
27	STAND	Phase III	1.CRIZANLIZUM AB 2.PLACEBO	30th September, 2019	1.Dr. Yvonne Dei Adomakoh Dr. Vivian Paintsil	1.Ghana Institute of Clinical Genetics, Korle-Bu Sickle Cell Office Directorate of Child Health,	Novartis Pharma AG	Application Approved. Enrolment closed, participants are receiving treatment 8 years 5 months	Sickle cell disease (SCD) is a genetic blood disorder, caused by a single missense mutation in the β-globin gene, progresses into a systemic disease. Vaso-occlusion is the hallmark of SCD and can lead to serious acute and chronic complications. Extensive preclinical data has established P-selectin as a key mediator of VOC in SCD and suggest that its blockade or genetic absence of P-selectin decreases or eliminates its interactions with its ligands, thereby reducing vaso-occlusion. Crizanlizumab is a monoclonal antibody that binds to P-selectin preventing it interactions with its ligands. The purpose of this study is to compare the efficacy and safety of 2 doses of crizanlizumab (5.0 mg/kg and 7.5 mg/kg) versus placebo in adolescent and adult SCD patients (12 years and older) with history of VOC leading to healthcare visit.
28	MAL 094	Phase IIb	1.RTS,S/AS01E 2.Rabies vaccine (Rabipur™)	21st November 2016	Prof. Tsiri Agbenyega	Malaria Research Center, Agogo	GlaxoSmithKline Biologicals SA	Enrollment ended; participants receiving treatment 72 months	As part of GSK and PATH's commitment to develop a malaria vaccine for reduction of malaria disease burden in children and contribution to the malaria elimination goal, characterization of an optimal dosing regimen and boosting schedules are critical. Results of previous efficacy study MAL 055, including the long term follow-up data and efficacy of a fourth dose administered 18 months after the third dose, and the preliminary results of MAL 071 study (recent controlled human malaria infection) were reviewed by the European Medicines Agency (EMA). There was evidence that demonstrated superior protection against malaria infection associated with the use of a fractional third dose in a 0, 1, 7-month schedule with a higher vaccine efficacy against malaria infection. This study intends to establish Proof of Concept for a fractional dose schedule under conditions of natural exposure. The study will be conducted in children 5-17 months old at first vaccination living in areas of mid to high malaria transmission, in line with the age group recommended by the World Health Organization. Results from study will be critical in informing future possibilities for the development of vaccine-based strategies which, in combination with other interventions, may contribute to the malaria elimination agenda.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
29	AVAREF TV ROTA	Phase III	1.Trivalent Rotavirus P2-VP8 Subunit Vaccine 2.Rotarix®	9th April, 2019	1.Prof. George E. Armah 2.Dr. Alberta Amu	Dodowa Health Research Centre	PATH	Approved study commenced 48 Months	Diarrhea is the second-leading cause of death worldwide among children under the age of five, killing an estimated three quarters of a million children annually and hospitalizing millions more in developing countries. The most common cause of infantile diarrhoea is rotavirus and almost all children are infected by their third birthday regardless of geographical area or economic status. Infection is primarily via fecal oral route and improved sanitation alone will not control infection. Oral rotavirus vaccines have traditionally shown lower efficacy in Low and Middle Income Countries (LMICs) as compared to developed countries. Several theories proposed for this observation includes interference by other intestinal viruses or bacteria, neutralization of vaccine by maternally virus by maternally derived antibodies in breastmilk, etc. Some of these challenges may be obviated by a parenteral administered rotavirus vaccine. This study is therefore to demonstrate the efficacy and safety of the parenteral trivalent rotavirus vaccine in healthy infants (≥6 and <8 weeks old) to prevent severe rotavirus gastroenteritis compared with the orally approved Rotarix®
30	ANTICOV	Phase III	1.Nitazoxanide 2.Ciclesonide 3.Paracetamol 4.Ivermectin 5.Artesunate Amodiaquine (ASAQ)	15th July, 2020	John Humphrey, AMUASI	Komfo Anokye Teaching Hospital	•Bernhard Nocht Institute for Tropical Medicine	Approved study commenced 24 Months	The purpose of this study is to compare the efficacy of alternative treatment strategies versus control on the risk of progression to severe respiratory disease. As there is no validated animal model for COVID-19, the efficacy of any potential treatment remains speculative beyond what is known about their pharmacokinetic and in-vitro data. Several repurposed drugs are currently being tested in severe cases or as prophylaxis, and the results may become available by the time the present study is initiated. At the same time, a number of other drug candidates are being evaluated for in-vitro efficacy or in small proof-of concept studies.13 In view of the rapidly evolving landscape in Africa, it was decided to select an adaptive design for the study in order to allow for the flexibility of adding or dropping arms or adjusting the randomisation ratio based on the data as it becomes available. Additionally, given that the control arm in the study may not be acceptable in some countries, it was decided to adopt a master platform-based approach to be allow for integration of data from all sites in the interim analyses, irrespective of their ability to have randomised patients in all treatment arms..
31	LETICIA	Phase II	1.LETICIA protocol diet (provided by study) 2. 3-Fer syrup 3. Usual or Typical diet	30th August, 2019	Dr. Lawrence Osei-Tutu	Agogo Presbyterian Hospital	Dr. Lawrence Osei-Tutu	Approved, yet to start 12 Months	Iron deficiency is the most common nutritional deficiency worldwide and an important public health problem in Low and Middle Income Countries (LMICs). Causes of anemia in LMICs like Ghana are usually multifactorial including malaria, hemolytic anemias, and chronic blood loss from chronic parasitic infections including schistosomiasis and hookworm. Factors accounting for inadequate supplies of dietary iron and micronutrients include poverty, a lack of nutritional supplementation, and food taboos. Anemia may result when iron deficiency is severe, after the body's iron stores are depleted and supply to the bone marrow is limited. This proof of concept study is to determine whether hospitalized children 6-59 months old who presented with moderate-to-severe anemia and given a combination of iron-rich food and standard iron replacement therapy (the intervention group) will demonstrate a greater final hemoglobin (Hb) concentration after two weeks compared to participants of similar characteristics in the control group who will receive oral iron supplementation in addition to their usual diet.
32	ROBOCOW		0.2% Chlorhexidine Digliconate	10th January 2023	Dr. Mohammed Sheriff	Tamale Teaching Hospital		5 Months	Primary Objective 1. To determine whether perioperative use of 0.2% chlorhexidine mouth wash reduces the rate of postoperative respiratory tract infections in 30 days postoperative period compared to placebo among patients undergoing midline laparotomy. Secondary Objectives 1.To assess the impact of the intervention on 30-day postoperative mortality 2.To determine the impact of the intervention on length of hospital stay 3.To determine whether the intervention impacts on the 30-day unplanned readmission rates due to a respiratory complication 4.To assess the effect of the intervention on time to return to normal activities

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
33	VERTEX Trial	Phase II/III	VX-147	23rd December 2022	Professor Sampson Antwi	Komfo Anokye Teaching Hospital (KATH)	Vertex Pharmaceuticals Incorporated	4 years	<p>Primary objectives</p> <ul style="list-style-type: none"> •To evaluate the efficacy of VX-147 to reduce proteinuria •To evaluate the efficacy of VX-147 on renal function as measured by eGFR slope <p>Secondary objectives</p> <ul style="list-style-type: none"> •To evaluate the efficacy of VX-147 to decrease the risk of the composite clinical outcome •To evaluate the safety and tolerability of VX-147 •To identify the optimal dose from Phase 2 to carry forward to Phase 3 •To characterize the plasma pharmacokinetics (PK) of VX-147
34	CIELO Trial	Phase III	Satralizumab	20th December 2022	Prof. Fred Stephen Sarfo	Komfo Anokye Teaching Hospital (KATH)	F-Hoffman LA Roche/ Chugai Pharma Co. LTD	5years 5months	<p>This study will evaluate the efficacy, safety, pharmacokinetics, and pharmacodynamics of satralizumab compared with placebo in each of the following cohorts:</p> <ul style="list-style-type: none"> •NMDAR autoimmune encephalitis (AIE) cohort: adults and adolescents with definite or probable NMDAR encephalitis •LG11 AIE cohort: adults with LG11 encephalitis <p>In addition, the study will assess the long-term safety and efficacy of satralizumab during an optional extension period. For efficacy analyses, each cohort will be treated as a separate population and will have independent Type I error control at a 5% significance level. Specific primary and secondary objectives and corresponding endpoints for the study are outlined below.</p>
35	SWIS (STERILE WATER INJECTION)		Sterile Water Injection	6th December 2022	Prof. Sue Kruske	Korle-Bu Teaching Hospital (KBTH)	Dr. Jonas Awuku Afari	40 Months	<p>Main Aim This study explores the feasibility, acceptability, and outcomes of implementing sterile water injections (SWI) for the management of lower back pain among birthing women in Ghana.</p> <p>Specific Objectives</p> <ol style="list-style-type: none"> 1. Develop and deliver a training package for midwives on sterile water injections for managing lower back pain. 2. Undertake implementation study in a tertiary hospital in Ghana to assess the feasibility and acceptability of implementing SWI for lower back pain. 3. Determine birth and neonatal outcomes of women with back pain who receive SWI 4. Explore the experiences of women who have had SWI for back pain in labour 5. Explore the experiences and perception of midwives and stakeholders regarding the implementation of SWI for managing back pain in labouring women.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
36	HU PHARMACOGEN OMICS		Hydroxyurea	5th October 2022	Prof Daniel Ansong	KNUST University hospital	Muhimbilla University of Health and Allied Science Haematology and clinical Research Lab Tanzania	27 Months Application Pending Approval,	<p>Specific Primary Objectives</p> <ol style="list-style-type: none"> To evaluate the pharmacogenomic response to hydroxyurea in SCD in the three SCD populations. The mechanism of action of Hydroxyurea (HU) is through increasing erythropoiesis and reducing hemolysis. However, there is variability in response with up to 20% of patients having poor or minimum response. We will evaluate genomic factors implicated in determining the response. To identify early predictive markers of HU response in the three SCD populations. The ability to predict HU response early enough is important in SCD management especially in low resource settings. We will evaluate potential markers of response including hematological markers (F cells and F-reticulocytes, (erythrocytes and reticulocytes containing considerable amount of HbF, respectively), molecular marker (expression of γ-globin mRNA) and genetic markers (pharmacogenomics). The ultimate goal is to be able to stratify patients based on the likelihood of responding to HU and hence facilitate precision medicine for HU in Tanzania.
37	ACTIV TRIAL	Phase III	S-217622	27th September 2022	Dr. Patrick Ansah	Navrongo Health Research Centre	SHIONOGI INC. & Co Ltd	Application Pending Approval,,16 Months	<p>Primary Objective</p> <p>To determine if S-217622 will reduce the time to sustained symptom resolution through Day 29. Time to sustained symptom resolution is defined as the time from start of study intervention to the first day of 4 consecutive days with complete resolution of 13 COVID-19 symptoms on participant self-assessment AND alive and without hospitalization for any reason by Day 29. Hospitalization is defined as ≥ 24 hours of acute care, in a hospital or similar acute care facility, including</p>
38	COPE TRIAL		(i) Healeanlo silicone lady Drain Valve menstrual Cup (ii) Foley catheter will connect the cup to a leg bag (cup+)		Dr. Gabriel Y.K. Ganyaglo	1. Mercy Women's Catholic Hospital in Mankessim 2. Tamale Fistula Center in Tamale	Korle Bu Teaching Hospital	Application Pending Approval, 15 Months	The aims of the study are to examine the effectiveness, comparative effectiveness, and acceptability of two vaginal menstrual cup models (cup and cup+) as a temporizing alternative to managing urinary leakage from vesico-vaginal fistula in both a clinical setting and a community setting, and to quantify non-surgical fistula management costs.
39	INO-9112 COVID 19	Phase I	1. INO-4800 followed by Electroporation (EP) 2. NO-4800 + INO-9112 followed by Electroporation	30th June 2022	Dr. Kwadwo Ansah Koram	Noguchi Memorial Institute for Medical Research, University of Ghana, Legon	Inovio Pharmaceuticals	Application Pending Approval, 15 Months	The overall purpose of this clinical trial is to identify a booster dose of INO-4800 or INO 4800 plus INO-9112 given 6 to 12 months following primary vaccination with an approved or authorized mRNA vaccine for future development.
40	ABDOV COVID-19 TRIAL	Phase III	SCTV01E (A COVID-19 Alpha/Beta/Delta/Omicron Variants S-Trimer Vaccine)	17th June 2022	1. Dr. Alberta Amu 2. Dr. Patrick Ansah 3. Dr. John Amuasi 4. Dr Kwaku Poku Asante	1. Dodowa Health Research Centre 2. Navrongo Health Research Centre 3. Kumasi Center for Collaborative Research (KCCR) 4. Kintampo Health Research Centre	Sinocelltech Ltd..	Application Pending Approval, 19 Months	<p>Stage 1 immunization</p> <p>To evaluate the protective efficacy of SCTV01E against symptomatic COVID-19 occurring from 14 days after the 2nd dose in population previously unvaccinated with COVID-19 vaccine. <input type="checkbox"/> To evaluate the protective efficacy of SCTV01E against moderate and above COVID-19, severe and above COVID-19, hospitalization due to COVID-19, and death due to COVID-19 occurring from 14 days. <input type="checkbox"/> To evaluate the protective efficacy of stage 1 immunization against different SARS-CoV-2 variants. <input type="checkbox"/> To evaluate the safety of SCTV01E in stage 1.</p> <p>Stage 2 immunization</p> <p>To evaluate the protective efficacy of SCTV01E against symptomatic COVID-19 occurring from 7 days after the 3rd dose in population previously unvaccinated with COVID-19 vaccine <input type="checkbox"/> To evaluate</p>

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
41	NOVIC TRIAL	Phase III	Jada System (Intrauterine Vacuum Induced Hemorrhage Control Device)	5th April 2022	Dr. Samuel A. Opong	1. Korle-Bu Teaching Hospital (KBTH) 2. Komfo Anokye Teaching Hospital (KATH)	Women and Infants Hospital of Rhode Island	Application Pending Approval, 48 Months	<p>Study Objectives</p> <ol style="list-style-type: none"> To evaluate the effectiveness of the Jada® System, compared to standard care, in treating PPH, as measured by maternal survival without surgical intervention. To assess the safety of the Jada® System, compared to standard care, in treating PPH, as measured by rate of composite adverse events potentially related to the device, including genital tract injury, uterine perforation or rupture and endometritis. To estimate the cost-effectiveness of the Jada® System, compared to standard care, in treating PPH, as measured by incremental cost per quality-adjusted life year.
42	POST MASTECTOMY PAIN RELIEF		Erector Spinae block using bupivacaine	2nd December 2021	Dr. Nana Addo Boateng	Komfo Anokye Teaching Hospital (KATH)	Self-Funding	Application Pending Approval	<p>General objective: The main objective of the study is to determine the postoperative analgesic effect of Erector Spinae Plane (ESP) Block after mastectomy.</p> <p>Specific objectives:</p> <ol style="list-style-type: none"> To compare the total morphine consumption within 24 postoperative hours between patients receiving ESP block with bupivacaine and ESP block with saline for mastectomy at the Komfo Anokye Teaching Hospital, Kumasi, Ghana. To compare the numeric rating score at 2, 4, 6, 12 and 24 hours between patients receiving ESP block with bupivacaine and ESP block with saline for mastectomy at the Komfo Anokye Teaching Hospital, Kumasi, Ghana. To compare the time to the first request of rescue analgesia between patients receiving ESP block with bupivacaine and ESP block with saline for mastectomy at the Komfo Anokye Teaching Hospital, Kumasi, Ghana. To compare patients satisfaction within the 24-hour postoperative analgesia between patients receiving ESP block with bupivacaine and ESP block with saline for mastectomy at the Komfo Anokye Teaching Hospital, Kumasi, Ghana.
43	GBT-2104-133	Phase III	Inclacumab	27 th August, 2021	Professor Alex Osei-Akoto	Komfo Anokye Teaching Hospital (KATH)	Global Blood Therapeutics, Inc.	Application Pending Approval 7years 5 months	The primary objective of this study is to evaluate the long-term safety of every 12-week dosing of inclacumab in participants with sickle cell disease (SCD) who have completed a prior inclacumab clinical trial. Additional objectives are to evaluate the incidence of vaso-occlusive crises (VOCs), hospitalizations, missed work/school days, red blood cell (RBC) transfusions, and quality of life (QoL) with long-term use of inclacumab.
44	BEMPU	Phase II	BempuBracelet	2nd November, 2020	Mr. Prince Owusu	•Achimota General Hospital •Greater Accra Regional Hospital •Eastern Regional Hospital •Korle-Bu Teaching Hospital •Central Regional Hospital Princess Marie Luis Children Hospital	Center for learning and childhood development	Application Pending Approval	<p>To determine the accuracy of the bracelet in identifying hypothermia and evaluate its effect on Kangaroo Mother Care (KMC) practices and neonatal health outcomes in Ghana.</p> <p>To assess the acceptability of the bracelet in Health providers and caregivers of Low Birth Weight (LBW) infants by conducting qualitative in-depth interviews.</p> <p>Determine the accuracy of the BEMPU bracelet in classifying hypothermia in the clinical setting.</p> <p>Evaluate the impact of the bracelet</p>

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
45	DOLF_IDA ONCHO SAFETY GHANA	Phase II	1.Diethylcarbamazine Citrate I. P 100mg 2.Ivermectin (Stromectol®) 3mg) 3.Albendazole (Zentel™ 400mg)	22nd February 2019	Dr. Nicholas Opoku	University of Health and Allied Sciences	Washington University School of Medicine	Study ended Final report yet to be submitted 24 Months	<p>Programs for control of onchocerciasis through community directed treatment with ivermectin (IVM) as a form of Mass Drug Administration (MDA) have been in place for almost 30 years. IVM is effective for clearing Mf and it temporarily sterilizes adult female worms, but it is not a microfilaricide and does not kill adult worms. For that reason, MDA with IVM must be repeated for the reproductive life of the adult worms, which is 10-15 years. Thus, there is a widely recognized need for new, safe, short-course treatment drug(s) that can kill or permanently sterilize adult worms.</p> <p>This study aims to provide preliminary data on the safety of ivermectin + diethylcarbamazine + albendazole (IDA) treatment in persons with onchocerciasis when administered after pre-treatment with IVM to clear or greatly reduce microfilariae from the skin and eyes. Widespread use of IDA following IVM pretreatment (I/IDA) has the potential to greatly accelerate elimination of LF in African countries that are endemic for LF and onchocerciasis</p>
46	SMAART	Phase II	1.POLYCAP 2.USUAL CARE	9th February, 2018	Dr. Fred Stephen Sarfo	Komfo Anokye Teaching Hospital	Kwame Nkrumah University of Science and Technology	Study ended Final report yet to be submitted 19 months	<p>Africa (SSA), which when compared to stroke profiles in high-income countries (HIC) is characterized by a younger age of onset, higher case fatality rates, and more severe disability among survivors. Stroke survivors in SSA are especially at high risk for recurrent vascular events or death due to several factors including uncoordinated health systems, undiagnosed and under-controlled vascular risk factors, and lack of care affordability. Fixed-dose combination pills, known as "polypills", containing Aspirin, a statin and blood pressure (BP) lowering medication(s) may improve medication adherence and consequently reduce vascular risk as a cost-effective intervention among high risk patients including stroke survivors.</p> <p>This trial is to assess whether a polypill containing fixed doses of 3 antihypertensives, a statin and antiplatelet therapy taken once daily orally would result in carotid intimal thickness regression, improved adherence, and tolerability compared with 'usual care' group on separate individual secondary preventive medications among Ghanaian first time stroke survivors (male or female above</p>
47	LEDoxy	Phase II	1.Doxycycline (Remycin®100mg 2.Placebo 3.Standard MDA Treatment	12th July, 2017	Prof. Alexander Yaw Debrah	1.Kumasi Centre for Collaborative Research (KCCR), Kwame Nkrumah University of Science and Technology (KNUST) 2.War Memorial Hospital, Navrongo	Kumasi Center For Collaborative Research (KCCR)	Study ended Final report submitted 40 months	<p>The previously demonstrated effect of doxycycline in reversing or stopping the progression of lymphedema of patients with stage 1-3, irrespective of their filarial infections being active or not, provides an opportunity to include the drug as a new tool in lymphatic filariasis (LF) morbidity management programs. However, before recommendations can be made regarding the frequency of its usage or alternate dosing patterns more trials need to be conducted. This multi-national trial is to show efficacy of a lower dosage of doxycycline and to confirm finding in patients with stages 1-3 lymphedema irrespective of active LF infection as well as in people with higher grades of lymphedema.</p> <p>The purpose of the study is to establish that Doxycycline can improve filarial lymphedema in healthy adolescents or adults (14 – 65 years)</p>
48	FALCON	Phase III	1.ChlorPrep™ stick 2.Videne® Antiseptic Solution 3.Triclosan Coated PDS and/or Vicryl sutures 4.Non-triclosan coated PDS and/or Vicryl sutures	10th April, 2019	Prof. Stephen Tabiri	Tamale Teaching Hospital	The University of Birmingham	Study ended Final report submitted 24 Months	<p>Improving surgical outcomes is a global health priority. Recent World Health Organisation (WHO) guidelines made 29 recommendations for intraoperative and postoperative measures to prevent SSI, including global perspectives relevant to LMICs., none of the evidence for the recommendations used was derived from resource limited settings, leading to uncertainty about implementation of measures in these settings. A randomised trial that has the potential to evaluate multiple interventions has particular value in this setting, and can establish a high quality evidence base that will inform guidance, and influence revisions to the WHO Surgical Safety Checklist</p> <p>This study assesses whether either (1) 2% alcoholic chlorhexidine versus 10% povidone-iodine for skin preparation, or (2) triclosan-coated suture versus non-coated suture for fascial closure, can reduce surgical site infection at 30-days post-surgery for each of (1) clean-contaminated and (2) contaminated/dirty surgery</p>

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
49	KNC 19 (NIBIMA)	Phase IIb	1.Nibima 2.WHO standard treatment for COVID-19	11th September 2020	Prof. Ellis Owusu-Dabo	Komfo Anokye Teaching Hospital	KNUST Office of Grants and Research	Study ended Yet to submit Final report From 3 months to 7 months	The purpose of this trial is to evaluate the: •Efficacy of Nibima in reducing >50% Covid-19 viral load per patient within 14 days of therapy. Evaluate the efficacy of Nibima in increasing the anti-inflammatory and interferon alpha/beta profiles of >50% of the Covid-19 patients within 14 days.
50	MULTIMAL	Phase II	Pyronaridine (Pyramax) 2.Atovaquone Proguanil (Malarone) 3.Clindamycin 4.Foscidomysin5	27th July 2020	PI(s) Dr. Oumou Maiga (KCCR)	St. Francis Xavier Hospital Assin Fosu, Ghana. Gabon	Tropical Medicine, Bernhard Nocht Institute for Tropical Medicine (BNITM)	Study ended Yet to submit Final report 7 months	outcome of this consideration was that the specific multi-therapeutic ACT combinations, discussed below, were decided on based on the following aspects: efficacy, potential for drug interactions, modes-of-action, half-life of the individual drugs, parasitological stages the drug acts on, dosing, availability of a paediatric formulation and cost.The two drug combinations envisaged to investigate during this study address two particular aspects of treatment of uncomplicated malaria in the sub-Saharan African region. Firstly,
51	STAR TRIAL	Phase IV	1.Paracetamol 2.Morphine	7th May 2021	Dr. Frank Enoch Gyamfi	Komfo Anokye Teaching Hospital, Kumasi	Dr. Frank Enoch Gyamfi	Study ended yet to submit Final report 10 months	with bimodal administration of i.m. morphine and i.v. paracetamol in managing postoperative pain in emergency abdominal surgery. To assess the response of patients to i.m. morphine in pain management after emergency abdominal surgery. To assess the response of patients to a combination of i.v. paracetamol and i.m. morphine in managing pain after emergency abdominal surgery. To determine the association between the administered analgesic and length of hospital stay. To determine the association
52	DIABETIC FOOT SELF CARE		1.Foot Selfcare Training and Education Plus usual care 2. Usual care.	28th October 2021	Dr. Joseph N. Suglo	Diabetes Clinic, Komfo Anokye Teaching Hospital (KATH) – Ghana	King's College London (KCL)	Study ended Final report in E3 format yet to be submitted, 7 months	The primary aim of this research is to evaluate the reasonability of conducting a randomised controlled trial to investigate the effectiveness of a hands-on skills training and education on foot self-care programme for persons with diabetes and their family caregivers in Ghana. The research question is 'can the provision of a family-oriented foot self-care skills training and education intervention improve foot care behaviour, foot care self- efficacy, knowledge of
53	CHEETAH	Pilot	1.Sterile Gloves 2.Sterile Surgical Instrument	1st June 2020	Professor Stephen Tabiri	•Cape Coast Teaching Hospital •Effiah Nkwanta Regional Hospital •Holy Family Hospital – Berekum •Holy Family Hospital – Techiman •KATH	Birmingham Clinical Trials Unit, University of Birmingham	Study ended Final report submitted. 24 Months	To purpose of this study is to assess whether the practice of using separate, sterile gloves and instruments to close wounds at the end of surgery can reduce surgical site infection at 30-days post-surgery for patients undergoing clean-contaminated, contaminated or dirty abdominal surgery, compared to current routine hospital practice.
54	KAE609	Phase II	1.KAE609 2.COARTEM TABLETS	1st September 2019	Dr. Abraham Rexford Oduro	1.Navrongo Health Center 2.Kintampo Health Research Centre	Novartis Pharma AG, Switzerland	Study ended; Final report submitted 14months	KAE609 will be evaluated primarily for hepatic safety of single and multiple doses in sequential cohorts with increasing doses. This study aims to determine the maximum safe dose of the investigational drug KAE609 in Adult patients with acute, uncomplicated Plasmodium falciparum malaria infection..

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
55	Saving Brains Navrongo	I	1.Small Quantity Lipid-based Nutrient Supplement for Pregnant and Lactating mothers (SQLNS P&L) 2. Enhanced Small Quantity Lipid-based Nutrient Supplement for Pregnant and Lactating mothers (eSQLNS P&L)	7th February 2019	Dr. Engelbert A. Nonterah	Navrongo Health Research Centre	Nutriset, SAS	Study ended; Final report yet to be submitted 6 months	Malnutrition continues to be a global problem. Globally 156 milion children less than 5 years are stunted, 50 million wasted, while simultaneously 42 million are overweight reflecting the double burden of malnutrition. Prevalence of malnutrition varies by region and country with Asia and Africa being the worst affected regions. This study is to ssess the acceptability and adherence to nutrient supplementation for 6 weeks among pregnant and lactating women and 6 monh old infants post weaning
56	SAVING BRAINS KUMASI	I	1.Small Quantity Lipid-based Nutrient Supplement for Pregnant and	1st November 2017	Prof. Jacob Plange-Rhule	1. Ifo Government Hospital 2.Suntreso Government Hospital	KNUST/Nutriset SAS	Study ended 6months	Malnutrition continues to be a global problem. Globally 156 milion children less than 5 years are stunted, 50 million wasted, while simultaneously 42 million are overweight reflecting the double burden of malnutrition. Prevalence of malnutrition varies by region and country with Asia and Africa being the worst affected regions. This study is to ssess the acceptability and adherence to nutrient supplementation
57	ALB_IVM	III	1. Ivermectin 2. Albendazole	1st April 2014	Dr. Nicholas Opoku	Onchocerciasis Chemotherapy Research Centre Government Hospital.	Case Western Reserve University School of Medicine, 10900 Euclid Ave Cleveland	Study ended; Final report submitted 38 months	To address whether IVM plus ALB given twice per year will be superior over annual treatment or IVM given biannually
58	MAL 055	III	RTS,S/AS01E	1st October 2008	1. Prof. E. Tsiri Agbenyaga 2. Prof. Seth Owusu Agyei 3. Dr. Kwaku Poku Asante	1. Malaria Research Centre, Agogo. 2. Kintampo Health Research Centre	GlaxoSmithKline Biologicals	Study ended; Final report submitted 60 months	This Phase III study of GSK Biologicals candidate malaria vaccine RTS,S/AS01E has been designed to address the key safety and efficacy information required for vaccine licensure. In addition, other disease endpoints that allow the evaluation of the full public health impact and cost effectiveness of vaccine implementation are included. Co-primary objectives will investigate the efficacy against clinical disease in children from 5-17 months of age at first dose and the efficacy in infants 6-12 weeks of age who receive the vaccine in co-administration with EPI antigens
59	MMS	III	1.Multiple micronutrient supplement 2.Iron + folic acid tablets	2nd October 2012	Prof. Tsiri Agbenyaga	1. Dabokoma Collaborative Community Development Project 2. C/O Komfo Anokye	Kirk Humanitarian	Study Ended; yet to submit report 48 months	
60	PRENABELT		1.Prenabelt™ 2. Sham prenabelt™ 3.Body Position Sensor	21st April 2015	Dr. Jerry Coleman	Korle-Bu Teaching Hospital, Accra – Korle Bu	Global Innovations for Reproductive Health and Life, USA	Study ended; Final report submitted 7 months	The purpose of this study is to determine the effect of the PrenaBelt on birth-weight and assess the feasibility of introducing it to Ghanaian third-trimester pregnant women in their home setting via an antenatal care clinic and local health-care staff. Data from this study will be used in effect size calculations for the design of a large-scale, epidemiological study targeted at reducing LBW and SB in Ghana and globally.
61	CPAP	Phase III	1.DeVilbiss IntelliPAP CPAP machine (Model DV5 Series) 2. Hudson RCI nasal cannulas	14th May 2013	1. Dr. Harry Tagbor 2. Dr. Frank Baiden 3. Dr. Damien Punguyire 4. Dr. Kwadwo Nyarko Jectey	1. Mampong Government Hospital, Mampong 2. Kintampo Municipal Hospital, Kintampo	(GE) Foundation's Systems Improvement at District Hospitals and Regional Training of	Study ended; yet to submit report in required format. 36 months	Evaluating the impact of using continuous positive airway pressure (CPAP) on mortality among children admitted into emergencies wards. an interventional trial to determine if CPAP reduces morality in children 1 month to 5 years of age with acute respiratory distress
62	AIMS	Phase III	1.Mirasol system for whole blood 2.Standard fresh whole blood	9th July 2013	Dr. Shirley Owusu-Ofori	Komfo Anokye Teaching Hospital	Terumo BCT Europe N.V.	Study ended; Final report submitted 6 months	The objective of this study was to evaluate the efficacy of Mirasol-treated fresh whole blood (WB) to prevent transfusion-transmitted malaria (TTM) by comparing the incidence of TTM between subjects receiving Mirasol-treated fresh WB and subjects receiving standard (untreated) fresh WB.
63		II	Meningococcal A Conjugate Vaccine	26th June 2007	Dr. Patrick Ansah	Navrongo Health Research Centre	SIIL PATH	Study ended; Final report submitted 54 months	To compare the immunogenicity at 28 days after vaccination of range dosages - 10, 5, and 2.5 µg of the PsA-TT vaccine, when administered to infants in a two-dose schedule at 14 weeks (window 14 to 18 weeks of age) and 9 months of age (window 9 to 12 months of age) concomitantly with EPI vaccines (Groups 1A vs.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
64	NON-INVASIVE HAEM DEVICE	III	1. Pronto & pronto-7 pulse co-oximeter pulse co-oximeter 2. Hemocue 201+3. Abx penra 60 hematology analyzer	9th April 2013	Dr. Sam Newton	Kintampo Health Research Centre, Kintampo	PATH	Study Ended Final report submitted 2 months	
65	ROTARIX	III	Rotarix™	6th February 2012	Prof. George Armah	Navrongo Health Research Centre	PATH	Study Ended 7 months Final Report submitted	To show the superiority of live, oral Rotarix vaccine administered at 6, 10, and 14 weeks of age versus live, oral Rotarix vaccine administered at 6 and 10 weeks of age in terms of serum rotavirus immunoglobulin A (IgA) seroconversion as the marker of vaccine-induced immunogenicity
66	ARTIMIST	III	ArTiMist	22nd October 2010	Dr. Patrick Ansah	Navrongo Health Research Centre	ProtoPharma Limited	Study Ended Final report submitted 5 months	The primary objective of this study was to demonstrate the superiority of ArTiMist™ over intravenous (iv) quinine in establishing parasite success (reduction of parasite counts by ≥ 90% within 24 hours) in children with severe or complicated falciparum malaria, or children with uncomplicated malaria with gastrointestinal complications.
67		III	Gardasil	1st November 2010	Dr. Nana Akosua Ansah	Navrongo Health Research Centre	Merck, Sharp and Dohme Corporation	Study Ended Final report submitted 20 months	To estimate the percentage of subjects who seroconvert to each of HPV 6, 11, 16, and 18 at Month 7 (4 weeks Postdose 3). To evaluate the safety and tolerability of GARDASIL in females 9 to 26 years of age in SubSaharan Africa. Secondary: To estimate Month 7 anti-HPV 6, 11, 16, and 18 geometric mean titers (GMTs) in vaccinated subjects
68	SMAC	III	1. Intravenous Artesunate 2. Intramuscular Artesunate	1st January 2013	Prof. Tsiri Agbenyega	Komfo Anokye Teaching Hospital, Kumasi	University Medical Centre Tubingen	Study Ended 15 months	
69	OXYTOCIN	III	1. Oxytocin in uniject™ 10 iu	12th May 2010	Dr. Sam Newton	Kintampo Health Research Centre	PATH	Study Ended Final report submitted 12 months	
70	AMARYL M	IV	Amaryl m oral tablets	16th October 2009	Dr. Frank Umeh	Korle-Bu Teaching Hospital	Sanofi Aventis	Study Ended 6 months	
71	MOXIDECTIN-IVERMECTIN	III	1. Moxidectin 2. Ivermectin	1st February 2004	Dr. Nicholas Opoku	Onchocerciasis Chemotherapy Research Centre Government Hospital.	1. Wyeth Research Division of Wyeth Pharmaceuticals Inc. 2. Product Development and Evaluation unit TDR	Study Ended Report submitted 25 months + (12 months ext.)	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
72	MOXIDECTIN	Phase II	Moxidectin 2mg Tablets	1st February 2004	Dr. Kwabla Awadzi	Onchocerciasis Chemotherapy Research Centre Government Hospital	1. Wyeth Research Division of Wyeth Pharmaceuticals	Study Ended Ended 60 months	
73	EBA	I	(EBA-175 RII-NG) malaria vaccine	1st March 2009	Prof. Kwadwo Ansah Koram	Noguchi Memorial Institute of Medical Research	Division of Microbiology and Infectious Diseases (DMID) National Institute of Allergy and Infectious	Study Ended Final report submitted 18 months	
74	IPT & SP	III	Sulfadoxine-pyrimethamine	1st May 2008	Dr. Abraham Hodgson	Health Facilities in the Kassena Nankana, Navrongo Health Research Centre	London School of Hygiene and Tropical Medicine	Study Ended 32 months	
75	IRON FORTIFICATION III		1. Sprinkles vitamin 2. mineral food supplement	1st July 2009	Prof. Seth Owusu Agyei	Kintampo Health Research Centre	National Institutes of Health	Study Ended 12 months	
76	ROTASHIELD	III	RRV-TV Vaccine (rotashield)	1st August 2009	1. Prof. George E. Armah 2. Prof. Fred N. Binka 3. Dr. Abraham Hodgson	1. War Memorial Hospital, Navrongo 2. Bongo Hospital	International Medica Foundation	Study Ended 16 months	
77	PLUS CHLOROQUINE PHOSPHATE	III	1. Azithromycin 2. Chloroquine Phosphate 3. Artemether-Lumefantrine	1st October 2007	Dr. Patrick Ansah	Navrongo Health Research Centre	Pfizer Laboratories Incorporated, Pfizer Global Research and Development	Study Ended Final report submitted 8 months	
78	CRASH-2	I	1. Tranexamic acid 2. Placebo	1st August 2007	Prof. J. C. B. Dakubo	Korle-Bu Teaching Hospital	London School of Hygiene & Tropical Medicine	Study Ended, Lancet publication submitted 24 months	
79	PYRONARIDINE ARTESUNATE VRS COARTEM	III	1. Pyronaridine Artesunate Tablet (PYRAMAX) 2. Artemether-Lumefantrine (COARTEM)	1st March 2007	Dr. G. Bedu-Adoo	Komfo Anokye Teaching Hospital	Medicines For Malaria Venture, Switzerland	Study Ended 3 months	
80	MAL 050	III	RTSS, AS10E Vaccine		Prof. Seth Owusu Adjei	Kintampo Health Research Centre	GlaxoSmithKline R&D	Study Ended 17 months	
81	PfCSP_MVACS_MALARIA	I	PfCSP DNA VACCINE (VCL-2510)	1st August 2005	Prof. Kwadwo A Koram	Tetteh Quarshie Memorial Hospital	Division of Microbiology and Infectious Diseases (DMID) National Institute of Allergy and Infectious Diseases (NIAID)	Study Ended 18 months	
82	ROTATEQ	III	Rotateq	1st September 2007	Prof. George E. Armah	Navrongo Health Research Centre	1. Merck & Co. 2. PATH	Study Ended Final report published in Lancet 18 months	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
83	MEFLOQCHLOA ZITH	III	1. Mefloquine 2. Chloroquine 3. Azythromycin	4th August 2004	Dr. Abraham Hodgson	Navrongo Health Research Centre	Pfizer Inc.	Study Ended Final report submitted 12 months	
84	MAL 047	II	1.RTS,S/AS02D 2.RTS,S/AS01E		Prof. Seth Owusu Adjei, Dr. Kwaku Poku Asante	Kintampo Health Research Centre	GlaxoSmithKline R&D	Study Ended 19 months	
85	CDA	III	1.Chorproguanil-Dapstone-Artesunate (CDA) 2.Artemether-Lumefantrine	19th July 2006	Prof. Seth Owusu Agyei Dr. Kwaku Poku Asante	Kintampo Health Research Centre	GlaxoSmithKline R & D	Study Ended 12 months	
86	CDA2	III	1.Chorproguanil-Dapstone-Artesunate (CDA) 2.Artemether-Lumefantrine	27,June 2006	Prof. Tsiri Agbenyega	Department of Physiology, School of Medical Sciences, KNUST	GlaxoSmithKline R & D	Study Ended 12 months	
87	NOVASIL	II	NovaSIL		Prof. David Ofori Agyei Dr. Nii- Ayi Ankrah	Ejura Sekyedumasi District, Ashanti Region	United States Agency for International Development (USAID)	Study Ended 9 months	
88	TENOFOVIR	II	Tenofovir Disoproxyl Fumarate (TDF)	1st February 2004	Dr. Edith Clarke	Ghana Health Service	Family Health International	Study Ended 20 months	
89	SAVVY	II	SAVVY (Microbicide)	1st February 2004	Dr. William Ampofo Dr. Baafuor Kofi Opoku	1. Noguchi Memorial Institution for Medical Research. 2. Komfo Anokye Teaching Hospital.	Family Health International	Study Ended 32 months	
90	MAL 063	III	RTS,S/AS01E	15th April 2011	Prof. E. Tsiri Agbenyaga	Malaria Research Centre, Agogo.	Malaria Research Centre, Agogo	Study Ended Final report submitted 52 months	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
91	PREGACT	III	1. Eurartesim oral tablets 2. Farmanguinhos artesunate+mefloquine fixed combination oral tablets 3. Coarsucum oral tablets		1.Dr. Harry Tagbor 2.Dr. Henry Opare Addo	1.Ejisu Government Hospital, Ejisu 2. Juaben Government Hospital, Juaben	Prince Leopold Institute of Tropical Medicine	Study Ended 60 months	
92	ALBIVIM K'SI	III	1. Ivermectin 2. Albendazole	10th November 2015	Prof. Alexander Yaw Debrah	Kumasi Centre for Collaborative Research in Tropical Medicine	University Hospitals Case medical Center	Study Ended, Yet to submit final report 4 years and 2 months	
93	RIFAMPIN VS ISONIAZID	III	1. Isoniazid 2. Rifampin	2nd March 2011	Dr. Joseph Baah Obeng	Komfo Anokye Teaching Hospital Chest Clinic, Kumasi	Canadian Institute of Health Research	Study Ended 60 months	
94	NOGUCHI FILARIASIS *		1. Alere filariasis test strip 2. Sd bioline lymphatic filariasis IgG4 3.Sd	7th June 2017	Prof. Daniel A. Boakye Dr. Nana – Kwadwo Biritwum	Noguchi Memorial Institute For Medical Research	World Health Organization - TDR	Study Ended Final report submitted 10 months	Development of a plan of action for strengthening LF elimination in Ghana, and where appropriate, a plan of action for integrating LF and onchocerciasis elimination efforts, to be proposed to the GHS decision makers.
95	ZIV AFFLIBERCEPT	I	1. Ziv-afibercept (ZALTRAP)	30th January 2017	Braimah Imoro Zeba	Keena Unit, Eye Centre, Korle-Bu, Teaching Hospital, Korle-Bu, Accra	Same as PI	Study ended Final report submitted 5 months	To evaluate the safety of 1.25mg and 2mg ziv-afibercept in Ghanaian population with retinal vascular diseases. To determine the safety of intravitreal injections of ziv-afibercept at 4 and 12 weeks in a Ghanaian population. To measure the visual outcome of treatment with 1.25mg and 2mg ziv-afibercept resulting in altered (sickle- shaped) red-blood cells. A vaso-occlusive crisis (VOC) is a severe, acute painful episode that occurs when sickle-shaped red blood cells obstruct the microcirculation and restrict blood flow to an organ or tissue, resulting in ischaemia, necrosis and organ damage. There is a high unmet need for treatment options in SCD and there is a data that platelet inhibition has the
96	HESTIA3	Phase III	1. Ticagrelor 2. Placebo	1st August, 2018	2. Dr Patrick Ansa 3. Dr. Catherine Segbefia 4. Dr Kokou Hefoume Amegan-Aho	Teaching Hospital, Department of Child Health 2. Navrongo Health Research Centre	AstraZeneca AB	Study Ended. Final Report submitted 29 Months	The lack of access to reliable tests for proteinuria measurement in low resource care settings, particularly at the periphery, remains a critical gap in the accurate identification of women at high risk for Pre-Eclampsia. In Low Resource Settings, a protein-only measurement via a urine dipstick is the most widely used proteinuria test due in part to its low complexity and low cost. However, the clinical utility of the protein-only dipstick is limited. Test results can be unreliable, as the test cannot adjust for daily fluctuation of body hydration. This leads to protein measurements that are either too low or too high due to the level of urine dilution. More accurate
97	PRCR DIPSTICK	Phase II	1. Creatinine Dipstick 2. Urinalysis Reagent Strips 3. Quantitative Spectrophotometric Method	16th February, 2018	Dr. Sam Newton	Kintampo Health Research Center	Program For Appropriate Technology In Health (PATH)	Study Ended. Final Report Submitted 19 months	In sub-Saharan Africa, most of the Expanded Program on Immunization (EPI) vaccines are given in early infancy while measles, rubella and yellow fever (YF) vaccines are given at 9 months of age. Between the first EPI vaccines and the measles, rubella and YF vaccines, children receive Vitamin A supplementation at 6 months of age. To limit the number of clinic visits for young children and to optimize vaccine implementation a schedule (0, 1.5, 3-month) is proposed. There are however no data of the anti-circumsporozoite protein of Plasmodium falciparum (anti-CS) immune response induced by RTS,S/AS01E when given in combination with measles, rubella and YF, in a 0, 1.5, 3-month schedule starting at an older age (5-17 months). This study intends to demonstrate that anti-CS immune response of the candidate malaria vaccine RTS,S/AS01E is not inferior when RTS,S/AS01E is administered at 6, 7.5 and 9 months of age with the third
98	MAL 073	Phase IIIb	1. RTS,S/AS01E 2. MR-VAC™ 3. STAMARIL4. VITAMIN A	11th December 2015	1. Prof. Tsiri Agbenyega Prof. Seth Owusu Adjei	1. Malaria Research Center, Agogo 2. Kintampo Health Research Centre Hospital	GlaxoSmithKline Pharmaceuticals	Study Ended Final Report submitted 43 months 16 days	
99	CEPHEID XPRT HIV-1	PILOT	Xpert HIV-1 VL XC Test Assay for detecting HIV-1 RNA in human plasma.	6th June 2019	Prof. Jacob Plange-Rhule	Atua Government Hospital Akosombo Hospital	CEPHEID	Study Ended Final Report yet to be submitted 6 Months	The Xpert® HIV-1 Viral Load XC test is an in vitro reverse transcriptase polymerase chain reaction (RT-PCR) assay for the quantification of Human Immunodeficiency Virus type 1 (HIV-1) RNA in human plasma using the automated GeneXpert® Instrument Systems. It is intended for use as an aid in the diagnosis of HIV-1 infection, as a confirmation of HIV-1 infection, and as an aid in clinical management of patients infected with HIV-1.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
100	INNOVATE	Phase III/II	1. Inn0-4800 2. Placebo		Susan Adu-Amankwah	Noguchi Memorial Institute for Medical Research	Inovio Pharmaceuticals, Inc	Study Closed/withdrawn by Sponsor months 24	1. Evaluate the cellular and humoral immune response to INO-4800 administered by ID injection followed immediately by electroporation EP 2. Evaluate the efficacy of INO-4800 in the prevention of COVID-19 disease in subjects who are SARS-CoV-2 negative at baseline
101	LIVZON	Phase III	1. SARS-CoV-2 fusion protein vaccine (code: V-0) 2. Placebo	2nd August 2021	1. Dr Seyram Kaali 2. Dr. Nana Akosua Ansah	1. Kintampo Health Research Centre 2. Navrongo Health Research Centre	Livzon Mabpharm Inc. Institution Pharmaceutical company	Study Closed by Sponsor before commencement. No recruitment was done. 20 months	Efficacy: To evaluate the efficacy of the recombinant SARS-CoV-2 fusion protein vaccine (V-01) for the prevention of symptomatic RT PCR positive COVID-19 (mild or above severity) starting from at least 14 days (≥15 days) after full-course immunization (completing all vaccinations) Safety: To evaluate the incidence of adverse events (AEs) of recombinant SARS-CoV-2 fusion protein vaccine (V-01) from the first vaccination to 28 days after full-course immunization
102	COVID 19 INTRANASAL SPRAY	Phase III	1. Influenza Virus Vector COVID-19 Vaccine for Intranasal Spray 2. Placebo	19th October 2021	Dr. Seyram Kaali	1. KHRC 2. NHRC 3. KCCR 4. Dodowa Health Research Center 5. Ghana Infectious Disease Center 6. KBTH	Beijing Wantai Biological Pharmacy Enterprise Co, Ltd	Study Closed by Sponsor before commencement. No recruitment was done. 20 months	1. To evaluate the protective efficacy of DelNS1-2019-nCoV-RBD-OPT1 for preventing virologically confirmed (RT-PCR positive) symptomatic COVID-19. 2. To evaluate the safety of DelNS1-2019-nCoV-RBD OPT1.
103	STEADFAST	Phase II	CRIZANLIZUMAB	15th February, 2021	Dr. Yvonne Dei Adomako	•Ghana Institute of Clinical Genetics Korlebu •Sickle cell office Directorate Child(KATH)	Novartis Pharma	Study closed by sponsor before commenced 21 Months	The purpose of this study is to explore the effect of P-selectin inhibition with crizanlizumab on renal function in SCD patients with CKD who are receiving standard of care for SCD-related CKD, have Grade A2-A3 albuminuria and Stage 1-3a CKD, and are at risk for rapid decline in their eGFR.
104	ESM UBT *		Uterine balloon tamponade	17th February, 2014	Dr. Ivy Frances Osei	Field Work	Bill and Melinda Gates Foundation, USA	Study not conducted; Funds from Sponsor withdrawn before initiation 8months	
105	FERROQUINE	II	1. Ferroquine 2. Amodiaquine 3. Artesunate	Apr-08	Dr. Josephine C. Ocran Prof. Kwadwo Ansah Koram	Noguchi Memorial Institute of Medical Research	Sanofi-Aventis Recherche And Development	Study Closed by Sponsor. No recruitment was done. 13Conths	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
106	HOPE SCD	III	GBT440 300mg	May-17	1. Dr. Yvonne Dei Adomakoh 2. Dr. Vivian Paintsil	1. Center for Clinical Genetics, Korle-Bu Teaching Hospital 2. Paediatric Sickle cell clinic, Komfo Anokye Teaching Hospital	Global Blood Therapeutics Inc. 400 East Jamie Court, Suite 101 South San Francisco, CA 94080, USA	Group 1 and 2 under current protocol completed (none recruited in Ghana); yet to start Main Population Study (Group 3) 17 months	The primary objective is to assess the efficacy of GBT440 in adolescents and adults with SCD as measured by improvement in anemia
107	VERO CELL COVID 19 TRIAL	Phase III	Inactivated (Vero Cell)	10th February 2022	1. Dr Alberta Amu Dr. Patrick Ansah	1. Dodowa Health Research Center 2. Navrongo Health Research Center	Institute of Medical Biology Chinese Academy of	Application Withdrawn, 18 Months	1. To evaluate the efficacy of SARS-CoV-2 Vaccine, Inactivated (Vero Cell) against symptomatic and laboratory-confirmed (RT PCR method) COVID-19 cases 2. To evaluate the solicited AEs within 7 days after each dose. 3. To evaluate the efficacy of SARS-CoV-2 Vaccine, Inactivated (Vero Cell) after at least
108	MEBENDAZOLE	IV	Menbendazole	Sep-17	Prof Michael David Wilson	Kintampo Health Research Centre	Program For Appropriate Technology In Health (PATH)	Application Withdrawn N/A	Soil-transmitted helminth (STH) infections are considered among the most pressing of global health problems, thought to parasitize some 2 billion people worldwide.[] The most recent estimates suggest that between 600 and 800 million people are infected with one or several of the common soil-transmitted helminths (STHs), which are Ascaris lumbricoides, Trichuris trichiura, and hookworm.[] Infection prevalence, incidence, and disease burden are particularly high in tropical and subtropical areas that are already burdened with poor living conditions, over-population, and inadequate sanitation, including some areas of sub-Saharan Africa, Asia, and Latin America.[1, ,] While adults represent a significant percentage of the infected population, it is children who are the most vulnerable
109	EBOLA Z	II	chimpanzee adenovirus Type 3 – vectored Ebola Zaire vaccine (ChAd3-EBO-Z)	Jan-15	1. Dr. Kwaku Poku Asante 2. Prof. Kwadwo A Koram	1. Kintampo Health Research Centre 2. OCRC, Hohoe	GlaxoSmithKline Biologicals	Application withdrawn N/A	
110	EBOLA Z (Paediatric)	II	chimpanzee adenovirus Type 3 – vectored Ebola Zaire vaccine (ChAd3-EBO-Z)	21st August 2015	Dr. Kwaku Poku Asante	OCRC, Hohoe	Glaxosmithkline Biologicals, Rue De L'institut, 89 – 1330 Rixensart, Belgium	Application withdrawn N/A	
111	ZEBOV	I	expressing the glycoprotein of the ebola virus mayinga variant [Ad26.ZEBOV	7th January 2015	Professor Fred Binka	OCRC, Hohoe	B.V, Represented by Janssen Pharmaceutica (Pty) Ltd	Approved but sponsor withdrew conduct N/A	
112	ZEBOV 2	II	expressing the glycoprotein of the ebola virus mayinga variant [Ad26.ZEBOV 2. Modified vaccinia ankara – bavarian nordic	6th April 2015	Professor Fred Binka	OCRC, Hohoe	Cruell Holland B.V, Represented by Janssen Pharmaceutica (Pty) Ltd	Application withdrawn N/A	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
113	HYDRANON	I	Hydranon solution	1st March 2008	Prof. David Ofori-Adjei	Noguchi Memorial Institute For Medical Research	General Resonance Technology 11lc	Application Withdrawn N/A	
114	SALIF,	IIIb	1.TDF/FTC/RPV 2.TDF/FTC/EFV	4th September 2013	1. Dr. Isaac Osei 2. Dr. Samuel Abara 3. Dr. Fred Adomako –	Navrongo Health Research Centre Upper East Regional Hospital	Janssen-Cilag International NV (Sponsor) represented by Clinical	Application Withdrawn N/A	
115	NOGUCHI SCD	Ib	NVX-508	1st May 2017	Amma Twumwaa Owusu Ansah	1. Noguchi Memorial Institute For Medical Research 2. College of Health Sciences 3. University of Ghana	University of Pittsburg, Representative: Amma Owusu-Ansah, MD	Application Withdrawn N/A	
116	PRCR SPOT	Phase II	PRCR Spot	15th March 2021	Dr. Hannah Brown Amoakoh	Ridge Hospital, Korlebu Teaching Hospital, Koforidua Regional Hospital	Emily Stephanie Zobrist, PATH, 2201 Westlake Avenue, Seattle, WA 98121, USA	Application Withdrawn by Sponsor	To address the gap in proteinuria measurement solutions, LifeAssay Diagnostics (LAD) has developed and commercialized a low-cost PrCr urine dipstick that has shown good laboratory and clinical performance and high usability within antenatal care (ANC) settings in previous studies. There is a need for further evidence on the clinical utility and operational fit of the LAD Test-it™ PrCr test to inform policy recommendation for its use in Ghana and other LMIC settings.
117	SAR97276A_SANOFI	II	SAR97276A	1st October, 2008	Prof. Seth Owusu-Agyei	Navrongo Health Research Centre	Sanofi Aventis Recherche & Developpement	Application Withdrawn by Sponsor before approval	
118	TENOFOVEK BEI	Bioequivalence	1. tenofovir (tenofovir) 300mg film coated tablets 2. Viread (tenofovir) 300mg	11th September 2015	1. Prof. Seth Owusu Agyei 2. Dr. Kwaku Poku Asante	Kintampo Health Research Centre	Danadams Pharmaceuticals Industry Limited, Accra-Ghana	Application closed by FDA since Sponsor failed to start study 3 years after approval.	
119	ELDON CARD NYN		1. Eldon card 2. Standard laboratory method	10th November 2015	Prof. Samuel Ameny Obed	Korle Bu Teaching Hospital, Accra.	Center for Global Child Health, Hospital for sick Children.	Incomplete CTA; Application closed by FDA. N/A	
120	AX-100 HIVI		1. AX-100 Immun 2. AX-100 ImmunPlus	9th december 2014	Dr. Kwaku Poku Asante	Kintampo Health Research Centre	Neopharmacie Limited, Germany	Incomplete CTA; Application closed by FDA. N/A	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
121	4P	III	Polypil	9th August 2013	1. Dr. Emmanuel Kwabla Srofenyoh 2. Dr. Patrick Frimpong	Ridge Hospital Accra La General Hospital	Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, The Netherlands	Incomplete CTA; Application closed by FDA. N/A	
122	INVACT	III	Artemisinin	13th may 2016	Prof. Kwadwo Ansah Koram	Noguchi Memorial Institute For Medical Research	Global Emerging Infections Surveillance and Response System of the US Armed Forces Health Surveillance Center	Incomplete CTA; Application closed by FDA. N/A	
123	INSUGENIV		Insugen	17th december 2013	N/A	Korle-Bu Teaching Hospital	BIOCON LTD	Incomplete CTA; Application closed by FDA. N/A	
124	AIM-LVRNA009	Phase II/III	1. SARS-CoV-2 mRNA vaccine (LVR) 2. Saline Placebo	21st June 2022	Dr. Patrick Odum Ansah	1. Navrongo Health Research Centre 2. Kumasi Centre for Collaborative Research 3. Dodowa Health Research Centre 4. Kintampo Health Research Centre 5. Ghana Infectious Disease Centre 6. Korle Bu Teaching Hospital (KBTH)	AIM Vaccine Co. Ltd.	Not Approved, 17-24 months.	Primary efficacy objective: To evaluate the protective efficacy of LVRNA009 (50 µg) in the prevention of first episodes of virologically-confirmed symptomatic cases of COVID-19 of any severity occurring from 14 days after 2nd dose in the initial set of vaccination in SARS-CoV-2 naive participants
125	MYCOPIROX_LA GRAY	III	Mycopirox Vaginal cream	15th June 2010	Dr. Luitgard Darko		Lagray Chemical Company, Ltd.	Not Approved N/A	
126	MoRiOn	II	1. Rifampentine (Pritin®) 2. Moxifloxacin (Avelox®) 3. Doxycycline	28th April, 2017	Prof. Alexander Yaw Debrah	1. Enchi Government Hospital 2. Communities of Aowin/Suaman District W/R	Kumasi Centre for Collaborative Research in Tropical Medicine	Study terminated by sponsor Yet to submit Final report 15 months	Onchocerciasis is caused by the parasite Onchocerca volvulus. More than 37 million people are estimated to be infected with O. Volvulus worldwide. The current therapeutic strategy relies on annual mass drug administration (MDA) based on the drug donation program for Ivermectin. Ivermectin is mainly microfilaricidal and after a few months female worms resume MF production levels high enough for transmission. Therefore, safe microfilaricidal drugs are needed to reach the goal of elimination. The study aims to show efficacy (Wolbachia depletion) of combination Rifampentine plus Moxifloxacin using immunohistology compared to no treatment and treatment with Doxycycline.

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
127	COVID MOUTHWASH	Phase III	1.Corsodyl Mouthwash 2.Wokadine mouthwash 3.Hydrogen Peroxide mouthwas	6th September 2021	Dr. George Boateng Kyei	Noguchi Memorial Institute for Medical Research	Dr. George Boateng Kyei	Study terminated by sponsor Yet to submit Final report 1 year 6 months	To investigate how long it takes for SARS-CoV-2 asymptomatic or presymptomatic persons to shed viable virus. It also seeks to evaluate among these patients the effect of a one-time mouth rinse on the detectable viral load of SARS-CoV-2 and to determine how long it takes for SARS-CoV-2 viral load to remain low after using the mouth rinse.
128	IMR SCD	Phase IIb	1.IMR-687 2.IMR-687 Placebo	13th August 2020	Dr. Seyram Kaali	•Korle-Bu Teaching Hospital •Kintampo Health Research Centre	IMARA Inc.	Early termination by Sponsor 1 Year 7 Months	study of subjects aged 18 to 65 years with SCD (HbSS, HbS0 thalassemia, or HbSB+ thalassemia) to evaluate the safety and efficacy of the PDE9 inhibitor, IMR-687, administered qd for 52 weeks. This study will provide data on IMR-687 doses of ≥ 3.0 to ≤ 4.5 mg/kg and > 4.5 to ≤ 6.7 mg/kg. In a relevant model of anemia (Hbbh1/th1 mice), oral administration of IMR-687 for 30 days at 30 mg/kg/day (human equivalent dose of 2.4 mg/kg/day) or 60 mg/kg/day (human equivalent dose of 4.9 mg/kg/day) increased RBCs and Hb, and reduced reticulocytes. The degree of these changes was dose dependent, with statistically significant improvement at the higher dose of 60 mg/kg. In addition, IMR-687 at 60
129	HESTIA4	Phase I	Ticagrelor	16th May, 2018	1. Dr. Patrick Ansaah 2. Dr. Catherine Segbefia 3. Dr. Kokou Hefoume Amegan-Aho	1. Navrongo Health Research Centre 2. Korle-Bu Teaching Hospital 3. Volta Regional Hospital	AstraZeneca AB	Study termination 31 Months	Complications of sickle cell disease (SCD) occur very early in life. Painful crises first appear in the fingers and toes (dactylitis) in very young children prior to their first birthday. In addition to painful crises occurring in the very young, SCD can affect organ function early in life. Loss of splenic function begins as early as 5 months of age with associated increase in infection risk. Stroke risk begins at age 2. Given the early onset of symptoms and complications of this disorder, therapies for SCD should be targeted at children, including the very young. There is a need to first establish the pharmacokinetics (PK) of ticagrelor in this age group to allow for modelling or extrapolation in this population. This goal of the study is to evaluate PK data in the 0-2 year old population in order to way for further studies and ultimately use of ticagrelor in this youngest population.
130	TADO	III	Prasugrel	20th may 2013	Prof. Tsiri Agbenyega Dr. Catherine Idara Segbefia	Center, Agogo Korle-Bu Teaching Hospital, Accra – Korle Bu	Eli Lilly and Company Indianapolis	Prematurely terminated 24 months	
131	WOMAN	III	Tranexamic acid(cyklokaprnr injection)	10th sept 2009	1. Dr. Anthony K. Dan 2. Dr. Opare Addo Henry Sakyi 3. Dr. Kwadwo Asamoah Nyarko-Jectey	1. Ashanti Mampong Municipal Hospital 2. Komfo Anokye Teaching Hospital	Clinical Trials Unit, London School of Hygiene and Tropical	Terminated by Sponsor Prematurely ended.	
132	NEOVITA	III	Vitamin A		Dr. Sam Newton	Kintampo Health Research Centre	PATH	Premature Termination 36 Months	
133	CALLASCOPE *	ii	Pocket Colposcope (CALLASCOPE)	12th February 2019	Dr. Emmanuel Srofenyoh	Ridge Hospital, Korle-Bu Teaching Hospital	Duke Global Health Institute	Study ended, FDA DISSOCIATED itself from any data or findings from the study due to violation of its guidelines for conducting clinical trials. 3 months	

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
SHORT AND DETAILED NAMES OF TRIALS									
1	4P		A strategy to reduce complications of Hypertensive disorders in Pregnancy and Maternal Mortality by 50% or more. - Polypill for the Prevention of Pregnancy Induced Hypertension and Preeclampsia (4P) Trial						
2	ABDOV COVID 19 TRIAL		A randomized, double-blind, positive-controlled Phase III clinical trial to evaluate the efficacy and safety of SCTV01E (A COVID-19 Alpha/Beta/Delta/Omicron Variants S Trimer Vaccine) in population previously unvaccinated with COVID-19 vaccine and aged ≥18 years						
3	ACTIVE TRIALS		A Phase 3, multicenter, randomized, double-blind, 24-week study of the clinical and antiviral effect of S-217622 compared with placebo in non-hospitalized participants with COVID-19						
4	AIM-LVRNA009		A Global Multi-center, Randomized, Blinded, Placebo-controlled Phase 2/3 Clinical Study to Evaluate the Efficacy, Safety and Immunogenicity of SARS-CoV-2 mRNA Vaccine (LVRNA009) for the Prevention of COVID-19 in Participants Aged 18 Years and Older						
5	AIMS		African Investigation Of Mirasol System For Whole Blood. Clinical And Biological Efficacy Of Mirasol Treated Fresh Whole Blood For The Prevention Of Transfusion Transmitted Malaria						
6	ALB_IVM		Comparison of Ivermectin alone with Albendazole (ALB) plus Ivermectin (IVM) in their efficacy against Onchocerciasis in the Volta Region, Ghana.						
7	ALBIVM K'SI		Comparism of Ivermectin Alone with Albendazole plus Ivermectin in Their Efficacy against Onchocerciasis						
8	AMARYL M		Clinical Efficacy and Safety of Amaryl M in Patients with Type 2 Diabetes who are inadequately treated by either Glimepride or Metformin Monotherapy or who are already treated With Free Combination Of Glimepride and Metformin in African Countries.						
9	ANTICOV		An Open-Label, Multicenter, Randomized, Adaptive Platform Trial of the Safety and Efficacy of Several Therapies, including Antiviral Therapies, Versus Control in Mild Cases of COVID-19						
10	ANTIPSYCHOTIC STUDY		A RANDOMIZED CONTROLLED TRIAL OF OMEGA-3 FATTY ACIDS IN THE TREATMENT OF ANTIPSYCHOTIC-INDUCED MOVEMENT DISORDERS IN GHANA						
11	AQUAMAT		An Open Randomized Comparism of Artesunate versus Quinine in the Treatment of Severe Falciparum Malaria in African Children.						
12	ARTIMIST		A Phase III, Randomized, Open Labelled, Active Controlled, Multicentre, Superiority Trial Of Artimisttm Versus Intravenous Quinine In Children With Severe Or Complicated Falciparum Malaria, Or Uncomplicated Falciparum Malaria With Gastrointestinal Complications						
13	ASAAP		A Multicentre Phase III Non-Inferiority Trial to Evaluate Safety, Tolerability and Efficacy of Artemether- Lumefantrine+Atovaquone-Proguanil Tri-TherapyVersus Artemether Lumefantrine Bi-Therapy for The Treatment of Uncomplicated Malaria in African Children Aged 6 To 59 Months (ASAAP PROJECT -STUDY II)						
14	ASTAWOL		The efficacy of Rifampicin 35mg/Kg/d plus Albendazole 400mg/d given for 7 or 14 days against Lymphatic Filariasis and Onchocerciasis- a randomized, controlled, parallel-group, open-label, phase II pilot trial						
15	AVAREF		A Phase 3 double-blind, randomized, active comparator-controlled, group-sequential, multinational trial to assess the safety, immunogenicity and efficacy of a trivalent rotavirus P2-VP8 subunit vaccine in prevention of severe rotavirus gastroenteritis in healthy infants.						
16	AX-100 HIV		A Double Blind Randomized Control Trial of AX-100 Immun (Liquid) and AX-100 Immun Plus Combination Among Adults Living with HIV In Ghana.						
17	AZI4YAWS		Randomized Controlled Trial Comparing Efficacy of a Single Dose of Treatment of Yaws with 20mg/kg versus 30mg/kg of Azithromycin.						
18	PLUS CHLOROQUINE		Azithromycin Plus Chloroquine Phosphate versus Artemether-Lumefantrine for the Treatment of Uncomplicated Plasmodium falciparum Malaria in Children in Africa.						
19	BEMPU		Hypothermia Prevention in low birth weight and preterm Infants						
20	BURULINOX		Evaluation of nitric oxide generating dressing (EDX) to improve management of buruli ulcer disease – a prospective randomized open-blinded end point.						
21	BURULIRIFDAC C		A randomized controlled trial to evaluate the effect of High Dose of Rifampicin and Dialkylcarbamoyl chloride (DACC)-coated dressings on outcomes in Mycobacterium ulcerans disease						

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
22	CDA								A Multicenter, Randomized, Double Blind Study to Compare the Efficacy and Safety of CDA Versus Artemether-Lumefantrine in the Treatment of Acute Uncomplicated P. Falciparum Malaria in Children and Adults in Africa.
23	CDA2								A Multicenter, Randomized, Double Blind Study to Compare the Efficacy and Safety of CDA Versus Chlorproguanil-Dapsone in the Treatment of Acute Uncomplicated P. Falciparum Malaria in Children and Adults in Africa.
24	CEREBETA								Efficacy of Beta-Glucans from Barley and Maintenance of Normal Blood LDL-Cholesterol Concentrations: A Randomized Control Study in Ghana.
25	CPAP								Clinical Trial Evaluating the Difference in Mortality Rates in Children in Ghana Receiving Continuous Positive Airway Pressure (CPAP) Versus Those Who Do Not.
26	CRASH-2								A Large Randomized Placebo Controlled Trial, among trauma patients with or at risk of significant Haemorrhage, of the Effects of Anti- Fibrinolytic treatment on Death and Transfusion requirement
27	CALLASCOPE								Clinical Studies and in-Depth Interviews for Portable, low-cost and Speculum-Free Cervical Cancer Screening in Ghana
28	CECOLIN								Phase 3 Randomized, Active-Comparator Controlled, Open-Label Trial to Evaluate the Immunogenicity and Safety of Alternate Two-Dose Regimens of a Bivalent Human Papillomavirus (HPV) Vaccine (Cecolin®) Compared to a Licensed Quadrivalent HPV Vaccine (Gardasil®) in Healthy 9-14 Year-Old Girls in Low and Low-Middle Income Countries
29	CEPHEIDXPRT HIV-1								An Investigation to Evaluate the Performance of the Cepheid XpertR HIV-1 VL XC Test
30	CIELO								A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter Basket Study to Evaluate the Efficacy, Safety, Pharmacokinetics, and Pharmacodynamics of Satralizumab in Patients with Anti-N-Methyl-D-Aspartic Acid Receptor (NMDAR) or Anti-Leucine-Rich Glioma-Inactivated 1 (LG1) Encephalitis
31	COPE TRIAL								Effectiveness and Acceptability of two models of an Insertable Vaginal Cup for Non-surgical management of obstetric fistula in Ghana: a hybrid type 1 randomized crossover trial
32	COVID ABDOV								A randomized, double-blind, positive-controlled Phase III clinical trial to evaluate the efficacy and safety of SCTV01E (A COVID-19 Alpha/Beta/Delta/Omicron Variants S Trimer Vaccine) in population previously unvaccinated with COVID-19 vaccine and aged ≥18 years* (COVID ABDOV).
33	CROWN CORONATION								An international, Bayesian platform adaptive, randomized, placebo-controlled trial assessing the effectiveness of candidate interventions in preventing COVID-19 disease in healthcare workers
34	CHEETAH								Cluster Randomized Trial of Sterile Glove and Instrument Change at the Time of Wound Closure to Reduce Site Infection: A Trial In Low- And Middle-Income Countries (LMICs)
35	COVID 19 CHO-CELL								A multicenter, randomized, double-blind, placebo-controlled Phase II/III trial to evaluate the efficacy, safety and immunogenicity of the recombinant two-component COVID-19 vaccine (CHO cell) in adults aged 18 years and older
36	COVID 19 INTRANASAL SPRAY								A Global, Multi-center, Randomized, Double-blind, Placebo-controlled Phase III Clinical Trial to Evaluate the Protective Efficacy and Safety of Influenza Virus Vector COVID-19 Vaccine for Intranasal Spray (DelNS1-2019-nCoV-RBD-OPT1) in Adults Aged 18 Years and Older
37	COVID 19 MOUTHWASH								Viral Shedding Dynamics and the Effect of Antimicrobial Mouthwashes on the Detection of SARS-CoV-2 in Ghana.
38	DIABETIC FOOT CARE								Family-oriented Diabetic Foot Self-care Programme in Ghana; A Feasibility Randomised Controlled Trial with nested qualitative interviews at the Komfo Anokye Teaching Hospital.
39	DOLF_IDA								Safety and Efficacy of Combination Therapy with Ivermectin, Diethylcarbamazine and Albendazole (IDA) for Individuals with Onchocerciasis
40	EBA								Double-Blinded, Placebo-Controlled Dosage-Escalation Study and Immunogenicity of EBA-175 RII-NG Malaria Vaccine Administered Intramuscularly in Semi Immune Adults
41	EBOLA Z								A Phase 2, Randomized, Observer-Blind, Placebo-Controlled, Multi-Country Study to Assess the Safety and Immunogenicity of a Single Intramuscular Dose of GSK Biologicals' Investigational Recombinant Chimpanzee Adenovirus Type 3 – Vectored Ebola Zaire Vaccine. (ChAd3-EBO-Z) (GSK3390107A), in Adults 18 years of age and older in Africa
42	EBOLA Z (PAEDIATRIC)								A Phase 2, Randomized, Observer-Blind, Placebo-Controlled, Multi-Country Study to Assess the Safety and Immunogenicity of a Single Intramuscular Dose of GSK Biologicals' Investigational Recombinant Chimpanzee Adenovirus Type 3 – Vectored Ebola Zaire Vaccine. (ChAd3-EBO-Z) (GSK3390107A), in children 1 to 17 years of age in Africa
43	EBSI-LSV								A Phase 1 Randomized, Blinded, Placebo Controlled, Dose-Escalation and Dosing Regimen Selection Study to Evaluate the Safety and Immunogenicity of rVSV-Vectored Lassa Virus Vaccine in Healthy Adults at Multiple Sites in West Africa
44	ELDON CARD								Using Eldon Card for Testing of Maternal and Newborn Blood Group in Comparison with the Standard Laboratory Method of Blood Group Testing in Accra, Ghana
45	EMODEPSIDE								A phase II, Randomised, double-blind, parallel – group trial to investigate Emodepside (BAY 44-4400) in subjects with onchocerca volvulus infection.
46	ESM UBT								A Multi-Centre Prospective Trial on the Impact of the Introduction of Condom-Based Uterine Balloon Tamponade for Uncontrolled Postpartum Hemorrhage
47	FALCON								Pragmatic Multicentre Factorial Randomized Controlled Trial Testing Measures to Reduce Surgical Site Infection in Low and Middle Income Countries
48	FERROQUINE								Randomized Multicentre Study Evaluating the Safety and Activity of Ferroquine Associated with Artesunate versus a Positive Calibrator (Amodiaquine Associated with Artesunate) In African Adult Patients with Uncomplicated Malaria

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
49	BUILLON CUBES STUDY								Effect of household use of multiple micronutrient-fortified bouillon on micronutrient status among women and children in two districts in the Northern region of Ghana
50	GARDASIL								Evaluation of Safety And Immunogenicity Of Gardasilm In Healthy Females Between 9 And 26 Years Of Age In Sub-Saharan Africa
51	GBT 2104-131								A Randomized, Double-blind, Placebo-controlled, Multicenter Study to Assess the Safety and Efficacy of Inclacumab in Participants with Sickle Cell Disease Experiencing Vasoocclusive Crises.
52	GBT-2104-132								A Randomized, Double-blind, Placebo-controlled, Multicenter Study of a Single Dose of Inclacumab to Reduce Re-admission in Participants with Sickle Cell Disease and Recurrent Vaso-occlusive Crises
53	GBT-2104-133								An Open-Label Extension Study to Evaluate the Long-Term Safety of Inclacumab Administered to Participants with Sickle Cell Disease Who Have Participated in an Inclacumab Clinical Trial.
54	GMZ 2								Randomized, Controlled, Double-Blind, Multicentre Study To Evaluate The Efficacy, Safety And Immunogenicity Of GMZ2 Candidate Malaria Vaccine In Gabonese, Burkinabe, Ghanaian And Ugandan Children Aged 12-60 Months
55	PHARMACOGEN OMICS TRIALS								Development of Precision Medicine Approaches to Improve Effectiveness of Hydroxyurea (HU) Treatment for Sickle Cell Disease (SCD) in 3 Low and Middle-Income Countries (LMIC)
56	ANTIMALARIAL								A Phase III of the Assessment of the Efficacy, Tolerability and Ease of Administration of, Dihydroartemisinin Plus Piperazine and Artesunate Plus Sulfamethoxypyrazine Plus Pyrimethamine for preventing Malaria in Ghanaian Children
57	HOPE SCD								A Phase 3, Double-blind, Randomized, Placebo-controlled, Multicenter Study of GBT440 Administered Orally to Patients With Sickle Cell Disease
58	HOPE KIDS 2								A phase 3, Randomised, Double-Blind, Placebo-Controlled Study of Voxelotor (GBT440) in Pediatric Participants with Sickle Cell Disease.
59	HYDRANON								Hydranon® solution (GR-08) in healthy adult volunteers
60	HESTIA4								A Multi-centre, Phase I, Open-label, Single-dose Study to Investigate Pharmacokinetics (PK) of Ticagrelor in Infants and Toddlers, Aged 0 to less than 24 Months, with Sickle Cell Disease
61	HESTIA3								A Randomised, Double-Blind, Parallel-Group, Multicentre, Phase III Study to Evaluate the Effect of Ticagrelor versus Placebo in Reducing the Rate of Vaso-Occlusive Crises in Paediatric Patients with Sickle Cell Disease
62	IMR-SCD-301								A Phase 2b Study to Evaluate the Safety and Efficacy of IMR-687 in Subjects with Sickle Cell Disease
63	INNOVATE								Phase 2/3 Randomized, Blinded, Placebo-Controlled Trial to Evaluate the Safety, Immunogenicity, and Efficacy of INO-4800, a Prophylactic Vaccine against COVID-19 Disease, Administered Intradermally Followed by Electroporation in Adults at High Risk of SARS-CoV-2 Exposure
64	INO-9112 COVID 19								Phase 1 Open Label, Randomized Study to Evaluate the Safety, Tolerability, and Immunogenicity of an Intradermal Booster Dose of INO-4800 alone or in combination with INO-9112 followed by Electroporation in Adults who Completed a Primary Immunization Series Against SARS-CoV-2 with mRNA Vaccines
65	INVACT								In Vivo Efficacy of Artemisinin Combination Therapy to Explore Laboratory and Parasitological Markers of Artemisinin Resistance in Uncomplicated Plasmodium falciparum Malaria in Ghana.
66	IPT & SP								Operational Research on Intermittent Preventive Treatment of Malaria in Infants (IPTi) with Sulfadoxine/Pyrimethamine (S/P)
67	INSUGEN								Post Market Surveillance Study of Insugen 30/70
68	INOVIO – LASSA FEVER								Study to evaluate the safety, tolerability and immunogenicity of INO-4500 in Healthy volunteers
69	IRON FORTIFICATION								Seasonal Impact Of Iron Fortification On Malaria Incidence In Ghanaian Children
70	IVERMECTIN GH								Safety and Efficacy of Ivermectin in the Prevention and Management of COVID- 19 among Ghanaian Populations

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
71	KAE609	A Phase 2, Multi-Center, Randomized, Open - Label, Dose Escalation Study To Determine Safety Of single (QD) and Multiple (3QD) Doses Of KAE609, Given To Adults With Uncomplicated Plasmodium Falciparum Malaria							
72	KNC 19(NIBIMA)	Repurposing the aqueous Extract of Cryptolepis for Covid-19 therapy							
73	LEDoxy	Doxycycline 200mg/d vs. 100mg/d for 6 weeks to improve filarial lymphedema - a multinational, double-blind, randomized, placebo-controlled trial.							
74	LETICIA	Combination Food-Based And Supplemental Iron Replacement Therapy For Children With Moderate-To-Severe Anemia In A Rural Ghanaian Setting:A Proof-Of-Concept Study							
75	LIVZON	A Global, Multi-Center, Randomized, Double-Blind, Placebo-Controlled, Phase III Clinical Study to Evaluate the Efficacy, Safety, and Immunogenicity of Recombinant SARS-CoV-2 Fusion Protein Vaccine (V01) in Adults Aged 18 Years and older.							
76	MAL 047	Randomized, Controlled, Partially-Blind Study Of The Safety And Immunogenicity Of Glaxosmithkline Biologicals' Candidate Plasmodium Falciparum Vaccines RTS,S/AS02D And RTS,S/AS01E, When Administered IM According To A Three Dose Schedules In Children Aged 5 To 17 Months Living In Ghana.							
77	MAL 050	Randomized, Open, Controlled Study Of The Safety Of The And Immunogenicity Of GSK Biologicals' Candidate Plasmodium Falciparum Malaria vaccine RTS, S/AS01E when incorporated into an expanded program on immunization (EPI) regimen that includes DTPWHEPB/HIB.OPV, Measles and yellow fever vaccination in infants living in malaria- Endemic Regions- 050							
78	MAL 055	Double Blind (Observer Blind), Randomised, Controlled Multicentre Study To Evaluate In Infants And Children, The Efficacy Of RTS,S/AS10E Candidate Vaccine Against Malaria Disease Caused By P. Falciparum Infection Across Diverse Malaria Transmission Settings In Africa							
79	MAL 063	Randomized, Open, Controlled Study To Evaluate The Immune Response To The Hepatitis B Antigen Of The RTS,S /AS01E Candidate Vaccine, When Administrated As Primary Vaccination Integrated Into An EPI Regimen To Infants Living In Sub-Saharan Africa							
80	MAL 073	Phase IIb randomized, open, controlled, multi-center study to evaluate the immunogenicity and safety of the RTS,S/AS01E candidate malaria vaccine, when administered as primary vaccination at 6, 7.5 and 9 months of age with or without co-administration of measles, rubella and yellow fever vaccines followed by an RTS,S/AS01E booster vaccination 18 months post Dose 3, to children living in sub-Saharan Africa							
81	MAL 094	Phase IIb Randomized, Open-Label, Controlled, Multi-Centre Study of the Efficacy, Safety and Immunogenicity of GSK Biologicals' Candidate Malaria Vaccine RTS,S/AS01E Evaluating Schedules with or without Fractional Doses, early Dose 4 and yearly Doses, in Children 5-17 Months of age Living in Sub-Saharan Africa.							
82	MDGH-MOX-1006	An open-label study of the pharmacokinetics and safety of a single dose of moxidectin per oral in subjects aged 4 to 17 years with (or at risk of) onchocerciasis to identify an optimal dose for treatment of children 4 to 11 years							
83	MEBENDAZOLE MEFLOQCHLOA	Efficacy and Safety Of A Single Dose Regimen And A Multi Dose Regimen Of Mebendazole Against Hookworm Infections In Children And Adolescents In Ghana : A Randomized Control Trail.							
84	ZITH	A Phase III, Randomized, Opened-Label, Comparative Trial Of Azithromycin Plus Chloroquine Versus Mefloquine For The Treatment Of Uncomplicated Plasmodium Falciparum Malaria In Africa.							
85	AL-A CONJUGATE	A Phase II, Double Blind, Randomized, Controlled, Dose Ranging Study to Evaluate the Safety, Immunogenicity Dose Response and Schedule Response of a Meningococcal A Conjugate Vaccine administered concomitantly with local EPI vaccines in Healthy Infants.							
86	MMS	The Use Of A Multiple Micronutrient Supplement In Women Of Reproductive Age							
87	MoRiOn	The Efficacy of Rifapentine 900mg/d plus Moxifloxacin 400mg/d given for 14 or 7 days against Onchocerciasis – a Randomized, Controlled, Parallel-Group, Open Label, Phase II Pilot Trial							
88	MOXIDECTIN	Randomized, single-ascending dose, Ivermectin-controlled, double-blind, safety, tolerability, pharmacokinetic and efficacy study of orally administered Moxidectin in subjects with Onchocerca volvulus Infection							
89	MOXIDECTIN-IVERMECTIN	A Phase III Randomized, Single-Ascending-Dose, Ivermectin-Controlled, Double-Blind, Safety, Tolerability, Pharmacokinetic, and Efficacy Study of Orally Administered Moxidectin in Subjects with Onchocerca volvulus Infection':							
90	MULTIMAL	Multi-Drug Combination-Therapies to prevent the Development of Drug Resistance: Phase II Controlled Clinical Trial Assessing Candidate Regimens of Multiple-Antimalarial Combinations for the Treatment of Uncomplicated Malarial in Africa							
91	MYCOPIROX_LA GRAY	Randomized, open labelled trial to evaluate the efficacy, safety and tolerability of mycopirox vaginal cream in the treatment of mixed infection vaginitis							
92	NEOVITA	Feasibility Studies							
93	NOGUCHI FILARIASIS	Determination of the Prevalence of LF Infection in Districts Not Included in LF Control Activities and of the Basis for Integrated Implementation of LF - Onchocerciasis Elimination Strategies in Potentially Co-endemic Areas							
94	NOGUCHI SCD	A Phase 1B Dose – Finding Pharmacokinetics and Pharmacodynamic Study Oof NVX – 508 In Sickle Cell Disease (SCD) Patients							
95	NON-INVASIVE HAEM DEVICE	A Comparison of Hemoglobin Values as Measured By The Pronto And Pronto 7 Non-Invasive Hemoglobin Devices, The Hemocue Hb 201+, And A Hematology Analyzer Among Pregnant Women Attending Antenatal Care Clinic In Ghana							

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
96	NOVASIL								Safety and Efficacy Evaluation of Novasil: Strategy for the Protection of Humans from Aflatoxin Toxicity
97	NOVIC TRIAL								Novel vacuum-induced Haemorrhage control for postpartum Haemorrhage: a multicentre randomised trial
98	OXYTOCIN								Determining the Effect of Prophylactic Administration Of Oxytocin In Unject™ By A Community Health Officer On Post-Partum Haemorrhage At Home Births In The Kintampo North And South Districts Of Ghana
99	PFCSP_MVACS_MALARIA								Partial Double-Blind, Randomized Study of PFCSP DNA/MVA Prime Boost Vaccine
100	PIVOT								Prospective Identification of Variables as Outcomes for Treatment (PIVOT): A Phase II clinical trial of hydroxyurea for children and adults with HbSC disease
101	POLYPHENOL-RICH COCOA POWDER TRIAL POST								Polyphenol-rich Cocoa Powder as Adjuvant Therapy in Patients with Covid-19.
102	MASTECTOMY								ULTRASOUND-GUIDED ERECTOR SPINAE PLANE BLOCK FOR POST-MASTECTOMY PAIN RELIEFve
103	PRAISE								An adaptive, Randomized, Placebo-controlled, Double-Blind, Multi-center Study of Oral FT-4202, a Pyruvate Kinase Activator in Patients with Sickle Cell disease (PRAISE)
104	PREGACT								Evaluating the Safety And Efficacy Of Artemisinin-Based Combination Treatments For African Pregnant Women With Malaria
105	PRENABELT								A Maternal Device to Reduce the Risk of Stillbirth and Low-Birth Weight
106	PROBIOTIC								A double-blind randomized control trial of a synbiotic vs. placebo among pregnant women to evaluate colonization of the gut microbiota of their infants with Lactobacillus plantarum (Probiotics pilot in Ghana)
107	ARTESUNATE VRS COARTEM								andomized multicentre clinical study to assess the safety and efficacy of fixed dose formulation of oral pyronaridine artesunate tablet versus coartem in children and adult patients with acute uncomplicated plasmodium falciparum malaria
108	PRCR DIPSTICK								Validation of a Protein Creatinine (PrCr) Dipstick Diagnostic Test for Proteinuria Screening on Antenatal Care Clinics in Ghana
109	PRCR SPOT								Evaluating the clinical utility and operational fit of the lifeAssay Diagnostics Test-It™ PrCr urinary dipstick test to assess risk of pre-eclampsia in referral hospitals in Ghana: A SPOT nested study, developing and VALidating a Severe Pre-eclampsia adverse Outcome Triage (SPOT) score
110	RECOVERY								Randomized Evaluation of Covid-19 Therapy (RECOVERY)
111	RIFAMPIN VS ISONIAZID								A Randomized Clinical Trial of 4 months Rifampin versus 9 months Isoniazid for treating Latent TB Infection
112	ROBOCOW								RANDOMIZED PLACEBO-CONTROLLED TRIAL TESTING 0.2% CHLORHEXIDINE MOUTHWASH TO REDUCE POSTOPERATIVE RESPIRATORY TRACT INFECTIONS IN ABDOMINAL SURGERIES
113	ROTARIX								Immunogenicity of The Human Rotavirus Vaccine (Rotarix™) At Varying Schedules and Ages in Rural Ghana
114	ROTASHIELD								The Randomized, Double-Blind, Placebo-Controlled Evaluation of The Efficacy, Immunogenicity, and Safety of 2 Single Doses of RRV-TV in Neonates/Infants
115	ROTATEQ								Efficacy, Safety and Immunogenicity of Rotateq™ Among Infants in Africa and Asia.
116	SALIF								A Phase 3b, Randomized, Open-label Clinical Study to Demonstrate non-inferiority in Virologic Response Rates of HIV-1 RNA Suppression <400 Copies/mL of TDF/FTC/RPV Versus TDF/FTC/EFV in First-line Antiretroviral NNRTI-based Suppressed Patients Switching At Low HIV-1 RNA Into Fixed Dose Combinations
117	SAR97276A_SA NOFI								A Multicentre, Open Label, Efficacy And Safety Of Parenteral Sar97276a In The Treatment Of Symptomatic Uncomplicated And Severe Malaria In Adults And Children
118	SAVVY								Randomised Controlled Trials of Savvy In HIV
119	SAVING BRAINS KUMASI								Saving Brains from Malnutrition: Implementation of Evidence-Based Nutritional Supplementation and Psychosocial Stimulation Program for Pregnant and Lactating Women and their Infants Post Weaning, To Improve Cognition and Behavioral Regulation to Deliver Better Social and Economic Prospects Later in Life
120	SAVING BRAINS NAVORONGO								Saving Brains from Malnutrition: Implementation of Evidence-Based Nutritional Supplementation and Psychosocial Stimulation Program for Pregnant and Lactating Women and their Infants Post Weaning, To Improve Cognition and Behavioral Regulation to Deliver Better Social and Economic Prospects Later in Life

CLINICAL TRIALS REGISTRY

N/O	TITLE OF STUDY	PHASE	Investigational Products (IPs)	DATE OF RECEIPT OF APPLICATION	PRINCIPAL INVESTIGATOR	STUDY CENTRE(S)	SPONSORS & APPLICANT	STATUS & DURATION OF STUDY	PURPOSE/AIM OF STUDY
121	SHEA LIDO	Comparison of Shea butter and Lidocaine gel for rectal examination- A Non-Inferiority Trial							
122	SMAC	A Comparative, Open Label, Dose And Regimen Optimization Follow-Up Study Of Intravenous And Intramuscular Artesunate In African Children With Severe Malaria.							
123	SMAART	Stroke Minimization through Additive Anti-atherosclerotic Agents in Routine Treatment							
124	SPUTNIK LIGHT	A phase III randomized double blind, placebo- controlled international multisite clinical trial in parallel assignment to evaluate efficacy, immunogenicity and safety of the sputnik light vector vaccine in adults in the sars-cov-2 infection prophylactic treatment							
125	STAND	A Phase III, Multi-Centre, Randomized, Double-Blind Study to Assess Efficacy and Safety of Two Doses of Crizanlizumab Versus Placebo With or Without Hydroxyurea/Hydroxycarbamide Therapy in Adolescent and Adult Sickle Cell Disease Patients with Vaso Occlusive Crises (STAND)							
126	STAR	POSTOPERATIVE PAIN MANAGEMENT IN EMERGENCY ABDOMINAL SURGERY: BIMODAL VERSUS UNIMODAL ANALGESIA							
127	STEADFAST	A Phase II, multicenter, randomized, open label two arm study comparing the effect of crizanlizumab + standard of care to standard of care alone on renal function in sickle cell disease patients ≥ 16 years with chronic kidney disease due to sickle cell nephropathy							
128	SWIS	Feasibility, Acceptability, and Outcomes of Sterile Water Injection (SWI) in Managing Lower Back Pain among Labouring Women in a Tertiary Hospital in Ghana: A Mixed-method Study							
129	TADO	Double-Blind, Randomized, Efficacy And Safety Comparison Of Prasugrel And Placebo In Pediatric Patients With Sickle Cell Disease							
130	TENOFOVEK BE	A balanced, randomized, two treatment, two-period, two-sequence single dose crossover, open-label, analyst blind and single centre bioequivalence study test product; Tenofovek of Danadams Pharmaceuticals Industry Ltd., Ghana and reference product; Viread (Gilead Sciences, Inc., CA, USA) in healthy, Ghanaian adult, male, human participants under fasting conditions.							
131	TENOFOVIR	A Phase II Study for Tenofovir Disoproxil Fumarate for Prevention of HIV							
132	TYVEGHA	A cluster-randomized controlled Phase IV trial assessing the impact of a Vi-Polysaccharide conjugate vaccine in preventing typhoid infection in Asante Akim, Ghana (TyVEGHA)*:							
133	VAT00008	A parallel-group, Phase III, multi-stage, modified double-blind, multi-armed study to assess the efficacy, safety, and immunogenicity of two SARS-CoV-2 Adjuvanted Recombinant Protein Vaccines (monovalent and bivalent) for prevention against COVID-19 in adults 18 years of age and older							
134	VERO CELL COVID 19 TRIAL	A Randomized, Double-Blinded, Placebo-Controlled, Phase III, Clinical Trial of SARS-CoV-2 Vaccine, Inactivated (Vero Cell) in Adults Aged 18 Years and Above							
135	VR-AD-1005 STUDY	Assessment of a novel fixed dose combination (FDC) drug VR-AD-1005 for the treatment of acute watery diarrhea in cholera: A phase II, multicenter, randomized, placebo controlled, double blinded efficacy and safety trial							
136	VERTEX	A Phase 2/3 Adaptive, double-blind, placebo-controlled study to evaluate the efficacy and safety of VX-147 in Subjects Aged 18 Years and Older with APOL1-mediated Proteinuric Kidney Disease							
137	WOMAN	Tranexamic Acid For The Treatment Of Postpartum Haemorrhage: An International, Randomized, Double Blind, Placebo Controlled Trial							
138	YAWS	Single Dose Oral Azithromycin Versus Injection Benzathine Penicillin For The Treatment Of Yaws – A Randomized Clinical Trial In Some Endemic Communities In Ghana							
139	ZEBOV	A Phase 1 Study to Evaluate the Safety, Tolerability and Immunogenicity of Heterologous Prime-Boost Regimens Using MVA-BN®-FILO and Ad26.ZEBOV Administered in Different Sequences and Schedules in Healthy Adults							
140	ZEBOV 2	A Randomised, Observer-blind, Placebo-controlled, Phase 2 Study to Evaluate the Safety, Tolerability and Immunogenicity of Three Prime-boost Regimens of the Candidate Prophylactic Vaccines for Ebola AD26ZEBOV and MVA-BN-Filo in Healthy Adults,							
141	ZIV AFFLIBERCEPT	Phase I, Safety of ZIV-AFLIBERCEPT in retinal diseases in Ghanaian population							
142	*	Feasibility Studies							
143	N/A	Study not Started/ Application Withdrawn /Not Approved / Terminated / FDA Dissociation from Trial data							
144	NYN	Not yet known							
145	Active Trials								
146	Applications pending approval								
147	Study ended								
148	Trials closed by Sponsor before commencement								
149	Application withdrawn by Sponsor before FDA approval								
150	Application closed by FDA								

