



Pfizer

**Campto®**

IRINOTECAN HYDROCHLORIDE TRIHYDRATE

CDS

Ghana, Kenya, Nigeria, Uganda

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

C AMPTO<sup>®</sup> injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Active ingredient: irinotecan hydrochloride

Each milliliter (mL) of sterile solution contains 20 mg of irinotecan hydrochloride (on the basis of the trihydrate salt).

## **3. PHARMACEUTICAL FORM**

Irinotecan sterile solution is a pale yellow, clear, aqueous solution requiring dilution for intravenous administration.

## **4. CLINICAL PARTICULARS**

### **4.1. Therapeutic indications**

Irinotecan is indicated for single-agent or combination treatment of patients with:

- Metastatic carcinoma of the colon or rectum that has recurred or progressed following 5-fluorouracil (5-FU)-based therapy
- Previously untreated metastatic carcinoma of the colon or rectum
- Non-small-cell lung cancer
- Small-cell lung cancer
- Cervical cancer
- Ovarian cancer
- Inoperable or recurrent gastric cancer
- Esophageal cancer

Irinotecan in combination with cetuximab is indicated for the treatment of patients with epidermal growth factor receptor (EGFR)-expressing, KRAS wild-type metastatic colorectal cancer, who had not received prior treatment for metastatic disease or after failure of irinotecan-including cytotoxic therapy (see Section 5.1 Pharmacodynamic properties).

Irinotecan in combination with 5-FU, folinic acid (FA) and bevacizumab is indicated for first-line treatment of patients with metastatic carcinoma of the colon or rectum (see Section 5.1 Pharmacodynamic properties).

Irinotecan in combination with capecitabine with or without bevacizumab is indicated for first-line treatment of patients with metastatic colorectal carcinoma (see Section 5.1 Pharmacodynamic properties).

Irinotecan is indicated for single-agent treatment of patients with:

- Inoperable or recurrent breast cancer
- Squamous cell carcinoma of skin
- Malignant melanoma
- Malignant lymphoma
- Pancreatic cancer
- Glioma

#### **4.2. Posology and method of administration**

All doses of irinotecan should be administered as an intravenous infusion over 30 to 90 minutes.

##### **Single-agent dosage schedules**

Single-agent dosage schedules have been extensively studied for metastatic colorectal cancer. These regimens may be used in the treatment of patients with other indicated cancers (see Section 4.1 Therapeutic indications).

##### Starting dose

*Weekly Dosage Schedule.* The recommended single-agent starting dose of irinotecan is 125 mg/m<sup>2</sup>. A lower starting dose may be considered (e.g., 100 mg/m<sup>2</sup>) for patients with any of the following conditions: prior extensive radiotherapy, performance status of 2, increased bilirubin levels, or gastric cancer. Treatment should be given in repeated 6-week cycles, comprising weekly treatment for 4 weeks, followed by a 2-week rest.

*Once-Every-2-Week Dosage Schedule.* The usual recommended starting dose of irinotecan is 250 mg/m<sup>2</sup> every 2 weeks by intravenous infusion. A lower starting dose may be considered (e.g., 200 mg/m<sup>2</sup>) for patients with any of the following conditions: age 65 years and older, prior extensive radiotherapy, performance status of 2, increased bilirubin levels, or gastric cancer.

*Once-Every-3-Week Dosage Schedule.* The usual recommended starting dose of irinotecan for the once-every-3-week dosage schedule is 350 mg/m<sup>2</sup>. A lower starting dose may be considered

(e.g., 300 mg/m<sup>2</sup>) for patients with any of the following conditions: age 65 years and older, prior extensive radiotherapy, performance status of 2, increased bilirubin levels, or gastric cancer.

### Special populations

Patients with impaired hepatic function In patients with hepatic dysfunction, the following starting doses are recommended:

**Table 1. Starting Doses in Patients with Hepatic Dysfunction: Single-Agent Weekly Regimen**

Serum Total Bilirubin Concentration	Serum ALT/AST Concentration	Starting Dose, mg/m <sup>2</sup>
1.5-3.0 x IULN	≤5.0 x IULN	60
3.1-5.0 x IULN	≤5.0 x IULN	50
<1.5 x IULN	5.1-20.0 x IULN	60
1.5-5.0 x IULN	5.1-20.0 x IULN	40

**Table 2. Starting Doses in Patients with Hepatic Dysfunction: Single-Agent Once-Every-3-Week Regimen**

Serum Total Bilirubin Concentration	Starting Dose, mg/m <sup>2</sup>
1.5-3.0 x IULN	200
>3.0 x IULN	Not Recommended <sup>a</sup>

a. The safety and pharmacokinetics of irinotecan given once-every-3-weeks have not been defined in patients with bilirubin >3.0 x institutional upper limit of normal (IULN) and this schedule cannot be recommended in these patients.

### Patients with impaired renal function

Studies in this population have not been conducted (see Section 5.2 Pharmacokinetic properties). Therefore, caution should be undertaken in patients with impaired renal function. Irinotecan is not recommended for use in patients on dialysis.

### Combination-agent dosage schedules

#### Starting dose

*Irinotecan in Combination with 5-Fluorouracil (5-FU) and Leucovorin in every 2 weeks Schedule.* Irinotecan in combination with 5-FU and leucovorin is recommended for use in patients with metastatic colorectal cancer. For all regimens, the dose of leucovorin should be administered immediately after irinotecan, with the administration of 5-FU to occur immediately after receipt of leucovorin. The currently recommended regimens are shown below:

- Regimen 1 (6-week cycle with bolus 5-FU/LV): The recommended starting dose is 125 mg/m<sup>2</sup> of irinotecan, 500 mg/m<sup>2</sup> bolus 5-FU, and 20 mg/m<sup>2</sup> bolus leucovorin.
- Regimen 2 (6-week cycle with infusional 5-FU/LV): The recommended starting dose is 180 mg/m<sup>2</sup> of irinotecan, 400 mg/m<sup>2</sup> bolus 5-FU, 600 mg/m<sup>2</sup> 5-FU infusion, and 200 mg/m<sup>2</sup> leucovorin.

Lower starting doses may be considered for irinotecan (e.g., 100 mg/m<sup>2</sup>) and 5-FU (e.g., 400 mg/m<sup>2</sup>) for patients with any of the following conditions: age 65 years and older, prior extensive radiotherapy, performance status of 2, increased bilirubin levels, or gastric cancer. Treatment should be given in repeated 6-week cycles, comprising weekly treatment for 4 weeks, followed by a 2-week rest.

*Irinotecan in Combination with Cisplatin.* Irinotecan has been studied in combination with cisplatin for non-small cell and small cell lung cancer, cervical cancer, gastric cancer, and esophageal cancer. This regimen may be used in the treatment of patients with other indicated cancers, except for colorectal cancer (see Section 4.1 Therapeutic indications).

The recommended starting dose is 65 mg/m<sup>2</sup> of irinotecan and 30 mg/m<sup>2</sup> of cisplatin. A lower starting dose of irinotecan (e.g., 50 mg/m<sup>2</sup>) may be considered for patients with any of the following conditions: age 65 years and older, prior extensive radiotherapy, performance status of 2, increased bilirubin levels, or gastric cancer. Treatment should be given in repeated 6-week cycles, comprising weekly treatment for 4 weeks, followed by a 2-week rest.

*Irinotecan in Combination with Cetuximab.* For the posology and method of administration of concomitant cetuximab, refer to the full prescribing information for cetuximab. Normally, the same dose of irinotecan is used as administered in the last cycles of the prior irinotecan-containing regimen. Irinotecan must not be administered earlier than 1 hour after the end of the cetuximab infusion.

*Irinotecan in Combination with Bevacizumab.* For the posology and method of administration of bevacizumab, refer to the full prescribing information for bevacizumab. Bevacizumab is recommended in combination with irinotecan (125 mg/m<sup>2</sup>) / bolus 5-FU (500 mg/m<sup>2</sup>) / folinic acid (20 mg/m<sup>2</sup>) given once weekly for 4 weeks every 6 weeks.

*Irinotecan in Combination with Capecitabine.* For the posology and method of administration of capecitabine, see section 5.1 and refer to the full prescribing information for capecitabine. Capecitabine is recommended in combination with irinotecan at a starting dose of 1000 mg/m<sup>2</sup> for 2 weeks every 3 weeks.

### **Duration of treatment**

For both single-agent and combination-agent regimens, treatment with additional cycles of irinotecan may be continued indefinitely in patients who attain a tumor response or in patients whose cancer remains stable. Patients should be carefully monitored for toxicity and should be

removed from therapy if unacceptable toxicity occurs that is not responsive to dose modification and routine supportive care.

### **Dose modification recommendations**

The recommended dose modifications during a cycle of therapy and at the start of each subsequent cycle of therapy for single-agent dosage schedules are described in Table 3. These recommendations are based on toxicities commonly observed with the administration of irinotecan. For modifications at the start of a subsequent cycle of therapy, the dose of irinotecan should be decreased relative to the initial dose of the previous cycle.

The recommended dose modifications during a cycle of therapy and at the start of each subsequent cycle of therapy for irinotecan, 5-FU, and leucovorin are described in Table 4.

The recommended dose modifications for irinotecan and cisplatin for the start of each cycle of therapy are described in Table 5, while recommended dose modifications during a cycle of therapy are described in Table 6.

Recommendations for dose modifications of cetuximab when administered in combination with irinotecan must be followed according to the full prescribing information for cetuximab.

Refer to the full prescribing information for bevacizumab for dose modifications of bevacizumab when administered in combination with irinotecan/5-FU/FA.

In combination with capecitabine for patients 65 years of age or more, a reduction of the starting dose of capecitabine to 800 mg/m<sup>2</sup> twice daily is recommended according to the full prescribing information for capecitabine. Refer also to the recommendations for dose modifications in combination regimen given in the full prescribing information for capecitabine.

All dose modifications should be based on the worst preceding toxicity. A new cycle of therapy should not begin until the toxicity has recovered to Grade 2 or less. Treatment may be delayed 1 to 2 weeks to allow for recovery from treatment-related toxicity. If the patient has not recovered, consideration should be given to discontinuing irinotecan.

**Table 3. Recommended Dose Modifications For Single-Agent Schedules**

A new cycle of therapy should not begin until the granulocyte count has recovered to $\geq 1500/\text{mm}^3$ , and the platelet count has recovered to $\geq 100,000/\text{mm}^3$ , and treatment-related diarrhea is fully resolved. Treatment should be delayed 1 to 2 weeks to allow for recovery from treatment-related toxicities. If the patient has not recovered after a 2-week delay, consideration should be given to discontinuing irinotecan.			
Toxicity NCI Grade <sup>b</sup> (Value)	During a Cycle of Therapy	At the Start of the Next Cycle of Therapy (After Adequate Recovery), Compared with the Starting Dose in the Previous Cycle <sup>a</sup>	
	Weekly	Weekly	Once Every 2 or 3 Week
No toxicity	Maintain dose level	$\uparrow$ 25 mg/m <sup>2</sup> up to a maximum dose of 150 mg/m <sup>2</sup>	Maintain dose level
Neutropenia 1 (1500 to 1999/mm <sup>3</sup> ) 2 (1000 to 1499/ mm <sup>3</sup> ) 3 (500 to 999/ mm <sup>3</sup> ) 4 (<500/ mm <sup>3</sup> )	Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> Omit dose, then $\downarrow$ 25 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2 Omit dose, then $\downarrow$ 50 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2	Maintain dose level Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>	Maintain dose level Maintain dose level $\downarrow$ 50 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>
Neutropenic fever (Grade 4 neutropenia & $\geq$ Grade 2 fever)	Omit dose, then $\downarrow$ 50 mg/m <sup>2</sup> when resolved	$\downarrow$ 50 mg/m <sup>2</sup>	$\downarrow$ 50 mg/m <sup>2</sup>
Other hematologic toxicities	Dose modifications for leukopenia, thrombocytopenia, and anemia during a cycle of therapy and at the start of subsequent cycles of therapy are also based on NCI toxicity criteria and are the same as recommended for neutropenia above.		
Diarrhea 1 (2-3 stools/day > pretx <sup>c</sup> ) 2 (4-6 stools/day > pretx <sup>c</sup> ) 3 (7-9 stools/day > pretx <sup>c</sup> ) 4 ( $\geq$ 10 stools/day > pretx <sup>c</sup> )	Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> Omit dose, then $\downarrow$ 25 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2 Omit dose, then $\downarrow$ 50 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2	Maintain dose level Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>	Maintain dose level Maintain dose level $\downarrow$ 50 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>
Other nonhematologic toxicities <sup>d</sup> 1 2 3 4	Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> Omit dose, then $\downarrow$ 25 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2 Omit dose, then $\downarrow$ 50 mg/m <sup>2</sup> when resolved to $\leq$ Grade 2	Maintain dose level $\downarrow$ 25 mg/m <sup>2</sup> $\downarrow$ 25 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>	Maintain dose level $\downarrow$ 50 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup> $\downarrow$ 50 mg/m <sup>2</sup>

- a. All dose modifications should be based on the worst preceding toxicity
- b. National Cancer Institute Common Toxicity Criteria
- c. Pretreatment
- d. Excludes alopecia, anorexia, asthenia

**Table 4. Recommended Dose Modifications for Irinotecan/5-Fluorouracil/Leucovorin Combination Schedules**

Patients should return to pre-treatment bowel function without requiring antidiarrhea medications for at least 24 hours before the next chemotherapy administration. A new cycle of therapy should not begin until the granulocyte count has recovered to $\geq 1500/\text{mm}^3$ , and the platelet count has recovered to $\geq 100,000/\text{mm}^3$ , and treatment-related diarrhea is fully resolved. Treatment should be delayed 1 to 2 weeks to allow for recovery from treatment-related toxicities. If the patient has not recovered after a 2-week delay, consideration should be given to discontinuing irinotecan.		
Toxicity NCI Grade <sup>b</sup> (Value)	During a Cycle of Therapy	At the Start of Subsequent Cycles of Therapy
No toxicity	Maintain dose level	Maintain dose level
Neutropenia 1 (1500 to 1999/ $\text{mm}^3$ ) 2 (1000 to 1499/ $\text{mm}^3$ ) 3 (500 to 999/ $\text{mm}^3$ ) 4 (< 500/ $\text{mm}^3$ )  Neutropenic fever (Grade 4 neutropenia & $\geq$ Grade 2 fever)	Maintain dose level <sup>c</sup> $\downarrow$ 1 dose level <sup>d</sup> Omit dose, then $\downarrow$ 1 dose level when resolved to $\leq$ Grade 2 Omit dose, then $\downarrow$ 2 dose levels when resolved to $\leq$ Grade 2 <sup>d</sup> Omit dose, then $\downarrow$ 2 dose levels when resolved	Maintain dose level <sup>c</sup> Maintain dose level $\downarrow$ 1 dose level <sup>d</sup> $\downarrow$ 2 dose levels  $\downarrow$ 2 dose levels
Other hematologic toxicities	Dose modifications for leukopenia or thrombocytopenia during a cycle of therapy and at the start of subsequent cycles of therapy are also based on NCI toxicity criteria and are the same as recommended for neutropenia above.	
Diarrhea 1 (2-3 stools/day > pret <sup>e</sup> ) 2 (4-6 stools/day > pret <sup>e</sup> ) 3 (7-9 stools/day > pret <sup>e</sup> ) 4 ( $\geq$ 10 stools/day > pret <sup>e</sup> )	Delay dose until resolved to baseline (bsl), then give same dose Omit dose, then $\downarrow$ 1 dose level when resolved to bsl Omit dose, then $\downarrow$ 1 dose level when resolved to bsl Omit dose, then $\downarrow$ 2 dose levels when resolved to bsl	Maintain dose level  Maintain dose level $\downarrow$ 1 dose level $\downarrow$ 2 dose levels
Other nonhematologic Toxicities <sup>f</sup>  1 2 3 4	Maintain dose level Omit dose, then $\downarrow$ 1 dose level when resolved to $\leq$ Grade 1 Omit dose, then $\downarrow$ 1 dose level when resolved to $\leq$ Grade 2 Omit dose, then $\downarrow$ 2 dose levels when resolved to $\leq$ Grade 2  For mucositis/stomatitis decrease only 5-FU, not irinotecan <sup>g</sup>	Maintain dose level Maintain dose level $\downarrow$ 1 dose level $\downarrow$ 2 dose levels  For mucositis/stomatitis decrease only 5-FU, not irinotecan. <sup>g</sup>

- a. Dose modification refers to irinotecan and 5-FU; LV dose remains fixed at 20 mg/m<sup>2</sup> (not adjusted).
- b. National Cancer Institute Common Toxicity Criteria
- c. Refers to initial dose used in previous cycle.
- d. Irinotecan: dose level reductions = 25 mg/m<sup>2</sup> decrements; 5-Fluorouracil: dose level reductions = 100 mg/m<sup>2</sup> decrements
- e. Pretreatment
- f. Excludes alopecia, anorexia, asthenia
- g. For mucositis/stomatitis decrease only 5-FU, not irinotecan.

**Table 5. Dose Modifications at the Start of a New Cycle of the Cisplatin and Irinotecan (mg/m<sup>2</sup>) Combination Schedule - Based on the Worst Toxicity Observed in the Prior Cycle**

A new cycle of therapy should not begin until the granulocyte count has recovered to  $\geq 1500/\text{mm}^3$ , and the platelet count has recovered to  $\geq 100,000/\text{mm}^3$ , and treatment-related diarrhea is fully resolved. Treatment should be delayed 1 to 2 weeks to allow for recovery from treatment-related toxicities. If the patient has not recovered after a 2-week delay, consideration should be given to discontinuing irinotecan.

Toxicity NCI Grade <sup>a</sup>	Cisplatin <sup>b</sup>	Irinotecan <sup>c</sup>
<b>Hematologic</b>		
Grade 0, 1, 2, or 3	Maintain dose level	Maintain dose level
Grade 4	↓ 1 dose level	↓ 1 dose level
Febrile neutropenia, <sup>d</sup> sepsis, thrombocytopenia requiring transfusion	↓ 1 dose level	↓ 1 dose level
<b>Non-Hematologic</b>		
Diarrhea		
Grade 0, 1 or 2	Maintain dose level	Maintain dose level
Grade 3 or 4	Maintain dose level	↓ 1 dose level
Vomiting		
Grade 0, 1 or 2	Maintain dose level	Maintain dose level
Grade 3	↓ 1 dose level	Maintain dose level
Grade 4	↓ 1 dose level	↓ 1 dose level
Serum Creatinine		
< 1.5 mg/dL	Maintain dose level	Maintain dose level
1.5-2.0 mg/dL	↓ 2 dose levels	Maintain dose level
> 2.0 mg/dL	Omit dose	Maintain dose level
Ototoxicity		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	↓ 1 dose level	Maintain dose level
Grade 3 or 4	Discontinue cisplatin	Maintain dose level
Neurotoxicity		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	↓ 1 dose level	Maintain dose level
Grade 3 or 4	Discontinue cisplatin	Maintain dose level
<b>Other Non-Hematologic Toxicities</b>		
Grade 0, 1 or 2	Maintain dose level	Maintain dose level
Grade 3 or 4	↓ 1 dose level	↓ 1 dose level

a. National Cancer Institute Common Toxicity Criteria

b. Cisplatin: dose level reductions = 7.5 mg/m<sup>2</sup> decrements

c. Irinotecan: dose level reductions = 10 mg/m<sup>2</sup> decrements

d. Febrile neutropenia is defined as in CTC version 2: temperature  $\geq 38.5^\circ\text{C}$  concomitant with an ANC  $< 1.0 \times 10^9/\text{L}$ .

**Table 6. Dose Modifications During a Cycle of the Cisplatin and Irinotecan (mg/m<sup>2</sup>) Combination Schedule Based on Worst Toxicity Observed Since the Start of Cycle**

<b>Toxicity NCI Grade<sup>a</sup></b>	<b>Cisplatin<sup>b</sup></b>	<b>Irinotecan<sup>c</sup></b>
<b>Hematologic</b>		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	↓ 1 dose level	↓ 1 dose level
Grade 3	↓ 2 dose levels	↓ 2 dose levels
Grade 4	Omit dose	Omit dose
Febrile neutropenia, <sup>d</sup> sepsis, thrombocytopenia requiring transfusion	Omit dose	Omit dose
<b>Non-Hematologic</b>		
<b>Diarrhea</b>		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	Maintain dose level	↓ 1 dose level
Grade 3	Maintain dose level	Omit dose
Grade 4	Omit dose	Omit dose
<b>Vomiting</b>		
Grade 0, 1 or 2	Maintain dose level	Maintain dose level
Grade 3	↓ 1 dose level	Maintain dose level
Grade 4	↓ 1 dose level	↓ 1 dose level
<b>Serum Creatinine</b>		
< 1.5 mg/dL	Maintain dose level	Maintain dose level
1.5-2.0 mg/dL	↓ 2 dose levels	Maintain dose level
> 2.0 mg/dL	Omit dose	Maintain dose level
<b>Ototoxicity</b>		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	↓ 1 dose level	Maintain dose level
Grade 3 or 4	Discontinue cisplatin	Maintain dose level
<b>Neurotoxicity</b>		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2	↓ 1 dose level	Maintain dose level
Grade 3 or 4	Discontinue cisplatin	Maintain dose level
<b>Other Non-Hematologic Toxicities</b>		
Grade 0 or 1	Maintain dose level	Maintain dose level
Grade 2, 3 or 4	Omit dose	Omit dose

a. National Cancer Institute Common Toxicity Criteria

b. Cisplatin: dose level reductions = 7.5 mg/m<sup>2</sup> decrements

c. Irinotecan: dose level reductions = 10 mg/m<sup>2</sup> decrements

d. Febrile neutropenia is defined as in CTC version 2: temperature ≥ 38.5°C concomitant with an ANC < 1.0 x 10<sup>9</sup>/L.

### 4.3. Contraindications

Irinotecan is contraindicated in patients with a known hypersensitivity to the drug or its excipients (see Section 4.4 Special warnings and precautions for use).



#### **4.4. Special warnings and precautions for use**

##### **Administration**

Irinotecan should be administered only under the supervision of a physician who is experienced in the use of cancer chemotherapeutic agents. Appropriate management of complications is possible only when adequate diagnostic and treatment facilities are readily available.

Irinotecan will only be prescribed in the following cases after the expected benefits have been weighted against the possible therapeutic risks:

- in patients presenting a risk factor, particularly those with a WHO performance status = 2.
- in the few rare instances where patients are deemed unlikely to observe recommendations regarding management of adverse events (need for immediate and prolonged antidiarrheal treatment combined with high fluid intake at onset of delayed diarrhea). Strict hospital supervision is recommended for such patients.

##### **Cholinergic symptoms**

Patients may have cholinergic symptoms of rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing (vasodilation), bradycardia, and intestinal hyperperistalsis that can cause abdominal cramping and early diarrhea (i.e., diarrhea generally occurring during or within 8 hours of administration of irinotecan). These symptoms may be observed during or shortly after infusion of irinotecan, are thought to be related to the anticholinesterase activity of the irinotecan parent compound, and are expected to occur more frequently with higher irinotecan doses. Therapeutic or prophylactic administration of 0.25 to 1 mg of intravenous or subcutaneous atropine should be considered (unless clinically contraindicated) in patients experiencing cholinergic symptoms.

##### **Extravasation**

While irinotecan is not a known vesicant, care should be taken to avoid extravasation, and the infusion site should be monitored for signs of inflammation. Should extravasation occur, flushing the site, and application of ice is recommended.

##### **Hepatic**

In clinical studies, National Cancer Institute (NCI) Common Toxicity Criteria Grade 3 or 4 liver enzyme abnormalities have been observed in fewer than 10% of patients. These events typically occur in patients with known hepatic metastases and are not clearly related to irinotecan.

##### **Hematology**



Irinotecan commonly causes neutropenia, leukopenia, and anemia, any of which may be severe and therefore should not be used in patients with severe bone marrow failure. Serious thrombocytopenia is uncommon. In clinical studies, the frequency of NCI Grade 3 and 4 neutropenia has been significantly higher in patients who received previous pelvic/abdominal irradiation than in those who had not received such irradiation. Patients with baseline serum total bilirubin levels of 1.0 mg/dL or more also have had a significantly greater likelihood of experiencing first-cycle Grade 3 or 4 neutropenia than those with bilirubin levels that were less than 1.0 mg/dL. There were no significant differences in the frequency of Grade 3 and 4 neutropenia by age or gender (see Section 4.2 Posology and method of administration).

Neutropenic fever (concurrent NCI Grade 4 neutropenia and  $\geq$  Grade 2 fever) occurred in fewer than 10% of patients in clinical studies; however, deaths due to sepsis following severe neutropenia have been reported in patients treated with irinotecan. Neutropenic complications should be managed promptly with antibiotic support. Therapy with irinotecan should be temporarily discontinued if neutropenic fever occurs or if the absolute neutrophil count drops below  $1000/\text{mm}^3$ . The dose of irinotecan should be reduced if clinically significant neutropenia occurs (see Section 4.2 Posology and method of administration).

#### **Patients with reduced UGT1A1 activity**

The metabolic conversion of irinotecan to the active metabolite SN-38 is mediated by carboxylesterase enzymes and primarily occurs in the liver. SN-38 subsequently undergoes conjugation to form the inactive glucuronide metabolite SN-38G. This glucuronidation reaction is mediated primarily by uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1), which is encoded by the UGT1A1 gene (see Section 5.2 Pharmacokinetic properties). The UGT1A1 gene is highly polymorphic, resulting in variable metabolic capacities among individuals. One specific variation of the UGT1A1 gene includes a polymorphism in the promoter region known as the UGT1A1 28 variant allele. This variant and other congenital deficiencies in UGT1A1 expression (such as Crigler-Najjar and Gilbert's syndrome) are associated with reduced enzyme activity and increased systemic exposure to SN-38. Higher plasma concentrations of SN-38 are observed in individuals who are homozygous for the UGT1A1\*28 allele (also referred to as UGT1A1 7/7 genotype) versus patients who have one or two wild-type alleles.

Data from a meta-analysis of nine studies involving a total of 821 patients indicate that individuals with Crigler-Najjar syndrome (types 1 and 2) or those who are homozygous for the UGT1A1\*28 allele (Gilbert's syndrome) are at increased risk of hematological toxicity (Grades 3 and 4) following administration of irinotecan at moderate or high doses ( $>150 \text{ mg}/\text{m}^2$ ). A relationship between UGT1A1 genotype and the occurrence of irinotecan induced diarrhea was not established.

Patients known to be homozygous for UGT1A1\*28 should be administered the normally indicated irinotecan starting dose. However, these patients should be monitored for hematologic toxicities. A reduced irinotecan starting dose should be considered for patients who have experienced prior hematologic toxicity with previous treatment. The exact



reduction in starting dose in this patient population has not been established and any subsequent dose modifications should be based on individual patient tolerance to treatment.

### **Hypersensitivity reactions**

Hypersensitivity reactions, including severe anaphylactic/anaphylactoid reactions, have been reported.

### **Immunosuppressant effects/increased susceptibility to infections**

Administration of live or live-attenuated vaccines in patients immunocompromised by chemotherapeutic agents including irinotecan, may result in serious or fatal infections. Vaccination with a live vaccine should be avoided in patients receiving irinotecan. Killed or inactivated vaccines may be administered; however, the response to such vaccines may be diminished.

### **Late diarrhea**

Late diarrhea (generally occurring more than 8 hours after administration of irinotecan) can be prolonged, may lead to dehydration, electrolyte imbalance, or sepsis and may be life-threatening. In the clinical studies testing the every 3-week-dosage schedule, the median time to the onset of late diarrhea was 5 days after irinotecan infusion. In the clinical studies evaluating the weekly dosage schedule, the median time to onset of late diarrhea was 11 days following administration of irinotecan. For patients starting treatment at the 125 mg/m<sup>2</sup> weekly dose, the median duration of any grade of late diarrhea was 3 days. Among those patients treated at the 125 mg/m<sup>2</sup> weekly dose who experienced Grade 3 or 4 late diarrhea, the median duration of the entire episode of diarrhea was 7 days. Results from a prospective study of the weekly dosage schedule did not demonstrate any difference in the rate of late onset diarrhea in patients ≥65 years of age than patients <65 years of age. However, patients ≥65 years of age should be closely monitored due to a greater risk of early diarrhea observed in this population. Colonic ulceration, sometimes with bleeding, has been observed in association with irinotecan-induced diarrhea.

Late diarrhea should be treated promptly with loperamide at the first episode of poorly formed or loose stools or the earliest onset of bowel movements more frequent than normally expected for the patient. The recommended dosage regimen for loperamide is 4 mg at the first onset of late diarrhea and then 2 mg every 2 hours until the patient is diarrhea-free for at least 12 hours. During the night, the patient may take 4 mg of loperamide every 4 hours. Loperamide is not recommended to be used for more than 48 consecutive hours at these doses, because of the risk of paralytic ileus, nor for less than 12 hours. Premedication with loperamide is not recommended. Patients with diarrhea should be carefully monitored and given fluid and electrolyte replacement if they become dehydrated and should be given antibiotic support if they develop ileus, fever, or severe neutropenia. In addition to the antibiotic treatment, hospitalization is recommended for management of the diarrhea, in the following cases:



- Diarrhea associated with fever,
- Severe diarrhea (requiring intravenous hydration),
- Patients with vomiting associated with delayed (i.e., late) diarrhea,
- Diarrhea persisting beyond 48 hours following the initiation of high-dose loperamide therapy.

After the first treatment, subsequent weekly chemotherapy treatments should be delayed in patients until return of pre-treatment bowel function for at least 24 hours without need for anti-diarrhea medication. If NCI Grade 2, 3, or 4 diarrhea occurs, subsequent doses of irinotecan should be reduced within the current cycle (see Section 4.2 Posology and method of administration).

### **Chronic inflammatory bowel disease and/or bowel obstruction**

Patients must not be treated with irinotecan until resolution of the bowel obstruction.

### **Nausea & vomiting**

Irinotecan is emetogenic. Nausea and vomiting can be severe and usually occurs during or shortly after infusion of irinotecan. It is recommended that patients receive premedication with antiemetic agents. Antiemetic agents should be given on the day of treatment, starting at least 30 minutes before administration of irinotecan. Physicians should also consider providing patients with an antiemetic regimen for subsequent use as needed. Patients with vomiting associated with delayed (i.e., late) diarrhea should be hospitalized as soon as possible for treatment.

### **Neurologic**

Dizziness has been observed and may sometimes represent symptomatic evidence of orthostatic hypotension in patients with dehydration.

### **Renal**

Increases in serum creatinine or blood urea nitrogen have been observed. There have been cases of acute renal failure. These events have generally been attributed to complications of infection or to dehydration related to nausea, vomiting, or diarrhea. Rare instances of renal dysfunction due to tumor lysis syndrome have also been reported.

### **Respiratory**

NCI Grade 3 or 4 dyspnea has been observed. The extent to which malignant pulmonary involvement or other preexisting lung disease may have contributed to dyspnea is unknown. A potentially life-threatening pulmonary syndrome, consisting of dyspnea, fever, and a

---



reticulonodular pattern on chest x-ray, was observed in a small percentage of patients in early Japanese studies. The contribution of irinotecan to these preliminary events was difficult to assess because these patients also had lung tumors and some had preexisting nonmalignant pulmonary disease.

Interstitial pulmonary disease presenting as pulmonary infiltrates is uncommon during irinotecan therapy. Interstitial pulmonary disease can be fatal. Risk factors possibly associated with the development of interstitial pulmonary disease include pre-existing lung disease, use of pneumotoxic drugs, radiation therapy, and colony stimulating factors. Patients with risk factors should be closely monitored for respiratory symptoms before and during irinotecan therapy.

### **Others**

Since this product contains sorbitol, it is unsuitable in hereditary fructose intolerance.

### **Special populations**

#### Pediatric

The effectiveness of irinotecan in pediatric patients has not been established (see Section 5.2 Pharmacokinetic properties). Results from two open-label, single arm studies were evaluated. One hundred and seventy children with refractory solid tumors were enrolled in one phase 2 trial in which 50 mg/m<sup>2</sup> of irinotecan was infused for 5 consecutive days every 3 weeks. Grade 3-4 neutropenia was experienced by 54 (31.8%) patients. Neutropenia was complicated by fever in 15 (8.8%) patients. Grade 3-4 diarrhea was observed in 35 (20.6%) patients. This adverse event profile was comparable to that observed in adults.

In the second phase 2 trial of 21 children with previously untreated rhabdomyosarcoma, 20 mg/m<sup>2</sup> of irinotecan was infused for 5 consecutive days on weeks 0, 1, 3, and 4. This single agent therapy was followed by multimodal therapy. Accrual to the single agent irinotecan phase was halted due to the high rate (28.6%) of progressive disease and the early deaths (14%). The adverse event profile was different in this study from that observed in adults; the most significant Grade 3 or 4 adverse events were dehydration experienced by 6 patients (28.6%) associated with severe hypokalaemia in 5 patients (23.8%) and hyponatremia in 3 patients (14.3%); in addition, Grade 3-4 infection was reported in 5 patients (23.8%) (across all courses of therapy and irrespective of causal relationship).

#### Geriatric

Specific dosing recommendations may apply to this population depending upon the regimen used (see Section 4.2 Posology and method of administration).

#### Hepatic insufficiency



In patients with hyperbilirubinemia, the clearance of irinotecan is decreased (see Section 5.2 Pharmacokinetic properties) and therefore the risk of hematotoxicity is increased. The use of irinotecan in patients with a serum total bilirubin concentration of  $>3.0 \times$  institutional upper limit of normal (IULN) given as a single-agent on the once-every-3-weeks schedule has not been established (see Section 4.2 Posology and method of administration). Liver function should be monitored before initiation of treatment and monthly, or as clinically indicated.

#### Irradiation therapy

Patients who have previously received pelvic/abdominal irradiation are at increased risk of myelosuppression following the administration of irinotecan. Physicians should use caution in treating patients with extensive prior irradiation. Specific dosing recommendations may apply to this population depending upon the regimen used (see Section 4.2 Posology and method of administration).

#### **Performance status**

Patients with poor performance status are at increased risk of irinotecan-related adverse events. Specific dosing recommendations for patients with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 may apply depending upon the regimen used (see Section 4.2 Posology and method of administration). Patients with performance status of 3 or 4 should not receive irinotecan. In patients receiving either irinotecan/5-FU/LV or 5-FU/LV in clinical trials comparing these agents, higher rates of hospitalisation, neutropenic fever, thromboembolism, first-cycle treatment discontinuation, and early deaths were observed in patients with a baseline performance status of 2 than in patients with a baseline performance status of 0 or 1.

#### **Gastric cancer**

Patients with gastric cancer appear to experience greater myelosuppression and other toxicities when given irinotecan. A lower starting dose should be considered in these patients (see Section 4.2 Posology and method of administration).

### **4.5. Interaction with other medicinal products and other forms of interaction**

#### **CYP3A4 and/or UGT1A1 inhibitors**

Irinotecan and active metabolite SN-38 are metabolized via the human cytochrome P450 3A4 isoenzyme (CYP3A4) and uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1) (see Section 5.2 Pharmacokinetic properties). Coadministration of irinotecan with inhibitors of CYP3A4 and/or UGT1A1 may result in increased systemic exposure to irinotecan and the active metabolite SN-38. Physicians should take this into consideration when administering irinotecan with these drugs.

#### Ketoconazole



Irinotecan clearance is greatly reduced in patients receiving concomitant ketoconazole, leading to increased exposure to SN-38. Ketoconazole should be discontinued at least 1 week prior to starting irinotecan therapy and should not be administered during irinotecan therapy.

#### Atazanavir sulfate

Coadministration of atazanavir sulfate, a CYP3A4 and UGT1A1 inhibitor has the potential to increase systemic exposure to SN-38, the active metabolite of irinotecan. Physicians should take this into consideration when coadministering these drugs.

#### **CYP3A4 inducers**

##### Anticonvulsants

Concomitant administration of CYP3A-inducing anticonvulsant drugs (e.g., carbamazepine, phenobarbital or phenytoin) leads to reduced exposure to the active metabolite SN-38. Consideration should be given to starting or substituting non-enzyme-inducing anticonvulsants at least one week prior to initiation of irinotecan therapy in patients requiring anticonvulsant treatment.

##### St. John's Wort (*Hypericum perforatum*)

Exposure to the active metabolite SN-38 is reduced in patients taking concomitant St. John's Wort. St. John's Wort should be discontinued at least 1 week prior to the first cycle of irinotecan, and should not be administered during irinotecan therapy.

#### **Other interactions**

##### Neuromuscular blocking agents

Interaction between irinotecan and neuromuscular blocking agents cannot be ruled out, since irinotecan has anticholinesterase activity. Drugs with anticholinesterase activity may prolong the neuromuscular blocking effects of suxamethonium and the neuromuscular blockade of non-depolarizing drugs may be antagonized.

##### Antineoplastic agents

The adverse effects of irinotecan, such as myelosuppression and diarrhea, would be expected to be exacerbated by other antineoplastic agents having a similar adverse-effect profile.

##### Dexamethasone

Lymphocytopenia has been reported in patients receiving irinotecan, and it is possible that the administration of dexamethasone as antiemetic prophylaxis may have enhanced the



likelihood of lymphocytopenia. However, serious opportunistic infections have not been observed and no complications have specifically been attributed to lymphocytopenia.

Hyperglycemia has been observed in patients with a history of diabetes mellitus or evidence of glucose intolerance prior to administration of irinotecan. It is probable that dexamethasone, given as antiemetic prophylaxis, contributed to hyperglycemia in some patients.

#### Laxatives

Laxative use during therapy with irinotecan is expected to worsen the incidence or severity of diarrhea.

#### Diuretics

Dehydration secondary to vomiting and/or diarrhea may be induced by irinotecan. The physician may wish to withhold diuretics during dosing with irinotecan and during periods of active vomiting or diarrhea.

#### Bevacizumab

Results from a dedicated drug-drug interaction trial demonstrated no significant effect of bevacizumab on the pharmacokinetics of irinotecan and its active metabolite SN-38.

### **4.6. Fertility, pregnancy and lactation**

#### **Pregnancy**

Irinotecan is teratogenic in rats and rabbits (see Section 5.3 Preclinical safety data). Irinotecan may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of irinotecan in pregnant women. If the drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with irinotecan.

#### **Lactation**

In rats, radioactivity appeared in the milk within 5 minutes of intravenous administration of radiolabeled irinotecan and was concentrated up to 65-fold at 4 hours after administration relative to plasma concentrations. Because many drugs are excreted in human milk, and because of the potential for serious adverse reactions in nursing infants, it is recommended that nursing be discontinued when receiving therapy with irinotecan.



#### **4.7. Effects on ability to drive and use machines**

The effect of irinotecan on the ability to drive or use machinery has not been evaluated. However, patients should be warned about the potential for dizziness or visual disturbances which may occur within 24 hours following the administration of irinotecan, and advised not to drive or operate machinery if these symptoms occur (see Section 4.4 Special warnings and precautions for use).

#### **4.8. Undesirable effects**

##### **Clinical studies**

Adverse reaction data has been extensively collected and analyzed for the clinical studies program in metastatic colorectal cancer that recurred or progressed following 5-FU-based therapy (second-line) and are presented below (patient population described below). The adverse reactions for other indications are expected to be similar to those for second-line colorectal cancer.

Adverse reactions detailed in this section refer to irinotecan. There is no evidence that the safety profile of irinotecan is influenced by cetuximab or vice versa. In combination with cetuximab, additional reported adverse reactions were those expected with cetuximab (such as acneform rash). Therefore, also refer to the full prescribing information for cetuximab.

Grade 3 hypertension was the principal significant risk involved with the addition of bevacizumab to bolus irinotecan/5-FU/FA. In addition, there was a small increase in the Grade 3/4 chemotherapy adverse events of diarrhea and leukopenia with this regimen compared to patients receiving bolus irinotecan/5-FU/FA alone. For other information on adverse reactions in combination with bevacizumab, refer to the bevacizumab full prescribing information.

Adverse drug reactions reported in patients treated with the combination irinotecan plus capecitabine in addition to those seen with capecitabine monotherapy or seen at a higher frequency grouping with the combination compared to capecitabine monotherapy include:

*Very common, all grade:* thrombosis/embolism

*Common, all grade:* hypersensitivity reaction, cardiac ischemia/infarction

*Common, Grade 3 and Grade 4:* febrile neutropenia

For complete information on adverse reactions of capecitabine, refer to the capecitabine full prescribing information.

Grade 3 and Grade 4 adverse drug reactions reported in patients treated with irinotecan in combination with capecitabine and bevacizumab in addition to those seen with capecitabine



monotherapy or seen at a higher frequency grouping with the combination compared to capecitabine monotherapy include:

*Common, Grade 3 and Grade 4:* neutropenia, thrombosis/embolism, hypertension, and cardiac ischemia/infarction

For complete information on adverse reactions of capecitabine and bevacizumab, refer to the respective capecitabine and bevacizumab full prescribing information.

### **Clinical studies of the 100- to 125-mg/m<sup>2</sup> single-agent weekly dosage schedule**

The weekly dosage schedule of irinotecan was evaluated in three clinical studies of 304 patients with metastatic carcinoma of the colon or rectum that had recurred or progressed following 5-FU-based therapy. Five (1.6%) deaths were potentially drug-related. These five patients experienced a constellation of medical events (myelosuppression, neutropenic sepsis without fever, small bowel obstruction, fluid accumulation, stomatitis, nausea, vomiting, diarrhea, and dehydration) that are known effects of irinotecan. Neutropenic fever, defined as NCI Grade 4 neutropenia and Grade 2 or greater fever, occurred in nine other patients; these patients recovered with supportive care.

Eighty-one (26.6%) patients were hospitalized for events judged to be related to administration of irinotecan. The primary reasons for drug-related hospitalization were diarrhea, with or without nausea and/or vomiting; neutropenia/leukopenia, with or without diarrhea and/or fever; and nausea and/or vomiting.

Adjustments in the dose of irinotecan were made during the cycle of treatment and for subsequent cycles based on individual patient tolerance. The most common reasons for dose reduction were late diarrhea, neutropenia, and leukopenia. Thirteen (4.3%) patients discontinued treatment with irinotecan because of adverse events.

### **Clinical studies of the 300- to 350-mg/m<sup>2</sup> once-every-3-week single-agent dosage schedule**

A total of 316 patients with metastatic colorectal cancer whose disease had progressed following prior 5-FU therapy received irinotecan in two studies involving once-every-3-week administration. Three (1%) deaths were potentially related to irinotecan treatment and were attributed to neutropenic infection, Grade 4 diarrhea, and asthenia, respectively. Hospitalizations due to serious adverse events, whether or not related to irinotecan administration, occurred at least once in 60% of patients who received irinotecan and, 8% of patients treated with irinotecan discontinued treatment due to adverse events.

### **Listing of adverse events**



The drug-related adverse events (NCI Grades 1- 4) as judged by the investigator that were reported in greater than 10% of the 304 patients enrolled in the three studies of the weekly dosage schedule are listed by body system in descending order of frequency in Table 7.

**Table 7. NCI Grade 1 to 4 Drug-Related Adverse Events Observed in over 10% of Patients in Clinical Studies**

<b>Gastrointestinal disorders:</b>	Late diarrhea, nausea, vomiting, early diarrhea, abdominal cramping/pain, anorexia, stomatitis
<b>Blood and lymphatic system disorders:</b>	Leukopenia, anemia, neutropenia
<b>General disorders and administration site conditions:</b>	Asthenia, fever
<b>Metabolism and nutrition disorders:</b>	Decreased weight, dehydration
<b>Skin and subcutaneous tissue disorders:</b>	Alopecia
<b>Vascular disorders:</b>	Thromboembolic events*

\*Includes angina pectoris, arterial thrombosis, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, embolus lower extremity, heart arrest, myocardial infarct, myocardial ischemia, peripheral vascular disorder, pulmonary embolus, sudden death, thrombophlebitis, thrombosis, vascular disorder.

NCI Grade 3 or 4 adverse events reported in the clinical studies of the weekly and once-every-3-week-dosage schedules (N=620) are listed in Table 8, Table 9, and Table 10.

**Table 8. NCI Grade 3 or 4 Drug-Related Adverse Events Observed in over 10% of Patients in Clinical Studies**

<b>Gastrointestinal disorders:</b>	Late diarrhea, nausea, abdominal cramping/pain
<b>Blood and lymphatic system disorders:</b>	Leukopenia, neutropenia
<b>Skin and subcutaneous tissue disorders:</b>	Alopecia



**Table 9. NCI Grade 3 or 4 Drug-Related Adverse Events Observed in 1% to 10% of Patients in Clinical Studies**

<b>Infections and infestations:</b>	Infection
<b>Gastrointestinal disorders:</b>	Vomiting, early diarrhea, constipation, anorexia, mucositis
<b>Blood and lymphatic system disorders:</b>	Anemia, thrombocytopenia
<b>General disorders and administration site conditions:</b>	Asthenia, fever, pain
<b>Metabolism and nutrition disorders:</b>	Dehydration, hypovolemia
<b>Hepatobiliary disorders:</b>	Bilirubinemia
<b>Respiratory, thoracic and mediastinal disorders:</b>	Dyspnea
<b>Investigations:</b>	Increased creatinine



**Table 10. NCI Grade 3 or 4 Drug-Related Adverse Events Observed in Fewer than 1% of Patients in Clinical Studies**

<b>Infections and infestations:</b>	Sepsis
<b>Gastrointestinal disorders:</b>	Rectal disorder, GI monilia
<b>General disorders and administration site conditions:</b>	Chills, malaise
<b>Metabolism and nutrition disorders:</b>	Decreased weight, hypokalemia, hypomagnesemia
<b>Skin and subcutaneous tissue disorders:</b>	Rash, cutaneous signs
<b>Nervous system disorders:</b>	Abnormal gait, confusion, headache
<b>Cardiac disorders:</b>	Hypotension, syncope, cardiovascular disorders
<b>Renal and urinary disorders:</b>	Urinary tract infection
<b>Reproductive system and breast disorders:</b>	Breast pain
<b>Investigations:</b>	Increased alkaline phosphatase, increased GGTP

The following additional drug-related events have been reported in clinical studies with irinotecan, but do not meet the criteria as defined above as either >10% drug-related NCI Grades 1-4 or as a NCI Grade 3 or 4 drug-related event: rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing, bradycardia, dizziness, extravasation, tumor lysis syndrome, and colonic ulceration.

### **Post-marketing surveillance**

#### Cardiac disorders

Myocardial ischemic events have been observed following irinotecan therapy predominantly in patients with underlying cardiac disease, other known risk factors for cardiac disease or previous cytotoxic chemotherapy (see also Table 7, thromboembolic events).

#### Gastrointestinal disorders



Infrequent cases of intestinal obstruction, ileus, megacolon, or gastrointestinal hemorrhage, and rare cases of colitis, including typhlitis, ischemic and ulcerative colitis were reported. In some cases, colitis was complicated by ulceration, bleeding, ileus, or infection. Cases of ileus without preceding colitis have also been reported. Rare cases of intestinal perforation were reported.

Rare cases of symptomatic pancreatitis or asymptomatic elevated pancreatic enzymes have been observed.

#### Hypovolemia

There have been rare cases of renal impairment and acute renal failure, generally in patients who became infected and/or volume depleted from severe gastrointestinal toxicities.

Infrequent cases of renal insufficiency, hypotension or circulatory failure have been observed in patients who experienced episodes of dehydration associated with diarrhea and/or vomiting, or sepsis. Infections and infestations

Bacterial, fungal and viral infections have been reported.

#### Immune system disorders

Hypersensitivity reactions including severe anaphylactic or anaphylactoid reactions have been reported (see Section 4.4 Special warnings and precautions for use).

#### Musculoskeletal and connective tissue disorders

Early effects such as muscular contraction or cramps and paresthesia have been reported.

#### Nervous system disorders

Speech disorders, generally transient in nature, have been reported in patients treated with irinotecan; in some cases, the event was attributed to the cholinergic syndrome observed during or shortly after infusion of irinotecan.

#### Respiratory, thoracic and mediastinal disorders

Interstitial pulmonary disease presenting as pulmonary infiltrates is uncommon during irinotecan therapy. Early effects such as dyspnea have been reported (see Section 4.4 Special warnings and precautions for use). Hiccups have also been reported.

#### Investigations

Rare cases of hyponatremia mostly related with diarrhea and vomiting have been reported. Increases in serum levels of transaminases (i.e., AST and ALT) in the absence of progressive liver metastasis have been very rarely reported.



## 4.9. Overdose

Single doses of up to 750 mg/m<sup>2</sup> irinotecan have been given to patients with various cancers. The adverse events in these patients were similar to those reported with the recommended dosages and regimens. There have been reports of overdosage at doses up to approximately twice the recommended therapeutic dose, which may be fatal. The most significant adverse reactions reported were severe neutropenia and severe diarrhea. Maximum supportive care should be instituted to prevent dehydration due to diarrhea and to treat any infectious complications. There is no known antidote for overdosage of irinotecan.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1. Pharmacodynamic properties

#### Therapeutic class

Irinotecan hydrochloride is an antineoplastic agent of the topoisomerase I inhibitor class, clinically investigated as CPT-11. Irinotecan is a semisynthetic derivative of camptothecin, an alkaloid extract from plants such as *Camptotheca acuminata*, or is chemically synthesized.

#### Mechanism of action

Irinotecan and its active metabolite SN-38 bind to the topoisomerase I – DNA complex and prevent re-ligation of these single-strand breaks. Current research suggests that the cytotoxicity of irinotecan is due to double-strand DNA damage produced during DNA synthesis when replication enzymes interact with the ternary complex formed by topoisomerase I, DNA, and either irinotecan or SN-38.

Irinotecan serves as a water-soluble precursor of the lipophilic metabolite SN-38. SN-38 is formed from irinotecan by carboxylesterase-mediated cleavage of the carbamate bond between the camptothecin moiety and the dipiperidino side chain. SN-38 is approximately 1000 times as potent as irinotecan as an inhibitor of topoisomerase I purified from human and rodent tumor cell lines. In vitro cytotoxicity assays show that the potency of SN-38 relative to irinotecan varies from 2- to 2000-fold. However, the plasma area under the concentration versus time curve (AUC) values for SN-38 are 2% to 8% of irinotecan and SN-38 is 95% bound to plasma proteins compared to approximately 50% bound to plasma proteins for irinotecan. The precise contribution of SN-38 to the activity of irinotecan is thus unknown. Both irinotecan and SN-38 exist in an active lactone form and an inactive hydroxy acid anion form. A pH-dependent equilibrium exists between the two forms such that an acid pH promotes the formation of the lactone, while a more basic pH favors the hydroxy acid anion form.

#### Clinical studies

In combination therapy for the first-line treatment of metastatic colorectal carcinoma

---

*Irinotecan in Combination With Cetuximab in Treatment-naïve Patients*: EMR 62 202-013: This randomized study in patients with metastatic colorectal cancer who had not received prior treatment for metastatic disease compared the combination of cetuximab and irinotecan plus infusional 5-fluorouracil/folinic acid (5-FU/FA) (599 patients) to the same chemotherapy alone (599 patients). The proportion of patients with KRAS wild-type tumors from the patient population evaluable for KRAS status comprised 64%.

The efficacy data generated in this study are summarized in Table 11 below:

**Table 11. Efficacy Results from Study EMR 62 202-013**

Variable/statistic	Overall population		KRAS wild-type population	
	Cetuximab plus FOLFIRI (N=599)	FOLFIRI (N=599)	Cetuximab plus FOLFIRI (N=172)	FOLFIRI (N=176)
ORR % (95% CI)	46.9 (42.9, 51.0)	38.7 (34.8, 42.8)	59.3 (51.6, 66.7)	43.2 (35.8, 50.9)
p-value	0.0038		0.0025	
PFS				
HR (95% CI)	0.85 (0.726, 0.998)		0.68 (0.501, 0.934)	
p-value	0.0479		0.0167	

CI=confidence interval; FOLFIRI=irinotecan plus infusional 5-FU/FA; ORR=objective response rate (patients with complete response or partial response); PFS=progression-free survival time.

*Irinotecan in Combination with Bevacizumab*: A phase III randomized, double-blind, active-controlled clinical trial evaluated bevacizumab in combination with irinotecan/5-FU/FA as first-line treatment for metastatic carcinoma of the colon or rectum (Study AVF2107g). The addition of bevacizumab to the combination of irinotecan/5-FU/FA resulted in a statistically significant increase in overall survival. The clinical benefit, as measured by overall survival, was seen in all pre-specified patient subgroups, including those defined by age, sex, performance status, location of primary tumor, number of organs involved, and duration of metastatic disease. Refer also to the bevacizumab full prescribing information. The efficacy results of Study AVF2107g are summarized in Table 12 below.

**Table 12. Efficacy Results from Study AVF2107g**

	AVF2107g	
	Arm 1 irinotecan/5-FU/FA + Placebo	Arm 2 irinotecan/5-FU/FA + bevacizumab <sup>a</sup>
Number of Patients	411	402
Overall survival		
Median time (months)	15.6	20.3

	AVF2107g	
	Arm 1 irinotecan/5-FU/FA + Placebo	Arm 2 irinotecan/5-FU/FA + bevacizumab <sup>a</sup>
95% CI	14.29 – 16.99	18.46 – 24.18
Hazard ratio <sup>b</sup>	-	0.660
p-value	-	0.00004
Progression-free survival		
Median time (months)	6.2	10.6
Hazard ratio	-	0.54
p-value	-	<0.0001
Overall response rate		
Rate (%)	34.8	44.8
95% CI	30.2 – 39.6	39.9 – 49.8
p-value	-	0.0036
Duration of response		
Median time (months)	7.1	10.4
25–75 percentile (months)	4.7 – 11.8	6.7 – 15.0

a. 5 mg/kg every 2 weeks.

b. Relative to control arm.

CI=confidence interval; 5-FU=5-fluorouracil; FA=folinic acid

*Irinotecan in Combination with Capecitabine:* Data from a randomized, controlled Phase III study (CAIRO) support the use of capecitabine at a starting dose of 1000 mg/m<sup>2</sup> for 2 weeks every 3 weeks in combination with irinotecan for the first-line treatment of patients with metastatic colorectal cancer. Eight hundred twenty (820) patients were randomized to receive either sequential treatment (n=410) or combination treatment (n=410). Sequential treatment consisted of first-line treatment with capecitabine (1250 mg/m<sup>2</sup> twice daily for 14 days), second-line irinotecan (350 mg/m<sup>2</sup> on day 1), and third-line combination of capecitabine (1000 mg/m<sup>2</sup> twice daily for 14 days) with oxaliplatin (130 mg/m<sup>2</sup> on day 1). Combination treatment consisted of first-line treatment of capecitabine (1000 mg/m<sup>2</sup> twice daily for 14 days) combined with irinotecan (250 mg/m<sup>2</sup> on day 1) (XELIRI) and second-line capecitabine (1000 mg/m<sup>2</sup> twice daily for 14 days) plus oxaliplatin (130 mg/m<sup>2</sup> on day 1). All treatment cycles were administered at intervals of 3 weeks. In first-line treatment the



median progression-free survival in the intent-to-treat population was 5.8 months (95% CI, 5.1-6.2 months) for capecitabine monotherapy and 7.8 months (95% CI, 7.0-8.3 months) for XELIRI (p=0.0002).

Data from an interim analysis of a multicenter, randomized, controlled Phase II study (AIO KRK 0604) support the use of capecitabine at a starting dose of 800 mg/m<sup>2</sup> for 2 weeks every 3 weeks in combination with irinotecan and bevacizumab for the first-line treatment of patients with metastatic colorectal cancer. One hundred fifteen (115) patients were randomized to treatment with capecitabine combined with irinotecan (XELIRI) and bevacizumab: capecitabine (800 mg/m<sup>2</sup> twice daily for 2 weeks followed by a 7-day rest period), irinotecan (200 mg/m<sup>2</sup> as a 30-minute infusion on day 1 every 3 weeks), and bevacizumab (7.5 mg/kg as a 30 to 90-minute infusion on day 1 every 3 weeks); a total of 118 patients were randomized to treatment with capecitabine combined with oxaliplatin plus bevacizumab; capecitabine (1000 mg/m<sup>2</sup> twice daily for 2 weeks followed by a 7-day rest period), oxaliplatin (130 mg/m<sup>2</sup> as a 2-hour infusion on day 1 every 3 weeks), and bevacizumab (7.5 mg/kg as a 30 to 90-minute infusion on day 1 every 3 weeks). Progression-free survival at 6 months in the intent-to-treat population was 80% (XELIRI plus bevacizumab) versus 74% (XELOX plus bevacizumab). Overall response rate (complete response plus partial response) was 45% (XELOX plus bevacizumab) versus 47% (XELIRI plus bevacizumab).

#### In combination therapy for the second-line treatment of metastatic colorectal carcinoma

*Irinotecan in Combination With Cetuximab After Failure of Irinotecan-including Cytotoxic Therapy:* The efficacy of the combination of cetuximab with irinotecan was investigated in two clinical studies. A total of 356 patients with EGFR-expressing metastatic colorectal cancer who had recently failed irinotecan-including cytotoxic therapy and who had a minimum Karnofsky performance status of 60, but the majority of whom had a Karnofsky performance status of ≥80 received the combination treatment.

EMR 62 202-007: This randomized study compared the combination of cetuximab and irinotecan (218 patients) with cetuximab monotherapy (111 patients).

IMCL CP02-9923: This single arm open-label study investigated the combination therapy in 138 patients.

The efficacy data from these studies are summarized in Table 13 below:

**Table 13. Efficacy Results from Studies EMR 62 202-007 and IMCL CP02-9923**

Study	N	ORR		DCR		PFS (months)		OS (months)	
		n (%)	95%CI	n (%)	95%CI	Median	95%CI	Median	95%CI
<b>Cetuximab + irinotecan</b>									
EMR 62 202-007	218	50 (22.9)	17.5, 29.1	121 (55.5)	48.6, 62.2	4.1	2.8, 4.3	8.6	7.6, 9.6



Study	N	ORR	DCR	PFS (months)	OS (months)				
IMCL CP02- 9923	138	21 (15.2)	9.7, 22.3	84 (60.9)	52.2, 69.1	2.9	2.6, 4.1	8.4	7.2, 10.3
<b>Cetuximab</b>									
EMR 62 202-007	111	12 (10.8)	5.7, 18.1	36 (32.4)	23.9, 42.0	1.5	1.4, 2.0	6.9	5.6, 9.1

CI=confidence interval; DCR=disease control rate (patients with complete response, partial response, or stable disease for at least 6 weeks); ORR=objective response rate (patients with complete response or partial response); OS=overall survival time; PFS=progression-free survival

The efficacy of the combination of cetuximab with irinotecan was superior to that of cetuximab monotherapy, in terms of objective response rate (ORR), disease control rate (DCR) and progression-free survival (PFS). In the randomized trial, no effects on overall survival were demonstrated (hazard ratio 0.91, p=0.48).

## 5.2. Pharmacokinetic properties

### Absorption and distribution

After intravenous infusion in humans, irinotecan plasma concentrations decline in a multiexponential manner, with a mean terminal elimination half-life of about 6 hours. The mean terminal elimination half-life of the active metabolite SN-38 is about 10 hours. The half-lives of the lactone (active) forms of irinotecan and SN-38 are similar to those of total irinotecan and SN-38, as the lactone and hydroxy acid forms are in equilibrium.

Over the dose range of 50 to 350 mg/m<sup>2</sup>, the AUC of irinotecan increases linearly with dose; the AUC of SN-38 increases less than proportionally with dose. Maximum concentrations of the active metabolite SN-38 are generally seen within 1 hour following the end of a 90-minute infusion of irinotecan.

Irinotecan exhibits moderate plasma protein binding (30% to 68% bound). SN-38 is highly bound to human plasma proteins (approximately 95% bound). The plasma protein to which irinotecan and SN-38 predominantly binds is albumin.

### Metabolism & excretion

Irinotecan (CPT-11) is subject to extensive metabolic conversion by various enzyme systems, including esterases to form the active metabolite SN-38, and UGT1A1 mediating glucuronidation of SN-38 to form the inactive glucuronide metabolite SN-38G. Irinotecan (CPT-11) can also undergo CYP3A4-mediated oxidative metabolism to several pharmacologically inactive oxidation products, one of which can be hydrolyzed by carboxylesterase to release SN-38. UGT1A1 activity is reduced in individuals with genetic polymorphisms that lead to reduced enzyme activity such as the UGT1A1\*28 polymorphism (see Section 4.4 Special warnings and precautions for use). SN-38 glucuronide had 1/50 to 1/100 the activity of SN-38 in cytotoxicity assays using two cell lines in vitro. The



disposition of irinotecan has not been fully elucidated in humans. The urinary excretion of irinotecan is 11% to 20%; SN-38, <1%; and SN-38 glucuronide, 3%. The cumulative biliary and urinary excretion of irinotecan and its metabolites (SN-38 and SN-38 glucuronide) over a period of 48 hours following administration of irinotecan in two patients ranged from approximately 25% (100 mg/m<sup>2</sup>) to 50% (300 mg/m<sup>2</sup>).

### Pharmacokinetics in special populations

*Geriatric:* The pharmacokinetics of irinotecan administered using the weekly schedule was evaluated in a study of 183 patients that was prospectively designed to investigate the effect of age on irinotecan toxicity. Results from this trial indicate that there are no differences in the pharmacokinetics of irinotecan, SN-38, and SN-38 glucuronide in patients <65 years of age compared with patients ≥65 years of age. In a study of 162 patients that was not prospectively designed to investigate the effect of age, small (less than 18%) but statistically significant differences in dose-normalized irinotecan pharmacokinetic parameters in patients <65 years of age compared to patients ≥65 years of age were observed. Although dose-normalized AUC<sub>0-24</sub> for SN-38 in patients ≥65 years of age was 11% higher than in patients <65 years of age, this difference was not statistically significant.

*Pediatric:* See Section 4.4 Special warnings and precautions for use.

The pharmacokinetics of irinotecan and its major metabolites in the pediatric population was investigated in clinical trials conducted in the US and Europe. Overall, results and general conclusions regarding irinotecan pharmacokinetics were comparable in the US and European studies. Any differences in the findings between these studies are probably attributable to differences in the doses investigated (20 to 200 mg/m<sup>2</sup> and 200 to 720 mg/m<sup>2</sup> in the US and European studies, respectively) and the marked inter-patient variability in values determined for the pharmacokinetic parameters of irinotecan and SN-38.

### US studies

Pharmacokinetic parameters for irinotecan and SN-38 were determined in 2 pediatric solid-tumor trials at dose levels of 50 mg/m<sup>2</sup> (60-min infusion, n=48) and 125 mg/m<sup>2</sup> (90-min infusion, n=6). Irinotecan clearance (mean ± S.D.) was 17.3 ± 6.7 L/h/m<sup>2</sup> for the 50 mg/m<sup>2</sup> dose and 16.2 ± 4.6 L/h/m<sup>2</sup> for the 125 mg/m<sup>2</sup> dose, which is somewhat greater than in adults. Minimal accumulation of irinotecan and SN-38 was observed in children on daily dosing regimens [daily x 5 every 3 weeks or (daily x 5) x 2 weeks every 3 weeks]. A finding that dose-normalized SN-38 AUC values were comparable between adults and children was inconsistent with the increase in irinotecan clearance seen in the pediatric population and was probably reflective of the marked inter-patient variability (%CV values for SN-38 AUC were 84 to 120%). Indeed SN-38 exposure in pediatric patients was approximately 30% lower than in adults when comparison was made without regard to the variability of the data.

### European studies



The pharmacokinetics of irinotecan and its major metabolites was investigated in pediatric patients with solid tumors in a phase I study at dose levels of 200 to 720 mg/m<sup>2</sup> (2-hour infusion, n=77). Systemic exposure to irinotecan, SN-38, APC, and NPC was dose proportional. Pharmacokinetic parameters of irinotecan and its metabolites demonstrated marked inter-patient variability with values (mean ± S.D.) for irinotecan plasma clearance of 18 ± 8 L/h/m<sup>2</sup> and volume of distribution at steady state of 104 ± 84 L/m<sup>2</sup>. Irinotecan clearance was 26% lower in adolescents than in children and dose normalized SN-38 and SN-38G exposures were 52% and 105% higher in adolescents than in children, respectively. Irinotecan clearance was higher and dose normalized values for SN-38, SN-38G and APC exposure were lower in the pediatric than in the adult population.

A population pharmacokinetic analysis of irinotecan was performed in 83 children and adolescents with relapsed or refractory rhabdomyosarcoma, primitive neuroectodermal tumor (PNET) including medulloblastoma or neuroblastoma receiving 600 mg/m<sup>2</sup> irinotecan as a 1-hour infusion once every 3 weeks as part of a phase II study. Mean values for irinotecan clearance and AUC demonstrated large inter- and intra-individual variability and were similar to those determined at the same dose in the European phase I pediatric study.

*Gender:* The pharmacokinetics of irinotecan do not appear to be influenced by gender.

*Race:* The influence of race on the pharmacokinetics of irinotecan has not been evaluated.

*Hepatic Insufficiency:* (see Section 4.2 Posology and method of administration) Irinotecan clearance is diminished in patients with hepatic dysfunction while relative exposure to the active metabolite SN-38 is increased. The magnitude of these effects is proportional to the degree of liver impairment as measured by elevations in serum total bilirubin and transaminase concentrations.

*Renal Insufficiency:* The influence of renal insufficiency on the pharmacokinetics of irinotecan has not been evaluated (see Section 4.2 Posology and method of administration).

### 5.3. Preclinical safety data

#### Toxicology

The acute intravenous toxicity of irinotecan in animals is shown below in Table 14. Lethality was observed after single intravenous irinotecan doses of approximately 111 mg/kg in mice and 73 mg/kg in rats (approximately 2.6 and 3.4 times the recommended human dose of 125 mg/m<sup>2</sup>, respectively). Death was preceded by cyanosis, tremors, respiratory distress, and convulsions. Subacute toxicity studies show that irinotecan affects tissues with rapid cell proliferation (bone marrow, intestinal epithelia, thymus, spleen, lymph nodes, and testes).

**Table 14. Acute Toxicity of Irinotecan (IV) in Animals**

Species	LD50 (mg/kg)
---------	--------------



Mouse	132-134
Rat	84-85
Dog	40-80

### **Carcinogenicity / Mutagenicity**

Long-term carcinogenicity studies with irinotecan were not conducted. Rats were, however, administered intravenous doses of 2 mg/kg or 25 mg/kg irinotecan once per week for 13 weeks (in separate studies, the 25 mg/kg dose produced an irinotecan  $C_{max}$  and AUC that were about 7.0 times and 1.3 times the respective values in patients administered 125 mg/m<sup>2</sup>) and were then allowed to recover for 91 weeks. Under these conditions, there was a significant linear trend with dose for the incidence of combined uterine horn endometrial stromal polyps and endometrial stromal sarcomas.

Neither irinotecan nor SN-38 was not mutagenic in the in vitro Ames assay. However, in the in vitro Chinese hamster cell chromosomal aberration assay, irinotecan produced a significant increase in the incidence of chromosomal aberrations in a concentration-dependent manner. Additionally, in the in vivo mouse micronucleus assay, a single intraperitoneal dose of irinotecan over the dosage range of 2.5 to 200 mg/kg caused a significant and dose-dependent increase in micronucleated polychromatic erythrocytes and a decrease in the reticulocyte/erythrocyte ratio in bone marrow cells.

### **Reproduction**

No significant adverse effects on fertility and general reproductive performance were observed after intravenous administration of irinotecan in doses of up to 6mg/kg/day to rats. However, atrophy of male reproductive organs was observed after multiple daily irinotecan doses both in rodents at 20 mg/kg (which in separate studies produced an irinotecan  $C_{max}$  and AUC about 5 and 1 times, respectively, the corresponding values in patients administered 125 mg/m<sup>2</sup>) and dogs at 0.4 mg/kg (which in separate studies produced an irinotecan  $C_{max}$  and AUC about one-half and 1/15th, respectively, the corresponding values in patients administered 125 mg/m<sup>2</sup>).

Radioactivity related to C-irinotecan crosses the placenta of rats following intravenous administration of 10 mg/kg (which in separate studies produced an irinotecan  $C_{max}$  and AUC about 3 and 0.5 times, respectively, the corresponding values in patients administered 125 mg/m<sup>2</sup>). Irinotecan was teratogenic in rats at doses greater than 1.2 mg/kg/day (which in separate studies produced an irinotecan  $C_{max}$  and AUC about 2/3 and 1/40th, respectively, of the corresponding values in patients administered 125 mg/m<sup>2</sup>) and in rabbits at 6 mg/kg/day (about one-half the recommended weekly human dose on a mg/m<sup>2</sup> basis). Teratogenic effects included a variety of external, visceral, and skeletal abnormalities. Irinotecan administered to



rat dams for the period following organogenesis through weaning at doses of 6 mg/kg/day caused decreased learning ability and decreased female body weights in the offspring.



## **6. PHARMACEUTICAL PARTICULARS**

### **6.1. List of excipients**

Not applicable

### **6.2. Incompatibilities**

Other drugs should not be added to the infusion solution.

### **6.3. Shelf life**

Do not use Campto after the expiry date which is stated on the carton/vial label after EXP:. The expiry date refers to the last day of that month.

### **6.4. Special precautions for storage**

CAMPTO Injection has been shown to be chemically and physically stable when diluted with infusion solutions (0.9% sodium chloride solution and 5% glucose solution) for up to 28 days when stored in low density polyethylene (LDPE) or polyvinyl chloride (PVC) containers at 5°C or at 30°C/ambient humidity and protected from light. When exposed to light, chemical and physical stability is indicated for up to 3 days.

It is recommended, however, that in order to reduce microbiological hazard, the infusion solutions should be prepared immediately prior to use and infusion commenced as soon as practicable after preparation. If not used immediately, in-use storage times and conditions prior to use should not be longer than 24 hours at 2°C to 8°C or 6 hours at room temperature (25°C).

Do not freeze CAMPTO or admixtures of CAMPTO as this may result in precipitation of the drug.

### **6.5. Nature and contents of container**

Single dose amber polypropylene vials: 40 mg/2 mL, 100 mg/5 mL and 300 mg/15 mL, 1s.

### **6.6. Special precautions for disposal and other handling**

#### **Preparation**

Irinotecan must be diluted prior to infusion in 5% Dextrose Injection, (preferred) or 0.9% Sodium Chloride Injection to a final concentration range of 0.12 to 2.8 mg/mL.

#### **Handling**

As with other potentially toxic anticancer agents, care should be exercised in the handling and preparation of infusion solutions prepared from irinotecan. The use of gloves is recommended. If irinotecan contacts the skin, wash the skin immediately and thoroughly



with soap and water. If irinotecan contacts the mucous membranes, flush thoroughly with water.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Inspect vial contents for particulate matter and repeat inspection when drug product is withdrawn from vial into syringe.

## **7. FURTHER INFORMATION**

### **Name and address of the Manufacturer**

Pfizer (Perth) PTY Ltd.  
15 Brodie Hall Drive, Technology Park  
Bentley WA 6102  
Australia

## **8. PRESCRIPTION STATUS**

Prescription only medicine.

## **9. DATE OF REVISION OF THE TEXT**

August 2018

® Registered Trademark.

## Document Approval Record

**Document Name:** NEAR Master Campto solution LPD

**Document Title:** NEAR Master Campto solution LPD

<b>Signed By:</b>	<b>Date(GMT)</b>	<b>Signing Capacity</b>
Oyinlola, Abimbola Folusho	14-Sep-2018 10:56:38	Regulatory Affairs Approval