	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

1. **Name of the medicinal product:**

Clopidogrel Tablets USP 75mg

2. **Qualitative and quantitative composition:**

Each film coated tablets contains:

Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg

3. **Pharmaceutical form:**

Tablet: A brown color circular shape film coated tablet, plain on both sides.

4. **Clinical particulars:**

**4.1 Therapeutic indications**

Prevention of atherothrombotic events


Clopidogrel is indicated in:

- Adult patients suffering from myocardial infarction (from a few days until less than 35 days), ischaemic stroke (from 7 days until less than 6 months) or established peripheral arterial disease.

- Adult patients suffering from acute coronary syndrome:

- Non-ST segment elevation acute coronary syndrome (unstable angina or non-Q-wave myocardial infarction), including patients undergoing a stent placement following percutaneous coronary intervention, in combination with acetylsalicylic acid (ASA).

- ST segment elevation acute myocardial infarction, in combination with ASA in medically treated patients eligible for thrombolytic therapy.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

Prevention of atherothrombotic and thromboembolic events in atrial fibrillation

In adult patients with atrial fibrillation who have at least one risk factor for vascular events, are not suitable for treatment with Vitamin K antagonists (VKA) and who have a low bleeding risk, clopidogrel is indicated in combination with ASA for the prevention of atherothrombotic and thromboembolic events, including stroke.

#### **4.2 Posology and method of administration**

Posology


- Adults and older people

Clopidogrel should be given as a single daily dose of 75 mg.

In patients suffering from acute coronary syndrome:

- Non-ST segment elevation acute coronary syndrome (unstable angina or non-Q-wave myocardial infarction): clopidogrel treatment should be initiated with a single 300-mg loading dose and then continued at 75 mg once a day (with acetylsalicylic acid (ASA) 75 mg-325 mg daily). Since higher doses of ASA were associated with higher bleeding risk it is recommended that the dose of ASA should not be higher than 100 mg. The optimal duration of treatment has not been formally established. Clinical trial data support use up to 12 months, and the maximum benefit was seen at 3 months.

- ST segment elevation acute myocardial infarction: clopidogrel should be given as a single daily dose of 75 mg initiated with a 300-mg loading dose in combination with ASA and with or without thrombolytics. For patients over 75 years of age clopidogrel should be initiated without a loading dose. Combined therapy should be started as early as possible after symptoms start and continued for at least four weeks. The benefit of the combination of clopidogrel with ASA beyond four weeks has not been studied in this setting.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

In patients with atrial fibrillation, clopidogrel should be given as a single daily dose of 75 mg. ASA (75-100 mg daily) should be initiated and continued in combination with clopidogrel.

If a dose is missed:

- Within less than 12 hours after regular scheduled time: patients should take the dose immediately and then take the next dose at the regular scheduled time.
- For more than 12 hours: patients should take the next dose at the regular scheduled time and should not double the dose.

- Paediatric population

Clopidogrel should not be used in children because of efficacy concerns.

- Renal impairment

Therapeutic experience is limited in patients with renal impairment.

- Hepatic impairment

Therapeutic experience is limited in patients with moderate hepatic disease who may have bleeding diatheses.


Method of administration

For oral use.

It may be given with or without food.

#### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients.
- Severe hepatic impairment.
- Active pathological bleeding such as peptic ulcer or intracranial haemorrhage.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

#### **4.4 Special warnings and precautions for use:**


##### Bleeding and haematological disorders

Due to the risk of bleeding and haematological adverse events, blood cell count determination and/or other appropriate testing should be promptly considered whenever clinical symptoms suggestive of bleeding arise during the course of treatment. As with other antiplatelet agents, clopidogrel should be used with caution in patients who may be at risk of increased bleeding from trauma, surgery or other pathological conditions and in patients receiving treatment with ASA, heparin, glycoprotein IIb/IIIa inhibitors or non-steroidal anti-inflammatory drugs (NSAIDs) including Cox-2 inhibitors, or selective serotonin reuptake inhibitors (SSRIs), or other medicinal products associated with bleeding risk such as pentoxifyllin. Patients should be followed carefully for any signs of bleeding including occult bleeding, especially during the first weeks of treatment and/or after invasive cardiac procedures or surgery. The concomitant administration of clopidogrel with oral anticoagulants is not recommended since it may increase the intensity of bleedings.

If a patient is to undergo elective surgery and antiplatelet effect is temporarily not desirable, clopidogrel should be discontinued 7 days prior to surgery. Patients should inform physicians and dentists that they are taking clopidogrel before any surgery is scheduled and before any new medicinal product is taken. Clopidogrel prolongs bleeding time and should be used with caution in patients who have lesions with a propensity to bleed (particularly gastrointestinal and intraocular).

Patients should be told that it might take longer than usual to stop bleeding when they take clopidogrel (alone or in combination with ASA), and that they should report any unusual bleeding (site or duration) to their physician.

Thrombotic Thrombocytopenic Purpura (TTP)

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

Thrombotic Thrombocytopenic Purpura (TTP) has been reported very rarely following the use of clopidogrel, sometimes after a short exposure. It is characterised by thrombocytopenia and microangiopathic haemolytic anaemia associated with either neurological findings, renal dysfunction or fever. TTP is a potentially fatal condition requiring prompt treatment including plasmapheresis.

#### Acquired haemophilia

Acquired haemophilia has been reported following use of clopidogrel. In cases of confirmed isolated activated Partial Thromboplastin Time (aPTT) prolongation with or without bleeding, acquired haemophilia should be considered. Patients with a confirmed diagnosis of acquired haemophilia should be managed and treated by specialists, and clopidogrel should be discontinued.

#### Recent ischaemic stroke


In view of the lack of data, clopidogrel cannot be recommended during the first 7 days after acute ischaemic stroke.

#### Cytochrome P450 2C19 (CYP2C19)

Pharmacogenetics: In patients who are poor CYP2C19 metabolisers, clopidogrel at recommended doses forms less of the active metabolite of clopidogrel and has a smaller effect on platelet function. Tests are available to identify a patient's CYP2C19 genotype.

Since clopidogrel is metabolized to its active metabolite partly by CYP2C19, use of medicinal products that inhibit the activity of this enzyme would be expected to result in reduced drug levels of the active metabolite of clopidogrel. The clinical relevance of this interaction is uncertain. As a precaution concomitant use of strong or moderate CYP2C19 inhibitors should be discouraged.

Caution is required in patients treated concomitantly with Clopidogrel and CYP2C8

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

substrate medicinal products.

Cross reactions among thienopyridines

Patients should be evaluated for history of hypersensitivity to thienopyridines (such as clopidogrel, ticlopidine, prasugrel) since cross-reactivity among thienopyridines has been reported. Thienopyridines may cause mild to severe allergic reactions such as rash, angioedema, or haematological cross-reactions such as thrombocytopenia and neutropenia. Patients who had developed a previous allergic reaction and/or haematological reaction to one thienopyridine may have an increased risk of developing the same or another reaction to another thienopyridine. Monitoring for signs of hypersensitivity in patients with a known allergy to thienopyridines is advised.

Renal impairment

Therapeutic experience with clopidogrel is limited in patients with renal impairment. Therefore clopidogrel should be used with caution in these patients.

Hepatic impairment


Experience is limited in patients with moderate hepatic disease who may have bleeding diatheses. Clopidogrel should therefore be used with caution in this population.

Excipients

This medicinal product contains hydrogenated castor oil which may cause stomach upset and diarrhoea.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Medicinal products associated with bleeding risk: There is an increased risk of bleeding due to the potential additive effect. The concomitant administration of medicinal products associated with bleeding risk should be undertaken with caution.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	


Oral anticoagulants: the concomitant administration of clopidogrel with oral anticoagulants is not recommended since it may increase the intensity of bleedings. Although the administration of clopidogrel 75 mg/day did not modify the pharmacokinetics of S-warfarin or International Normalised Ratio (INR) in patients receiving long-term warfarin therapy, coadministration of clopidogrel with warfarin increases the risk of bleeding because of independent effects on hemostasis.

Glycoprotein IIb/IIIa inhibitors: clopidogrel should be used with caution in patients who receive concomitant glycoprotein IIb/IIIa inhibitors.

Acetylsalicylic acid (ASA): ASA did not modify the clopidogrel-mediated inhibition of ADP-induced platelet aggregation, but clopidogrel potentiated the effect of ASA on collagen-induced platelet aggregation. However, concomitant administration of 500 mg of ASA twice a day for one day did not significantly increase the prolongation of bleeding time induced by clopidogrel intake. A pharmacodynamic interaction between clopidogrel and acetylsalicylic acid is possible, leading to increased risk of bleeding. Therefore, concomitant use should be undertaken with caution. However, clopidogrel and ASA have been administered together for up to one year.

Heparin: in a clinical study conducted in healthy subjects, clopidogrel did not necessitate modification of the heparin dose or alter the effect of heparin on coagulation. Co-administration of heparin had no effect on the inhibition of platelet aggregation induced by clopidogrel. A pharmacodynamic interaction between clopidogrel and heparin is possible, leading to increased risk of bleeding. Therefore, concomitant use should be undertaken with caution.

Thrombolytics: the safety of the concomitant administration of clopidogrel, fibrin or non-fibrin specific thrombolytic agents and heparins was assessed in patients with acute myocardial infarction. The incidence of clinically significant bleeding was similar to that

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

observed when thrombolytic agents and heparin are co-administered with ASA.

NSAIDs: in a clinical study conducted in healthy volunteers, the concomitant administration of clopidogrel and naproxen increased occult gastrointestinal blood loss. However, due to the lack of interaction studies with other NSAIDs it is presently unclear whether there is an increased risk of gastrointestinal bleeding with all NSAIDs. Consequently, NSAIDs including Cox-2 inhibitors and clopidogrel should be co-administered with caution.

SSRIs: since SSRIs affect platelet activation and increase the risk of bleeding, the concomitant administration of SSRIs with clopidogrel should be undertaken with caution.

Other concomitant therapy: Since clopidogrel is metabolized to its active metabolite partly by CYP2C19, use of medicinal products that inhibit the activity of this enzyme would be expected to result in reduced drug levels of the active metabolite of clopidogrel. The clinical relevance of this interaction is uncertain. As a precaution concomitant use of strong or moderate CYP2C19 inhibitors should be discouraged.


Medicinal products that are strong or moderate CYP2C19 inhibitors include, for example, omeprazole and esomeprazole, fluvoxamine, fluoxetine, moclobemide, voriconazole, fluconazole, ticlopidine, carbamazepine, and efavirenz.

Proton Pump Inhibitors (PPI):

Omeprazole 80 mg once daily administered either at the same time as clopidogrel or with 12 hours between the administrations of the two drugs decreased the exposure of the active metabolite by 45% (loading dose) and 40% (maintenance dose). The decrease was associated with a 39% (loading dose) and 21% (maintenance dose) reduction inhibition of platelet aggregation. Esomeprazole is expected to give a similar interaction with clopidogrel.

Inconsistent data on the clinical implications of this pharmacokinetic



	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

(PK)/pharmacodynamic (PD) interaction in terms of major cardiovascular events have been reported from both observational and clinical studies. As a precaution, concomitant use of omeprazole or esomeprazole should be discouraged.

Less pronounced reductions of metabolite exposure has been observed with pantoprazole or lansoprazole.

The plasma concentrations of the active metabolite was 20% reduced (loading dose) and 14% reduced (maintenance dose) during concomitant treatment with pantoprazole 80 mg once daily. This was associated with a reduction of the mean inhibition of platelet aggregation by 15% and 11%, respectively. These results indicate that clopidogrel can be administered with pantoprazole.


There is no evidence that other medicinal products that reduce stomach acid such as H<sub>2</sub> blockers or antacids interfere with antiplatelet activity of clopidogrel.

Other medicinal products: A number of other clinical studies have been conducted with clopidogrel and other concomitant medicinal products to investigate the potential for pharmacodynamic and pharmacokinetic interactions. No clinically significant pharmacodynamic interactions were observed when clopidogrel was co-administered with atenolol, nifedipine, or both atenolol and nifedipine. Furthermore, the pharmacodynamic activity of clopidogrel was not significantly influenced by the co administration of phenobarbital, or oestrogen.

The pharmacokinetics of digoxin or theophylline were not modified by the co-administration of clopidogrel. Antacids did not modify the extent of clopidogrel absorption.

Data from the CAPRIE study indicate that phenytoin and tolbutamide which are metabolized by CYP2C9 can be safely co-administered with clopidogrel.

CYP2C8 substrate medicinal products: clopidogrel has been shown to increase repaglinide

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

exposure in healthy volunteers. In vitro studies have shown the increase in repaglinide exposure is due to inhibition of CYP2C8 by the glucuronide metabolite of clopidogrel. Due to the risk of increased plasma concentrations, concomitant administration of clopidogrel and drugs primarily cleared by CYP2C8 metabolism (e.g., repaglinide, paclitaxel) should be undertaken with caution.

Apart from the specific medicinal product interaction information described above, interaction studies with clopidogrel and some medicinal products commonly administered in patients with atherothrombotic disease have not been performed. However, patients entered into clinical trials with clopidogrel received a variety of concomitant medicinal products including diuretics, beta blockers, ACEI, calcium antagonists, cholesterol lowering agents, coronary vasodilators, antidiabetic agents (including insulin), antiepileptic agents, and GPIIb/IIIa antagonists without evidence of clinically significant adverse interactions.

#### **4.6 Fertility, Pregnancy and lactation**


##### **Pregnancy**

As no clinical data on exposure to clopidogrel during pregnancy are available, it is preferable not to use clopidogrel during pregnancy as a precautionary measure.

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development.

##### **Breast-feeding**

It is unknown whether clopidogrel is excreted in human breast milk. Animal studies have shown excretion of clopidogrel in breast milk. As a precautionary measure, breast-feeding should not be continued during treatment with Clopidogrel 75 mg Tablets.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

#### **4.7 Effects on ability to drive and use machines**

Undesirable effects such as dizziness, vertigo, drowsiness, fatigue, visual disturbances or other central nervous system disorders are possible after taking NSAIDs. If affected, patients should not drive or operate machinery.

#### **4.8 Undesirable effects**


Summary of the safety profile

Clopidogrel has been evaluated for safety in more than 44,000 patients, who have participated in clinical studies, including over 12,000 patients treated for 1 year or more. The clinically relevant adverse reactions observed in the CAPRIE, CURE, CLARITY, COMMIT and ACTIVE-A studies are discussed below. Overall, clopidogrel 75 mg/day was comparable to ASA 325 mg/day in CAPRIE regardless of age, gender and race. In addition to clinical studies experience, adverse reactions have been spontaneously reported. Bleeding is the most common reaction reported both in clinical studies as well as in post-marketing experience where it was mostly reported during the first month of treatment.

In CAPRIE, in patients treated with either clopidogrel or ASA, the overall incidence of any bleeding was 9.3%. The incidence of severe cases was similar for clopidogrel and ASA.

In CURE, there was no excess in major bleeds with clopidogrel plus ASA within 7 days after coronary bypass graft surgery in patients who stopped therapy more than five days prior to surgery. In patients who remained on therapy within five days of bypass graft surgery, the event rate was 9.6% for clopidogrel plus ASA, and 6.3% for placebo plus ASA.

In CLARITY, there was an overall increase in bleeding in the clopidogrel plus ASA group

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

vs. the placebo plus ASA group. The incidence of major bleeding was similar between groups. This was consistent across subgroups of patients defined by baseline characteristics, and type of fibrinolytic or heparin therapy.

In COMMIT, the overall rate of noncerebral major bleeding or cerebral bleeding was low and similar in both groups.

In ACTIVE-A, the rate of major bleeding was greater in the clopidogrel + ASA group than in the placebo + ASA group (6.7% versus 4.3%). Major bleeding was mostly of extracranial origin in both groups (5.3% in the clopidogrel + ASA group; 3.5% in the placebo +ASA group), mainly from the gastrointestinal tract (3.5% vs. 1.8%). There was an excess of intracranial bleeding in the clopidogrel + ASA treatment group compared to the placebo + ASA group (1.4% versus 0.8%, respectively). There was no statistically significant difference in the rates of fatal bleeding (1.1% in the clopidogrel + ASA group and 0.7% in the placebo +ASA group) and haemorrhagic stroke (0.8% and 0.6%, respectively) between groups.

Tabulated list of adverse reactions

Adverse reactions that occurred either during clinical studies or that were spontaneously reported are presented in the table below. Their frequency is defined using the following conventions: common ( $\geq 1/100$  to  $<1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $<1/1,000$ ); very rare ( $<1/10,000$ ), not known (cannot be estimated from the available data). Within each system organ class, adverse drug reactions are presented in order of decreasing seriousness.



**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

System Class	Organ	Common	Uncommon	Rare	Very rare, not known*
Blood and the lymphatic system disorders			Thrombocytopenia, leucopenia, eosinophilia	Neutropenia, including severe neutropenia	Thrombotic thrombocytopenic purpura (TTP), aplastic anaemia, pancytopenia, agranulocytosis, severe thrombocytopenia, acquired haemophilia A, granulocytopenia, anaemia
Cardiac disorders					Kounis syndrome (vasospastic allergic angina / allergic myocardial infarction) in the context of a hypersensitivity reaction due to clopidogrel*



**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

Immune system disorders				Serum sickness, anaphylactoid reactions, cross- reactive drug hypersensitivity among thienopyridines (such as ticlopidine, prasugrel), insulin autoimmune syndrome, which can lead to severe hypoglycemia, particularly in patients with HLA DRA4 subtype (more frequent in the Japanese population)*
Psychiatric				Hallucinations,



**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

disorders				confusion
Nervous system disorders		Intracranial bleeding (some cases were reported with fatal outcome), headache, paraesthesia, dizziness		Taste disturbances, ageusia
Eye disorders		Eye bleeding (conjunctival, ocular, retinal)		
Ear and labyrinth disorders			Vertigo	
Vascular disorders	Haematoma			Serious haemorrhage, haemorrhage of operative wound, vasculitis, hypotension
Respiratory, thoracic and mediastinal disorders	Epistaxis			Respiratory tract bleeding (haemoptysis, pulmonary



**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

				haemorrhage), bronchospasm, interstitial pneumonitis, eosinophilic pneumonia
Gastrointestinal disorders	Gastrointestinal haemorrhage, diarrhoea, abdominal pain, dyspepsia	Gastric ulcer and duodenal ulcer, gastritis, vomiting, nausea, constipation, flatulence	Retroperitoneal haemorrhage	Gastrointestinal and retroperitoneal haemorrhage with fatal outcome, pancreatitis, colitis (including ulcerative or lymphocytic colitis), stomatitis
Hepato-biliary disorders				Acute liver failure, hepatitis, abnormal liver function test
Skin and	Bruising	Rash, pruritus,		Bullous





**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

subcutaneous tissue disorders		skin bleeding (purpura)		dermatitis (toxic epidermal necrolysis, Stevens Johnson Syndrome, erythema multiforme, acute generalised exanthematous pustulosis (AGEP)), angioedema, drug-induced hypersensitivity syndrome, drug rash with eosinophilia and systemic symptoms (DRESS), rash erythematous or exfoliative, urticaria,
----------------------------------	--	----------------------------	--	---




**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

				eczema, lichen planus
Reproductive systems and breast disorders			Gynaecomastia	
Musculoskeletal, connective tissue and bone disorders				Musculo-skeletal bleeding (haemarthrosis), arthritis, arthralgia, myalgia
Renal and urinary disorders		Haematuria		Glomerulonephritis, blood creatinine increased
General disorders and administration site conditions	Bleeding at puncture site			Fever
Investigations		Bleeding time prolonged, neutrophil count decreased, platelet count decreased		

\*Information related to clopidogrel with frequency “not known”.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

#### **4.9 Overdose:**

Overdose following clopidogrel administration may lead to prolonged bleeding time and subsequent bleeding complications. Appropriate therapy should be considered if bleedings are observed. No antidote to the pharmacological activity of clopidogrel has been found. If prompt correction of prolonged bleeding time is required, platelet transfusion may reverse the effects of clopidogrel.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties:**

Pharmacotherapeutic group: platelet aggregation inhibitors excl. heparin,


ATC code: B01AC-04.

Mechanism of action

Clopidogrel is a prodrug, one of whose metabolites is an inhibitor of platelet aggregation.

Clopidogrel must be metabolized by CYP450 enzymes to produce the active metabolite that inhibits platelet aggregation. The active metabolite of clopidogrel selectively inhibits the binding of adenosine diphosphate (ADP) to its platelet P2Y<sub>12</sub> receptor and the subsequent ADP-mediated activation of the glycoprotein GPIIb/IIIa complex, thereby inhibiting platelet aggregation. Due to the irreversible binding, platelets exposed are affected for the remainder of their lifespan (approximately 7-10 days) and recovery of normal platelet function occurs at a rate consistent with platelet turnover. Platelet aggregation induced by agonists other than ADP is also inhibited by blocking the amplification of platelet activation by released ADP.

Because the active metabolite is formed by CYP450 enzymes, some of which are polymorphic or subject to inhibition by other medicinal products, not all patients will have

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

adequate platelet inhibition.

Pharmacodynamic effects

Repeated doses of 75 mg per day produced substantial inhibition of ADP-induced platelet aggregation from the first day; this increased progressively and reached steady state between Day 3 and Day 7. At steady state, the average inhibition level observed with a dose of 75 mg per day was between 40% and 60%. Platelet aggregation and bleeding time gradually returned to baseline values, generally within 5 days after treatment was discontinued.

## **5.2 Pharmacokinetic properties**

### **Absorption**


After single and repeated oral doses of 75 mg per day, clopidogrel is rapidly absorbed. Mean peak plasma levels of unchanged clopidogrel (approximately 2.2-2.5 ng/ml after a single 75 mg oral dose) occurred approximately 45 minutes after dosing. Absorption is at least 50%, based on urinary excretion of clopidogrel metabolites.

### **Distribution**

Clopidogrel and the main circulating (inactive) metabolite bind reversibly in vitro to human plasma proteins (98% and 94% respectively). The binding is non-saturable in vitro over a wide concentration range.

### **Biotransformation**

Clopidogrel is extensively metabolized by the liver. In vitro and in vivo, clopidogrel is metabolized according to two main metabolic pathways: one mediated by esterases and leading to hydrolysis into its inactive carboxylic acid derivative (85% of circulating metabolites), and one mediated by multiple cytochromes P450. Clopidogrel is first

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

metabolized to a 2 oxo-clopidogrel intermediate metabolite. Subsequent metabolism of the 2 oxo-clopidogrel intermediate metabolite results in formation of the active metabolite, a thiol derivative of clopidogrel. The active metabolite is formed mostly by CYP2C19, with contributions from several other CYP enzymes, including CYP1A2, CYP2B6 and CYP3A4. The active thiol metabolite which has been isolated in vitro, binds rapidly and irreversibly to platelet receptors, thus inhibiting platelet aggregation.

The C<sub>max</sub> of the active metabolite is twice as high following a single 300-mg clopidogrel loading dose as it is after four days of 75-mg maintenance dose. C<sub>max</sub> occurs approximately 30 to 60 minutes after dosing.


#### **Elimination**

Following an oral dose of <sup>14</sup>C-labelled clopidogrel in man, approximately 50% was excreted in the urine and approximately 46% in the faeces in the 120-hour interval after dosing. After a single oral dose of 75 mg, clopidogrel has a half-life of approximately 6 hours. The elimination half-life of the main circulating (inactive) metabolite was 8 hours after single and repeated administration.

#### **Pharmacogenetics**

CYP2C19 is involved in the formation of both the active metabolite and the 2 oxo-clopidogrel intermediate metabolite. Clopidogrel active metabolite pharmacokinetics and antiplatelet effects, as measured by ex vivo platelet aggregation assays, differ according to CYP2C19 genotype.


The CYP2C19\*1 allele corresponds to fully functional metabolism while the CYP2C19\*2 and CYP2C19\*3 alleles are nonfunctional. The CYP2C19\*2 and CYP2C19\*3 alleles account for the majority of reduced function alleles in Caucasian (85%) and Asian (99%) poor metabolisers. Other alleles associated with absent or reduced metabolism are less frequent and include CYP2C19\*4, \*5, \*6, \*7 and \*8. A patient with poor metaboliser

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

status will possess two loss-of-function alleles as defined above. Published frequencies for the poor CYP2C19 metaboliser genotypes are approximately 2% for Caucasians, 4% for Blacks and 14% for Chinese. Tests are available to determine a patient's CYP2C19 genotype.

A crossover study in 40 healthy subjects, 10 each in four CYP2C19 metaboliser groups (ultrarapid, extensive intermediate and poor), evaluated pharmacokinetic and antiplatelet responses using 300 mg followed by 75 mg/day and 600 mg followed by 150mg/day, each for a total of 5 days (steady state). No substantial differences in active metabolite exposure and mean inhibition of platelet aggregation (IPA) were observed between ultrarapid, extensive and intermediate metabolisers. In poor metabolisers, active metabolite exposure was decreased by 63-71% compared to extensive metabolisers. After the 300 mg/75 mg dose regimen, antiplatelet responses were decreased in the poor metabolisers with mean IPA (5µM ADP) of 24% (24 hours) and 37% (Day 5) as compared to IPA of 39% (24 hours) and 58% (Day 5) in the extensive metabolisers and 37% (24 hours) and 60% (Day 5) in the intermediate metabolisers. When poor metabolisers received the 600 mg/150 mg regimen, active metabolite exposure was greater than with the 300 mg/75 mg regimen. In addition, IPA was 32% (24 hours) and 61% (day 5), which were greater than in the poor metabolisers receiving the 300 mg/75 mg regimen, and were similar to the other CYP2C19 metaboliser groups receiving the 300 mg/75 mg regimen. An appropriate dose regimen for this patient population has not been established in clinical outcome trials.

Consistent with the above results, in a meta-analysis including 6 studies of 335 clopidogrel-treated subjects at a steady state, it was shown that active metabolite exposure was decreased by 28% for intermediate metabolisers, and 72% for poor metabolisers while platelet aggregation inhibition (5 µM ADP) was decreased with differences in IPA of 5.9% and 21.4%, respectively, when compared to extensive metabolisers.

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

The influence of CYP2C19 genotype on clinical outcomes in patients treated with clopidogrel has not been evaluated in prospective, randomised, controlled trials. There have been a number of retrospective analyses, however, to evaluate this effect in patients treated with clopidogrel for whom there are genotyping results: CURE (n=2721), CHARISMA (n=2428), CLARITY-TIMI 28 (n=227), TRITON-TIMI 38 (n=1477), and ACTIVE-A (n=601), as well as a number of published cohort studies.

In TRITON-TIMI 38 and 3 of the cohort studies (Collet, Sibbing, Giusti) the combined group of patients with either intermediate or poor metaboliser status had a higher rate of cardiovascular events (death, myocardial infarction and stroke) or stent thrombosis compared to extensive metabolisers.

In CHARISMA and one cohort study (Simon), an increased event rate was observed only in poor metabolisers when compared to extensive metabolisers.

In CURE, CLARITY, ACTIVE-A and one of the cohort studies (Trenk), no increased event rate was observed based on metaboliser status.


None of these analyses were adequately sized to detect differences in outcome in poor metabolisers.

#### Special populations

The pharmacokinetics of the active metabolite of clopidogrel is not known in these special populations

#### Renal impairment

After repeated doses of 75 mg clopidogrel per day in subjects with severe renal disease (creatinine clearance from 5 to 15 ml/min), inhibition of ADP-induced platelet aggregation was lower (25%) than that observed in healthy subjects, however, the prolongation of bleeding time was similar to that seen in healthy subjects receiving 75 mg of clopidogrel

	<b>ATOZ Pharmaceuticals Pvt. Ltd.</b> <b>Chennai – India.</b>
Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

per day. In addition, clinical tolerance was good in all patients.

#### Hepatic impairment

After repeated doses of 75 mg clopidogrel per day for 10 days in patients with severe hepatic impairment, inhibition of ADP-induced platelet aggregation was similar to that observed in healthy subjects. The mean bleeding time prolongation was also similar in the two groups.

#### **5.3 Preclinical safety data:**

During non-clinical studies in rat and baboon, the most frequently observed effects were liver changes. These occurred at doses representing at least 25 times the exposure seen in humans receiving the clinical dose of 75 mg/day and were a consequence of an effect on hepatic metabolising enzymes. No effect on hepatic metabolising enzymes was observed in humans receiving clopidogrel at the therapeutic dose.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of Excipients**

Maize Starch

Microcrystalline Cellulose

Croscarmellose Sodium

Povidone K30

Purified Talc

Colloidal Anhydrous Silica

Crospovidone

Magnesium Stearate

Hypromellose E15





**ATOZ Pharmaceuticals Pvt. Ltd.**  
Chennai – India.

Product Name	<b>FRELET-75 (Clopidogrel Tablets USP 75mg)</b>
Composition	Each film coated tablet contains Clopidogrel Bisulphate USP Equivalent to Clopidogrel 75mg
<b>SUMMARY OF PRODUCT CHARACTERISTICS</b>	

Titanium Dioxide

Iron Oxide of Red

**6.2 Incompatibilities:** Not applicable.

**6.3 Shelf life:** 24 months

**6.4 Special precautions for storage:**

Store below 30°C. Protect from light and Moisture.

**6.5 Nature and contents of container**

Alu-Alu Blister pack of 3x10's Tablets

**6.5 Special precautions for disposal and other handling:**

KEEP OUT OF THE REACH OF CHILDREN