

SUMMARY OF PRODUCT CHARACTERISTICS

Table of Contents

- 1. Name of the medicinal product
- 2. Qualitative and quantitative composition
- 3. Pharmaceutical form
- 4. Clinical particulars
 - 4.1 Therapeutic indications
 - 4.2 Posology and method of administration
 - 4.3 Contraindications
 - 4.4 Special warnings and precautions for use
 - 4.5 Interaction with other medicinal products and other forms of interaction
 - 4.6. Fertility, pregnancy and lactation
 - 4.7 Effects on ability to drive and use machines
 - 4.8 Undesirable effects
 - 4.9 Overdose
- 5. Pharmacological properties
 - 5.1 Pharmacodynamic properties
 - 5.2 Pharmacokinetic properties
 - 5.3 Preclinical safety data
- 6. Pharmaceutical particulars
 - 6.1 List of excipients
 - 6.2 Incompatibilities
 - 6.3 Shelf life
 - 6.4 Special precautions for storage
 - 6.5 Nature and contents of container
 - 6.6 Special precautions for disposal and other handling
- 7. Marketing authorisation holder

1. Name of the medicinal product

MEBEDOL

MEBENDAZOLE ORAL SUSPENSION USP 100 MG/5 ML

2. Qualitative and quantitative composition

Each 5 ml contains:

Mebendazole USP 100 mg

Flavoured Syrupy Base..... Q.S.

Colour: - Quinoline Yellow

3. Pharmaceutical form

Oral suspension.

4. Clinical particulars

4.1 Therapeutic indications

Broad spectrum gastrointestinal anthelmintic indicated for the treatment of:

Enterobius vermicularis (threadworm/pinworm)

Oxyuris vermicularis

Trichuris trichuria (whipworm)

Ascaris lumbricoides (large roundworm)

Ancylostoma duodenale (common hookworm)

Necator americanus (American hookworm)

There is no evidence that Mebendazole suspension is effective in the treatment of cysticercosis.

4.2 Posology and method of administration

Adults and children over 2 years:

Enterobiasis:

1 x 5 ml (1 dosing cup).

It is highly recommended that a second dose is taken after 2 weeks, if reinfection is suspected.

Ascariasis, trichuriasis, ancylostomiasis, necatoriasis and mixed infections:

1 x 5 ml (1 dosing cup) bd for three days.

Children under 2 years:

See section 4.4 Special warnings and precautions for use

Method of administration.

Oral Use

4.3 Contraindications

Mebendazole suspension is contraindicated in pregnancy and in patients who have shown hypersensitivity to the product or any components.

4.4 Special warnings and precautions for use

Not recommended in the treatment of children under 2 years

There have been rare reports of reversible liver function disturbances, hepatitis and neutropenia described in patients who were treated with mebendazole at standard dosages for indicated conditions (see section 4.8 'Undesirable effects'). These events, along with glomerulonephritis and agranulocytosis, have also been reported with dosages substantially above those recommended and with treatment for prolonged periods of time.

A case-control study of a single outbreak of Stevens-Johnson syndrome /toxic epidermal necrolysis (SJS/TEN) suggested a possible association with the concomitant use of metronidazole with mebendazole. Although there are no additional data on this potential interaction, concomitant use of mebendazole and metronidazole should be avoided.

Convulsions in children, including in infants below 1 year of age, have been reported very rarely during post-marketing experience (see section 4.8 'Undesirable effects'). Because of the risk of convulsions, Mebendazole suspension should not be used particularly in children below the age of 1 year. Mebendazole suspension has not been extensively studied in children below the age of 2 years.

Mebendazole suspension should only be given to very young children if their worm infestation interferes significantly with their nutritional status and physical development.

Mebendazole suspension oral suspension contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Concomitant treatment with cimetidine may inhibit the metabolism of mebendazole in the liver, resulting in increased plasma concentrations of the drug.

Concomitant use of mebendazole and metronidazole should be avoided (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

Since Mebendazole suspension is contraindicated in pregnancy, patients who think they are or may be pregnant should not take this preparation.

Breast-feeding

Limited data from case reports demonstrate that a small amount of mebendazole is present in human milk following oral administration. Therefore, caution should be exercised when Mebendazole suspension is administered to breast-feeding women.

4.7 Effects on ability to drive and use machines

Mebendazole suspension has no influence on the ability to drive and use machines.

4.8 Undesirable effects

Throughout this section adverse reactions are reported. Adverse reactions are adverse events that were considered to be reasonably associated with the use of Mebendazole suspension based on the comprehensive assessment of the available adverse event information. A causal relationship with Mebendazole suspension cannot be reliably established in individual cases. Further, because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety of Mebendazole suspension was evaluated in 6276 subjects who participated in 39 clinical trials for the treatment of single or mixed parasitic infestations of the gastrointestinal tract. In these 39 clinical trials, no adverse drug reactions (ADRs) occurred in $\geq 1\%$ of Mebendazole suspension-treated subjects.

ADRs identified from clinical trials and post-marketing experience with Mebendazole suspension are included in Table 1. The displayed frequency categories use the following convention:

Very common ($\geq 1/10$); Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1000$ to $< 1/100$); Rare ($\geq 1/10,000$ to $< 1/1000$); Very rare ($< 1/10,000$), Not known (cannot be estimated from the available data).

Table 1: Adverse Drug Reactions Reported in Clinical Trials and Post-marketing Experience for Mebendazole suspension

System Organ Class	Adverse Drug Reactions		
	Frequency Category		
	Common ($\geq 1/100$ to $< 1/10$)	Uncommon ($\geq 1/1000$ to $< 1/100$)	Rare ($\geq 1/10,000$ to $< 1/1000$)
Blood and Lymphatic System Disorders			Neutropenia ^b Agranulocytosis ^{b*}
Immune System Disorders			Hypersensitivity including anaphylactic reaction and anaphylactoid reaction ^b
Nervous System Disorders			Convulsions ^b Dizziness ^a
Gastrointestinal Disorders	Abdominal pain ^a	Abdominal discomfort ^a ; Diarrhoea ^a ; Flatulence ^a Nausea ^a , Vomiting ^a	
Hepatobiliary Disorders			Hepatitis; ^b Abnormal liver function tests ^b
Skin and Subcutaneous Tissue Disorders			Rash ^a Toxic epidermal necrolysis ^b ; Stevens-Johnson syndrome ^b ; Exanthema ^b ; Angioedema ^b ; Urticaria ^b ; Alopecia ^b
Renal and Urinary Disorders			Glomerulonephritis ^{b*}

^a ADR frequency data derived from Clinical Trials or Epidemiological Studies

^b ADRs not observed in clinical trials and frequency calculated based on 6276 patients exposed in clinical trials and epidemiological studies, divided by 3 (Frequency = 1/2092).

* Observed in higher and prolonged doses

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

In patients treated at dosages substantially higher than recommended or for prolonged periods of time, the following adverse reactions have been reported rarely: alopecia, reversible liver function disturbances, hepatitis, agranulocytosis, neutropenia and glomerulonephritis. With the exception of agranulocytosis and glomerulonephritis, these also have been reported in patients who were treated with mebendazole at standard dosages (see section 4.8).

Signs and symptoms

In the event of accidental overdosage, abdominal cramps, nausea, vomiting and diarrhoea may occur.

Treatment

There is no specific antidote. Activated charcoal may be given if considered appropriate.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: anthelmintic for oral administration, benzimidazole derivatives; ATC code: P02CA01.

In vitro and *in vivo* work suggests that mebendazole blocks the uptake of glucose by adult and larval forms of helminths, in a selective and irreversible manner. Inhibition of glucose uptake appears to lead to endogenous depletion of glycogen stores within the helminth. Lack of glycogen leads to decreased formation of ATP and ultrastructural changes in the cells.

There is no evidence that Mebendazole suspension is effective in the treatment of cysticercosis.

5.2 Pharmacokinetic properties

Absorption

Following oral administration, < 10% of the dose reaches the systemic circulation, due to incomplete absorption and pre-systemic metabolism (first-pass effect). The majority of an orally administered dose remains in the gastrointestinal tract. Maximum plasma concentrations are generally seen 2 to 4 hours after administration. Administration with a high fat meal increases the bioavailability of mebendazole, but the overall effect of food on the amount of drug remaining in the gastrointestinal tract is not expected to be substantial.

Distribution

The plasma protein binding of mebendazole is 90 to 95%. The volume of distribution is 1 to 2 L/kg, indicating that mebendazole penetrates areas outside the vascular space. This is supported by data in patients on chronic mebendazole therapy (e.g., 40 mg/kg/day for 3-21 months) that show drug levels in tissue.

Metabolism

Orally administered mebendazole is extensively metabolised primarily by the liver. Plasma concentrations of its major metabolites (hydrolysed and reduced forms of mebendazole) are substantially higher than those of mebendazole. Impaired hepatic function, impaired metabolism, or impaired biliary elimination may lead to higher plasma levels of mebendazole.

Elimination

Mebendazole, the conjugated forms of mebendazole, and its metabolites likely undergo some degree of enterohepatic recirculation and are excreted in the urine and bile. The apparent elimination half-life after an oral dose ranges from 3 to 6 hours in most patients.

Steady-state pharmacokinetics

During chronic dosing (e.g., 40 mg/kg/day for 3-21 months), plasma concentrations of mebendazole and its major metabolites increase, resulting in approximately 3-fold higher exposure at steady-state compared to single dosing.

5.3 Preclinical safety data

In animal reproduction studies, adverse developmental effects (i.e., skeletal malformations, soft tissue malformations, decreased pup weight, embryoletality) were observed when mebendazole was administered to pregnant rats and mice throughout the period of organogenesis or as a single oral dose as low as 10 mg/kg in rats (approximately 0.2-fold the maximum recommended human dose (MRHD)). Maternal toxicity was present at the highest of these doses. Dosing of hamsters and rabbits did not result in embryotoxicity or teratogenicity. Doses up to 40 mg/kg in rats (0.8-fold the MRHD, based on mg/m²), given to males for 60 days and to females for 14 days prior to gestation, had no effect upon foetuses and offspring.

No mutagenic activity was observed with mebendazole in bacterial reverse mutation tests. Mebendazole was mutagenic when tested in the mouse lymphoma thymidine kinase assay and

aneugenic in vitro in mammalian somatic cells. In the *in vivo* mouse micronucleus assay, orally administered mebendazole induced an increased frequency of micronucleated polychromatic erythrocytes with evidence suggestive of aneugenicity.

Mebendazole had no carcinogenic effects at doses as high as 40 mg/kg/day when administered daily in the diet over 2 years in carcinogenicity tests in mice and rats (0.4 to 0.8-fold the MRHD, based on mg/m²).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

1. Sucrose BP
2. Sorbitol 70% BP
3. Glycerin BP
4. Sodium Methyl Hydroxybenzoate BP
5. Sodium Propyl Hydroxybenzoate BP
6. Citric Acid Monohydrate BP
7. Poly Sorbate-80 BP
8. Sodium Citrate BP
9. Colloidal Anhydrous Silica BP
10. Xanthan Gum BP
11. Color: Quinoline yellow IHS
12. Flavour Vanilla liquid IHS
13. Purified water BP

6.2 Incompatibilities

Not applicable

6.3 Shelf life

2 years

6.4 Special precautions for storage

Store at a temperature not exceeding 30°C. Protected from light.

6.5 Nature and contents of container

Primary Packing: 30 ml Amber glass bottles with PP cap and measuring cup.

Secondary Packing: 1 such bottle in one carton along with a package insert.

6.6 Special precautions for disposal and other handling

Shake well before use

7. MARKETING AUTHORISATION HOLDER

Company Name : GB PHARMA LIMITED
Address : 65 Chatsworth Road, London NW2 4BG
Country : United Kingdom
Phone : +44 (0)2088301057
Fax : + 44 (0) 2088304807
E-mail : info@gbpharma.co.uk