

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

**Tulip® 10 mg film-coated tablets**

**Tulip® 20 mg film-coated tablets**

**Tulip® 30 mg film-coated tablets**

**Tulip® 40 mg film-coated tablets**

**Tulip® 80 mg film-coated tablets**

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

*Tulip® 10 mg film-coated tablets*

Each 10 mg film-coated tablet contains 10 mg of atorvastatin, equivalent to 10.34 mg of atorvastatin calcium.

Excipients: lactose monohydrate. Each tablet contains 33.06 mg of lactose, equivalent to 34.80 mg of lactose monohydrate.

*Tulip® 20 mg film-coated tablets*

Each 20 mg film-coated tablet contains 20 mg of atorvastatin, equivalent to 20.68 mg of atorvastatin calcium.

Excipients: lactose monohydrate. Each tablet contains 33.06 mg of lactose, equivalent to 34.80 mg of lactose monohydrate.

*Tulip® 30 mg film-coated tablets*

Each 30 mg film-coated tablet contains 30 mg of atorvastatin, equivalent to 31.0 mg of atorvastatin calcium.

Excipients: lactose monohydrate. Each tablet contains 49.59 mg of lactose, equivalent to 52.20 mg of lactose monohydrate.

*Tulip® 40 mg film-coated tablets*

Each 40 mg film-coated tablet contains 40 mg of atorvastatin, equivalent to 41.36 mg of atorvastatin calcium.

Excipients: lactose monohydrate. Each tablet contains 66.12 mg of lactose, equivalent to 69.60 mg of lactose monohydrate.

*Tulip® 80 mg film-coated tablets*

Each 80 mg film-coated tablet contains 80 mg of atorvastatin, equivalent to 82.7 mg of atorvastatin calcium.

Excipients: lactose monohydrate. Each tablet contains 132.24 mg of lactose, equivalent to 139.20 mg of lactose monohydrate.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Film-coated tablet

Tulip 10 mg tablets are white to off-white, round, biconvex, debossed with »HLA 10« on one side.

Tulip 20 mg tablets are light yellow, round, biconvex, debossed with »HLA 20« on one side.

Tulip 30 mg tablets are light yellow, round, biconvex, debossed with »HLA 30« on one side.

Tulip 40 mg tablets are light yellow, round, biconvex, debossed with »HLA 40« on one side.

Tulip 80 mg tablets are light yellow, oval, biconvex, debossed with »HLA 80« on one side.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Hypercholesterolaemia

Tulip is indicated as an adjunct to diet for reduction of elevated total cholesterol (total-C), LDL-cholesterol (LDL-C), apolipoprotein B, and triglycerides in adults, adolescents and children aged 10 years or older with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate.

Tulip is also indicated to reduce total-C and LDL-C in adults with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.

#### Prevention of cardiovascular disease

Prevention of cardiovascular events in patients estimated to have a high risk for a first cardiovascular event (see section 5.1), as an adjunct to correction of other risk factors.

### **4.2 Posology and method of administration**

#### Posology

The patient should be placed on a standard cholesterol-lowering diet before receiving Tulip and should continue on this diet during treatment with Tulip.

The dose should be individualised according to baseline LDL-C levels, the goal of therapy, and patient response.

The usual starting dose is 10 mg once a day. Adjustment of dose should be made at intervals of 4 weeks or more. The maximum dose is 80 mg once a day.

#### Primary hypercholesterolaemia and combined (mixed) hyperlipidaemia

The majority of patients are controlled with Tulip 10 mg once a day.

A therapeutic response is evident within 2 weeks, and the maximum therapeutic response is usually achieved within 4 weeks. The response is maintained during chronic therapy.

#### Heterozygous familial hypercholesterolaemia

Patients should be started with Tulip 10 mg daily. Doses should be individualised and adjusted every 4 weeks to 40 mg daily. Thereafter, either the dose may be increased to a

maximum of 80 mg daily or a bile acid sequestrant may be combined with 40 mg atorvastatin once daily.

#### Homozygous familial hypercholesterolaemia

Only limited data are available (see section 5.1).

The dose of atorvastatin in patients with homozygous familial hypercholesterolaemia is 10 to 80 mg daily (see section 5.1). Atorvastatin should be used as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) in these patients or if such treatments are unavailable.

#### Prevention of cardiovascular disease

In the primary prevention trials the dose was 10 mg/day. Higher doses may be necessary in order to attain LDL cholesterol levels according to current guidelines.

#### Renal impairment

No adjustment of dose is required (see section 4.4).

#### Hepatic impairment

Tulip should be used with caution in patients with hepatic impairment (see sections 4.4 and 5.2). Tulip is contraindicated in patients with active liver disease (see section 4.3).

#### Use in the elderly

Efficacy and safety in patients older than 70 years of age using recommended doses are similar to those seen in the general population.

#### Paediatric use

##### Hypercholesterolaemia:

Paediatric use should only be carried out by physicians experienced in the treatment of paediatric hyperlipidaemia and patients should be re-evaluated on a regular basis to assess progress.

For patients aged 10 years and above, the recommended starting dose of atorvastatin is 10 mg per day with titration up to 20 mg per day. Titration should be conducted according to the individual response and tolerability in paediatric patients. Safety information for paediatric patients treated with doses above 20 mg, corresponding to about 0.5 mg/kg, is limited.

There is limited experience in children between 6 to 10 years of age (see section 5.1). Atorvastatin is not indicated in the treatment of patients below the age of 10 years.

Other pharmaceutical forms/strengths may be more appropriate for this population.

#### Method of administration

Tulip is for oral administration. Each daily dose of atorvastatin should be taken all at once and may be taken at any time of day with or without food. It is advisable to take the medicinal product each day at the same time of the day.

### **4.3 Contraindications**

Tulip is contraindicated:

- in patients with hypersensitivity to the active substance or to any of the excipients of this medicinal product
- in patients with active liver disease or unexplained persistent elevations of serum transaminases exceeding 3 times the upper limit of normal (ULN)
- during pregnancy, while breast-feeding and in women of child-bearing potential not using appropriate contraceptive measures (see section 4.6).

### **4.4 Special warnings and precautions for use**

#### Liver effects

Liver function tests should be performed before the initiation of treatment and periodically thereafter. Patients who develop any signs or symptoms suggestive of liver injury should have liver function tests performed. Patients who develop increased transaminase levels should be monitored until the abnormality(ies) resolve. Should an increase in transaminases of greater than 3 times the upper limit of normal (ULN) persist, reduction of dose or withdrawal of Tulip is recommended (see section 4.8).

Tulip should be used with caution in patients who consume substantial quantities of alcohol and/or have a history of liver disease.

#### SPARCL study (Stroke Prevention by Aggressive Reduction in Cholesterol Levels)

In a *post-hoc* analysis of stroke subtypes in patients without coronary heart disease (CHD) who had a recent stroke or transient ischaemic attack (TIA) there was a higher incidence of haemorrhagic stroke in patients initiated on atorvastatin 80 mg compared to placebo. The increased risk was particularly noted in patients with prior haemorrhagic stroke or lacunar infarct at study entry. For patients with prior haemorrhagic stroke or lacunar infarct, the balance of risks and benefits of atorvastatin 80 mg is uncertain, and the potential risk of haemorrhagic stroke should be carefully considered before initiating treatment (see section 5.1).

#### Skeletal muscle effects

Atorvastatin, like other 3-hydroxy-3-methyl-glutaryl-coenzyme A (HMG-CoA) reductase inhibitors, may in rare occasions affect the skeletal muscle and cause myalgia, myositis, and myopathy that may progress to rhabdomyolysis, a potentially life-threatening condition characterised by markedly elevated creatinine kinase (CK) levels (> 10 times ULN), myoglobinaemia and myoglobinuria which may lead to renal failure.

#### **Before the treatment**

Atorvastatin should be prescribed with caution in patients with pre-disposing factors for rhabdomyolysis.

A CK level should be measured before starting statin treatment in the following situations:

- renal impairment
- hypothyroidism
- personal or familial history of hereditary muscular disorders

- previous history of muscular toxicity with a statin or fibrate
- previous history of liver disease and/or where substantial quantities of alcohol are consumed
- in elderly patients (age > 70 years), the necessity of such measurement should be considered, according to the presence of other predisposing factors for rhabdomyolysis
- situations where an increase in plasma levels may occur, such as interactions (see section 4.5) and special populations including genetic subpopulations (see section 5.2)

In such situations, the risk of treatment should be considered in relation to possible benefit, and clinical monitoring is recommended. If CK levels are significantly elevated (> 5 times ULN) at baseline, treatment should not be started.

### **Creatine kinase measurement**

Creatine kinase (CK) should not be measured following strenuous exercise or in the presence of any plausible alternative cause of CK increase as this makes value interpretation difficult. If CK levels are significantly elevated at baseline (> 5 times ULN), levels should be remeasured within 5 to 7 days later to confirm the results.

### **Whilst on treatment**

- Patients must be asked to promptly report muscle pain, cramps, or weakness especially if accompanied by malaise or fever.
- If such symptoms occur whilst a patient is receiving treatment with atorvastatin, their CK levels should be measured. If these levels are found to be significantly elevated (> 5 times ULN), treatment should be stopped.
- If muscular symptoms are severe and cause daily discomfort, even if the CK levels are elevated to  $\leq 5 \times$  ULN, treatment discontinuation should be considered.
- If symptoms resolve and CK levels return to normal, then re-introduction of atorvastatin or introduction of an alternative statin may be considered at the lowest dose and with close monitoring.
- Atorvastatin must be discontinued if clinically significant elevation of CK levels (> 10 x ULN) occur, or if rhabdomyolysis is diagnosed or suspected.

### Concomitant treatment with other medicinal products

Risk of rhabdomyolysis is increased when atorvastatin is administered concomitantly with certain medicinal products that may increase the plasma concentration of atorvastatin such as potent inhibitors of cytochrome P450 3A4 (CYP3A4) or transport proteins (e.g. cyclosporine, telithromycin, clarithromycin, delavirdine, stiripentol, ketoconazole, voriconazole, itraconazole, posaconazole and HIV protease inhibitors including ritonavir, lopinavir, atazanavir, indinavir, darunavir, etc.). The risk of myopathy may also be increased with the concomitant use of gemfibrozil and other fibric acid derivatives, erythromycin, niacin and ezetimibe. If possible, alternative (non-interacting) therapies should be considered instead of these medicinal products.

In cases where co-administration of these medicinal products with atorvastatin is necessary, the benefit and the risk of concurrent treatment should be carefully considered. When patients are receiving medicinal products that increase the plasma concentration of atorvastatin, a lower maximum dose of atorvastatin is recommended. In addition, in the case of potent CYP3A4 inhibitors, a lower starting dose of atorvastatin should be considered and appropriate clinical monitoring of these patients is recommended (see section 4.5).

The concurrent use of atorvastatin and fusidic acid is not recommended, therefore, temporary suspension of atorvastatin may be considered during fusidic acid therapy (see section 4.5).

### Paediatric use

Developmental safety in the paediatric population has not been established (see section 4.8).

### Interstitial lung disease

Exceptional cases of interstitial lung disease have been reported with some statins, especially with long term therapy (see section 4.8). Presenting features can include dyspnoea, non-productive cough and deterioration in general health (fatigue, weight loss and fever). If it is suspected a patient has developed interstitial lung disease, statin therapy should be discontinued.

### Diabetes mellitus

Some evidence suggests that statins as a class raise blood glucose and in some patients, at high risk of future diabetes, may produce a level of hyperglycaemia where formal diabetes care is appropriate. This risk, however, is outweighed by the reduction in vascular risk with statins and therefore should not be a reason for stopping statin treatment. Patients at risk (fasting glucose 5.6 to 6.9 mmol/L, BMI > 30kg/m<sup>2</sup>, raised triglycerides, hypertension) should be monitored both clinically and biochemically according to national guidelines.

### **Lactose**

Tulip contains inactive ingredient lactose. Patients with rare hereditary problems of galactose intolerance, Lapp lactose deficiency or glucose-galactose malabsorption should not take this medicinal product.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Effect of co-administered medicinal products on atorvastatin

Atorvastatin is metabolised by cytochrome P450 3A4 (CYP3A4) and is a substrate to transport proteins e.g. the hepatic uptake transporter OATP1B1. Concomitant administration of medicinal products that are inhibitors of CYP3A4 or transport proteins may lead to increased plasma concentrations of atorvastatin and an increased risk of myopathy. The risk might also be increased at concomitant administration of atorvastatin with other medicinal products that have a potential to induce myopathy, such as fibric acid derivatives and ezetimibe (see section 4.4).

### CYP3A4 inhibitors

Potent CYP3A4 inhibitors have been shown to lead to markedly increased concentrations of atorvastatin (see Table 1 and specific information below). Co-administration of potent CYP3A4 inhibitors (e.g. cyclosporine, telithromycin, clarithromycin, delavirdine, stiripentol, ketoconazole, voriconazole, itraconazole, posaconazole and HIV protease inhibitors including ritonavir, lopinavir, atazanavir, indinavir, darunavir, etc.) should be avoided if possible. In cases where co-administration of these medicinal products with atorvastatin cannot be avoided lower starting and maximum doses of atorvastatin should be considered and appropriate clinical monitoring of the patient is recommended (see Table 1).

Moderate CYP3A4 inhibitors (e.g. erythromycin, diltiazem, verapamil and fluconazole) may increase plasma concentrations of atorvastatin (see Table 1). An increased risk of myopathy has been observed with the use of erythromycin in combination with statins. Interaction studies evaluating the effects of amiodarone or verapamil on atorvastatin have not been conducted. Both amiodarone and verapamil are known to inhibit CYP3A4 activity and co-administration with atorvastatin may result in increased exposure to atorvastatin. Therefore, a lower maximum dose of atorvastatin should be considered and appropriate clinical monitoring of the patient is recommended when concomitantly used with moderate CYP3A4 inhibitors. Appropriate clinical monitoring is recommended after initiation or following dose adjustments of the inhibitor.

#### CYP3A4 inducers

Concomitant administration of atorvastatin with inducers of cytochrome P450 3A (e.g. efavirenz, rifampin, St. John's Wort) can lead to variable reductions in plasma concentrations of atorvastatin. Due to the dual interaction mechanism of rifampin (cytochrome P450 3A induction and inhibition of hepatocyte uptake transporter OATP1B1), simultaneous co-administration of atorvastatin with rifampin is recommended, as delayed administration of atorvastatin after administration of rifampin has been associated with a significant reduction in atorvastatin plasma concentrations. The effect of rifampin on atorvastatin concentrations in hepatocytes is, however, unknown and if concomitant administration cannot be avoided, patients should be carefully monitored for efficacy.

#### Transport protein inhibitors

Inhibitors of transport proteins (e.g. cyclosporine) can increase the systemic exposure of atorvastatin (see Table 1). The effect of inhibition of hepatic uptake transporters on atorvastatin concentrations in hepatocytes is unknown. If concomitant administration cannot be avoided, a dose reduction and clinical monitoring for efficacy is recommended (see Table 1).

#### Gemfibrozil / fibric acid derivatives

The use of fibrates alone is occasionally associated with muscle related events, including rhabdomyolysis. The risk of these events may be increased with the concomitant use of fibric acid derivatives and atorvastatin. If concomitant administration cannot be avoided, the lowest dose of atorvastatin to achieve the therapeutic objective should be used and the patients should be appropriately monitored (see section 4.4).

#### Ezetimibe

The use of ezetimibe alone is associated with muscle related events, including rhabdomyolysis. The risk of these events may therefore be increased with concomitant use of ezetimibe and atorvastatin. Appropriate clinical monitoring of these patients is recommended.

#### Colestipol

Plasma concentrations of atorvastatin and its active metabolites were lower (by approx. 25 percent) when colestipol was co-administered with Tulip. However, lipid effects were greater when Tulip and colestipol were co-administered than when either medicinal product was given alone.

### Fusidic acid

Interaction studies with atorvastatin and fusidic acid have not been conducted. As with other statins, muscle related events, including rhabdomyolysis, have been reported in post-marketing experience with atorvastatin and fusidic acid given concurrently. The mechanism of this interaction is not known. Patients should be closely monitored and temporary suspension of atorvastatin treatment may be appropriate.

### Effect of atorvastatin on co-administered medicinal products

#### Digoxin

When multiple doses of digoxin and 10 mg atorvastatin were co-administered, steady-state digoxin concentrations increased slightly. Patients taking digoxin should be monitored appropriately.

#### Oral contraceptives

Co-administration of Tulip with an oral contraceptive produced increases in plasma concentrations of norethindrone and ethinyl oestradiol.

#### Warfarin

In a clinical study in patients receiving chronic warfarin therapy, coadministration of atorvastatin 80 mg daily with warfarin caused a small decrease of about 1.7 seconds in prothrombin time during the first 4 days of dosing which returned to normal within 15 days of atorvastatin treatment. Although only very rare cases of clinically significant anticoagulant interactions have been reported, prothrombin time should be determined before starting atorvastatin in patients taking coumarin anticoagulants and frequently enough during early therapy to ensure that no significant alteration of prothrombin time occurs. Once a stable prothrombin time has been documented, prothrombin times can be monitored at the intervals usually recommended for patients on coumarin anticoagulants. If the dose of atorvastatin is changed or discontinued, the same procedure should be repeated. Atorvastatin therapy has not been associated with bleeding or with changes in prothrombin time in patients not taking anticoagulants.

### Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

**Table 1: Effect of co-administered medicinal products on the pharmacokinetics of atorvastatin**

Co-administered medicinal product and dosing regimen	Atorvastatin		
	Dose (mg)	Change in AUC <sup>&amp;</sup>	Clinical recommendation <sup>#</sup>

Tipranavir 500 mg BID/ Ritonavir 200 mg BID, 8 days (days 14 to 21)	40 mg on day 1, 10 mg on day 20	↑ 9.4 fold	In cases where coadministration with atorvastatin is necessary, do not exceed 10 mg atorvastatin daily. Clinical monitoring of these patients is recommended.
Cyclosporine 5.2 mg/kg/day, stable dose	10 mg OD for 28 days	↑ 8.7 fold	
Lopinavir 400 mg BID/ Ritonavir 100 mg BID, 14 days	20 mg OD for 4 days	↑ 5.9 fold	In cases where co-administration with atorvastatin is necessary, lower maintenance doses of atorvastatin are recommended. At atorvastatin doses exceeding 20 mg, clinical monitoring of these patients is recommended.
Clarithromycin 500 mg BID, 9 days	80 mg OD for 8 days	↑ 4.4 fold	
Saquinavir 400 mg BID/ Ritonavir (300 mg BID from days 5-7, increased to 400 mg BID on day 8), days 5-18, 30 min after atorvastatin dosing	40 mg OD for 4 days	↑ 3.9 fold	In cases where co-administration with atorvastatin is necessary, lower maintenance doses of atorvastatin are recommended. At atorvastatin doses exceeding 40 mg, clinical monitoring of these patients is recommended.
Darunavir 300 mg BID/Ritonavir 100 mg BID, 9 days	10 mg OD for 4 days	↑ 3.3 fold	
Itraconazole 200 mg OD, 4 days	40 mg SD	↑ 3.3 fold	
Fosamprenavir 700 mg BID/ Ritonavir 100 mg BID, 14 days	10 mg OD for 4 days	↑ 2.5 fold	
Fosamprenavir 1400 mg BID, 14 days	10 mg OD for 4 days	↑ 2.3 fold	
Nelfinavir 1250 mg BID, 14 days	10 mg OD for 28 days	↑ 1.7 fold <sup>^</sup>	No specific recommendation
Grapefruit Juice, 240 mL OD*	40 mg, SD	↑ 37 %	Concomitant intake of large quantities of grapefruit juice and atorvastatin is not recommended.
Diltiazem 240 mg OD, 28 days	40 mg, SD	↑ 51 %	After initiation or following dose adjustments of diltiazem, appropriate clinical monitoring of these patients is recommended.
Erythromycin 500 mg QID, 7 days	10 mg, SD	↑ 33 % <sup>^</sup>	Lower maximum dose and clinical monitoring of these patients is recommended.
Amlodipine 10 mg, single dose	80 mg, SD	↑ 18 %	No specific recommendation.
Cimetidine 300 mg QID, 2 weeks	10 mg OD for 4 weeks	↓ less than 1 % <sup>^</sup>	No specific recommendation.
Antacid suspension of magnesium and aluminium hydroxides, 30 mL QID, 2 weeks	10 mg OD for 4 weeks	↓ 35 % <sup>^</sup>	No specific recommendation.
Efavirenz 600 mg OD, 14 days	10 mg for 3 days	↓ 41 %	No specific recommendation.
Rifampin 600 mg OD, 7 days (co-administered)	40 mg SD	↑ 30 %	If co-administration cannot be avoided, simultaneous co- administration of atorvastatin with rifampin is recommended, with clinical monitoring.
Rifampin 600 mg OD, 5 days (doses separated)	40 mg SD	↓ 80 %	

Gemfibrozil 600 mg BID, 7 days	40 mg SD	↑ 35 %	Lower starting dose and clinical monitoring of these patients is recommended.
Fenofibrate 160 mg OD, 7 days	40 mg SD	↑ 3 %	Lower starting dose and clinical monitoring of these patients is recommended.

& Data given as x-fold change represent a simple ratio between co-administration and atorvastatin alone (i.e., 1-fold = no change). Data given as % change represent % difference relative to atorvastatin alone (i.e., 0 % = no change).

# See sections 4.4 and 4.5 for clinical significance.

\* Contains one or more components that inhibit CYP3A4 and can increase plasma concentrations of medicinal products metabolised by CYP3A4. Intake of one 240 ml glass of grapefruit juice also resulted in a decreased AUC of 20.4 % for the active orthohydroxy metabolite. Large quantities of grapefruit juice (over 1.2 l daily for 5 days) increased AUC of atorvastatin 2.5 fold and AUC of active (atorvastatin and metabolites).

^ Total atorvastatin equivalent activity

Increase is indicated as "↑", decrease as "↓"

OD = once daily; SD = single dose; BID = twice daily; QID = four times daily

Table 2: Effect of atorvastatin on the pharmacokinetics of co-administered medicinal products

Atorvastatin and dosing regimen	Co-administered medicinal product		
	Medicinal product/Dose (mg)	Change in AUC&	Clinical recommendation
80 mg OD for 10 days	Digoxin 0.25 mg OD, 20 days	↑ 15 %	Patients taking digoxin should be monitored appropriately.
40 mg OD for 22 days	Oral contraceptive OD, 2 months - norethindrone 1 mg - ethinyl estradiol 35 µg	↑ 28 % ↑ 19 %	No specific recommendation.
80 mg OD for 15 days	* Phenazone, 600 mg SD	↑ 3 %	No specific recommendation

& Data given as % change represent % difference relative to atorvastatin alone (i.e., 0 % = no change)

\* Co-administration of multiple doses of atorvastatin and phenazone showed little or no detectable effect in the clearance of phenazone.

Increase is indicated as "↑", decrease as "↓"

OD = once daily; SD = single dose

#### 4.6 Fertility, pregnancy and lactation

##### Women of childbearing potential

Women of child-bearing potential should use appropriate contraceptive measures during treatment (see section 4.3).

## Pregnancy

Tulip is contraindicated during pregnancy (see section 4.3). Safety in pregnant women has not been established. No controlled clinical trials with atorvastatin have been conducted in pregnant women. Rare reports of congenital anomalies following intrauterine exposure to HMG-CoA reductase inhibitors have been received. Animal studies have shown toxicity to reproduction (see section 5.3).

Maternal treatment with atorvastatin may reduce the foetal levels of mevalonate which is a precursor of cholesterol biosynthesis. Atherosclerosis is a chronic process, and ordinarily discontinuation of lipid-lowering medicinal products during pregnancy should have little impact on the long-term risk associated with primary hypercholesterolaemia.

For these reasons, Tulip should not be used in women who are pregnant, trying to become pregnant or suspect they are pregnant. Treatment with Tulip should be suspended for the duration of pregnancy or until it has been determined that the woman is not pregnant (see section 4.3.).

## Lactation

It is not known whether atorvastatin or its metabolites are excreted in human milk. In rats, plasma concentrations of atorvastatin and its active metabolites are similar to those in milk (see section 5.3). Because of the potential for serious adverse reactions, women taking Tulip should not breast-feed their infants (see section 4.3). Atorvastatin is contraindicated during breastfeeding (see section 4.3).

## Fertility

In animal studies atorvastatin had no effect on male or female fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Tulip has negligible influence on the ability to drive and use machines.

### **4.8 Undesirable effects**

In the atorvastatin placebo-controlled clinical trial database of 16,066 (Tulip n=8755; placebo n=7311) patients treated for a mean period of 53 weeks, 5.2 % of patients on atorvastatin discontinued due to adverse reactions compared to 4.0 % of the patients on placebo.

Based on data from clinical studies and extensive post-marketing experience with atorvastatin, the following table presents the adverse reaction profile for atorvastatin.

Estimated frequencies of reactions are ranked according to the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $1/10$ ); uncommon ( $\geq 1/1,000$  to  $1/100$ ); rare ( $\geq 1/10,000$  to  $1/1,000$ ); very rare ( $\leq 1/10,000$ ); not known (cannot be estimated from the available data)

#### Infections and infestations

Common: nasopharyngitis

#### Blood and lymphatic system disorders

Rare: thrombocytopenia

### Immune system disorders

Common: allergic reactions

Very rare: anaphylaxis

### Metabolism and nutrition disorders

Common: hyperglycaemia

Uncommon: hypoglycaemia, weight gain, anorexia

### Psychiatric disorders

Uncommon: nightmares, insomnia

### Nervous system disorders

Common: headache

Uncommon: dizziness, paraesthesia, hypoesthesia, dysgeusia, amnesia

Rare: peripheral neuropathy

### Eye disorders

Uncommon: blurred vision

Rare: visual disturbances

### Ear and labyrinth disorders

Uncommon: tinnitus

Very rare: hearing loss

### Respiratory, thoracic and mediastinal disorders

Common: pharyngolaryngeal pain, epistaxis

### Gastrointestinal disorders

Common: constipation, flatulence, dyspepsia, nausea, diarrhoea

Uncommon: vomiting, abdominal pain upper and lower, eructation, pancreatitis

### Hepatobiliary disorders

Uncommon: hepatitis

Rare: cholestatic jaundice

Very rare: hepatic failure

### Skin and subcutaneous tissue disorders

Uncommon: urticaria, skin rash, pruritus, alopecia

Rare: angioneurotic oedema, bullous dermatitis (including erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis)

### Musculoskeletal and connective tissue disorders

Common: myalgia, arthralgia, pain in extremity, muscle spasms, joint swelling, back pain

Uncommon: neck pain, muscle fatigue

Rare: myositis, rhabdomyolysis, myopathy, tendinopathy, sometimes complicated by tendon rupture

### Reproductive system and breast disorders

Very rare: gynaecomastia

### General disorders and administration site conditions

Uncommon: malaise, asthenia, chest pain, peripheral oedema, fatigue, pyrexia

### Investigations

Common: liver function test abnormal, blood creatine phosphokinase increased

Uncommon: white blood cells urine positive

As with other HMG-CoA reductase inhibitors elevated serum transaminases have been reported in patients receiving Tulip. These changes were usually mild, transient, and did not require interruption of treatment. Clinically important (> 3 times upper normal limit) elevations in serum transaminases occurred in 0.8 % patients on Tulip. These elevations were dose related and were reversible in all patients.

Elevated serum creatine phosphokinase (CPK) levels greater than 3 times upper limit of normal occurred in 2.5 % of patients on Tulip, similar to other HMG-CoA reductase inhibitors in clinical trials. Levels above 10 times the normal upper range occurred in 0.4 % Tulip-treated patients (see section 4.4).

### Paediatric population

The clinical safety database includes safety data for 249 paediatric patients who received atorvastatin, among which 7 patients were less than 6 years old, 14 patients were in the age range of 6 to 9, and 228 patients were in the age range of 10 to 17.

### *Nervous system disorders*

Common: headache

### *Gastrointestinal disorders*

Common: abdominal pain

### Investigations

Common: alanine aminotransferase increased, blood creatine phosphokinase increased

Based on the data available, frequency, type and severity of adverse reactions in children are expected to be the same as in adults. There is currently limited experience with respect to long-term safety in the paediatric population.

### Class effect of statins

The following undesirable effects have been reported with some statins:

- Sexual dysfunction
- Depression
- Exceptional cases of interstitial lung disease, especially with long term therapy (see section 4.4)
- Diabetes mellitus: frequency will depend on the presence or absence of risk factors (fasting blood glucose  $\geq 5.6$  mmol/L, BMI  $> 30$ kg/m<sup>2</sup>, raised triglycerides, history of hypertension).

### Reporting of suspected undesirable effects

Reporting of suspected undesirable effects after the marketing authorisation is important. It enables continuous monitoring of the drug's risk/benefit ratio. Health care professionals are required to report any suspected undesirable effect of the medicinal product to the national centre for reporting:

Univerzitetni klinični center Ljubljana, Interna klinika  
Center za zastrupitve  
Zaloška cesta 2

SI-1000 Ljubljana  
Telefax: + 386 (0)1 434 76 46  
E-mail: farmakovigilanca@kclj.si

#### **4.9 Overdose**

Specific treatment is not available for Tulip overdose. Should an overdose occur, the patient should be treated symptomatically and supportive measures instituted, as required. Liver function tests should be performed and serum CPK levels should be monitored. Due to extensive atorvastatin binding to plasma proteins, haemodialysis is not expected to significantly enhance atorvastatin clearance.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Lipid modifying agents, without combinations, HMG-CoA-reductase inhibitors.  
ATC code: C10AA05

Atorvastatin is a selective, competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme responsible for the conversion of 3-hydroxy-3-methyl-glutaryl-coenzyme A to mevalonate, a precursor of sterols, including cholesterol. Triglycerides and cholesterol in the liver are incorporated into very low-density lipoproteins (VLDL) and released into the plasma for delivery to peripheral tissues. Low-density lipoprotein (LDL) is formed from VLDL and is catabolised primarily through the receptor with high affinity to LDL (LDL receptor).

Atorvastatin lowers plasma cholesterol and lipoprotein serum concentrations by inhibiting HMG-CoA reductase and subsequently cholesterol biosynthesis in the liver and increases the number of hepatic LDL receptors on the cell surface for enhanced uptake and catabolism of LDL.

Atorvastatin reduces LDL production and the number of LDL particles. Additionally, atorvastatin produces a profound and sustained increase in LDL receptor activity coupled with a beneficial change in the quality of circulating LDL particles. Atorvastatin is effective in reducing LDL-C in patients with homozygous familial hypercholesterolaemia, a population that has not usually responded to lipid-lowering medicinal products.

Atorvastatin has been shown to reduce concentrations of total-C (30 % - 46 %), LDL-C (41 % - 61 %), apolipoprotein B (34 % - 50 %), and triglycerides (14 % - 33 %) while producing variable increases in HDL-C and apolipoprotein A1 in a dose response study. These results are consistent in patients with heterozygous familial hypercholesterolaemia, nonfamilial forms of hypercholesterolaemia, and mixed hyperlipidaemia, including patients with noninsulin-dependent diabetes mellitus.

Reductions in total-C, LDL-C, and apolipoprotein B have been proven to reduce risk for cardiovascular events and cardiovascular mortality.

## Homozygous familial hypercholesterolaemia

In a multicenter 8 week open-label compassionate-use study with an optional extension phase of variable length, 335 patients were enrolled, 89 of which were identified as homozygous familial hypercholesterolaemia patients. From these 89 patients, the mean percent reduction in LDL-C was approximately 20 %. Atorvastatin was administered at doses up to 80 mg/day.

## Atherosclerosis

In the REVERSAL study (*Reversing Atherosclerosis with Aggressive Lipid- Lowering Study*), the effect of intensive lipid lowering with atorvastatin 80 mg and standard degree of lipid lowering with pravastatin 40 mg on coronary atherosclerosis was assessed by intravascular ultrasound, during angiography, in patients with coronary heart disease. In this randomised, double- blind, multicenter, controlled clinical trial, intravascular ultrasound was performed at baseline and at 18 months in 502 patients. In the atorvastatin group (n=253), there was no progression of atherosclerosis.

The median percent change, from baseline, in total atheroma volume (the primary study criteria) was -0.4 % (p=0.98) in the atorvastatin group and +2.7 % (p=0.001) in the pravastatin group (n=249). When compared to pravastatin the effects of atorvastatin were statistically significant (p=0.02). The effect of intensive lipid lowering on cardiovascular endpoints (e. g. need for revascularisation, non fatal myocardial infarction, coronary death) was not investigated in this study.

In the atorvastatin group, LDL-C was reduced to a median of 2.04 mmol/L  $\pm$  0.8 (78.9 mg/dl  $\pm$  30) from baseline 3.89 mmol/l  $\pm$  0.7 (150 mg/dl  $\pm$  28) and in the pravastatin group, LDL-C was reduced to a median of 2.85 mmol/l  $\pm$  0.7 (110 mg/dl  $\pm$  26) from baseline 3.89 mmol/l  $\pm$  0.7 (150 mg/dl  $\pm$  26) (p<0.0001). Atorvastatin also significantly reduced mean TC by 34.1 % (pravastatin: -18.4 %, p<0.0001), median TG levels by 20 % (pravastatin: -6.8 %, p<0.0009), and median apolipoprotein B by 39.1 % (pravastatin: -22.0 %, p<0.0001). Atorvastatin increased mean HDL-C by 2.9 % (pravastatin: +5.6 %, p=NS). There was a 36.4 % mean reduction in the reactive protein (CRP) in the atorvastatin group compared to a 5.2 % reduction in the pravastatin group (p<0.0001).

Study results were obtained with the 80 mg dose strength. Therefore, they cannot be extrapolated to the lower dose strengths.

The safety and tolerability profiles of the two treatment groups were comparable.

The effect of intensive lipid lowering on major cardiovascular endpoints was not investigated in this study. Therefore, the clinical significance of these imaging results with regard to the primary and secondary prevention of cardiovascular events is unknown.

## Acute coronary syndrome

In the MIRACL study, atorvastatin 80 mg has been evaluated in 3,086 patients (atorvastatin n=1,538; placebo n=1,548) with an acute coronary syndrome (non Q-wave myocardial infarction or unstable angina). Treatment was initiated during the acute phase after hospital admission and lasted for a period of 16 weeks. Treatment with atorvastatin 80 mg/day increased the time to occurrence of the combined primary endpoint, defined as death from any cause, nonfatal myocardial infarction, resuscitated cardiac arrest, or angina pectoris with evidence of myocardial ischaemia requiring hospitalisation,

indicating a risk reduction by 16 % (p=0.048). This was mainly due to a 26 % reduction in re-hospitalisation for angina pectoris with evidence of myocardial ischaemia (p=0.018). The other secondary endpoints did not reach statistical significance on their own (overall: Placebo: 22.2 %, Atorvastatin: 22.4 %).

The safety profile of atorvastatin in the MIRACL study was consistent with what is described in section 4.8.

#### Prevention of cardiovascular disease

The effect of atorvastatin on fatal and non-fatal coronary heart disease was assessed in a randomised, double-blind, placebo-controlled study, the Anglo-Scandinavian Cardiac Outcomes Trial Lipid Lowering Arm (ASCOT-LLA). Patients were hypertensive, 40 to 79 years of age, with no previous myocardial infarction or treatment for angina, and with TC levels  $\leq 6.5$  mmol/l (251 mg/dl). All patients had at least 3 of the pre-defined cardiovascular risk factors: male gender, age  $\geq 55$  years, smoking, diabetes, history of coronary heart disease in a first-degree relative, TC:HDL-C  $>6$ , peripheral vascular disease, left ventricular hypertrophy, prior cerebrovascular event, specific ECG abnormality, proteinuria/albuminuria. Not all included patients were estimated to have a high risk for a first cardiovascular event.

Patients were treated with anti-hypertensive therapy (either amlodipine or atenolol-based regimen) and either atorvastatin 10 mg daily (n=5,168) or placebo (n=5,137).

The absolute and relative risk reduction effect of atorvastatin was as follows:

Event	Relative risk reduction (%)	No. of events (atorvastatin vs. placebo)	Absolute risk reduction <sup>1</sup> (%)	p-value
Fatal CHD plus non-fatal MI	36 %	100 vs. 154	1.1 %	0.0005
Total cardiovascular events and revascularisation procedures	20 %	389 vs. 483	1.9 %	0.0008
Total coronary events	29 %	178 vs. 247	1.4 %	0.0006
<sup>1</sup> Based on difference in crude events rates occurring over a median follow-up of 3.3 years. CHD = coronary heart disease; MI = myocardial infarction.				

Total mortality and cardiovascular mortality were not significantly reduced (185 vs. 212 events, p=0.17 and 74 vs. 82 events, p=0.51). In the subgroup analyses by gender (81 % males, 19 % females), a beneficial effect of atorvastatin was seen in males but could not be established in females possibly due to the low event rate in the female subgroup. Overall and cardiovascular mortality were numerically higher in the female patients (38 vs. 30 and 17 vs. 12), but this was not statistically significant. There was significant treatment interaction by antihypertensive baseline therapy. The primary endpoint (fatal CHD plus non-fatal myocardial infarction) was significantly reduced by atorvastatin in patients treated with amlodipine (hazard ratio 0.47 (0.32-0.69), p=0.00008), but not in those treated with atenolol (hazard ratio 0.83 (0.59-1.17), p=0.287).

The effect of atorvastatin on fatal and non-fatal cardiovascular disease was also assessed in a randomised, double-blind, multicenter, placebo-controlled trial, the Collaborative Atorvastatin Diabetes Study (CARDS) in patients with type 2 diabetes, 40 to 75 years of age, without prior history of cardiovascular disease, and with LDL-C  $\leq$  4.14 mmol/l (160 mg/dl) and TG  $\leq$  6.78 mmol/l (600 mg/dl). All patients had at least one of the following risk factors: hypertension, current smoking, retinopathy, microalbuminuria or macroalbuminuria.

Patients were treated with either atorvastatin 10 mg daily (n=1,428) or placebo (n=1,410) for a median follow-up of 3.9 years.

The absolute and relative risk reduction effect of atorvastatin was as follows:

Event	Relative risk reduction (%)	No. of events (atorvastatin vs. placebo)	Absolute risk reduction <sup>1</sup> (%)	p-value
Major cardiovascular events (fatal and non-fatal AMI, silent MI, acute CHD death, unstable angina, CABG, PTCA, revascularisation, stroke)	37 %	83 vs. 127	3.2 %	0.0010
MI (fatal and non-fatal AMI, silent MI)	42 %	38 vs. 64	1.9 %	0.0070
Strokes (Fatal and non-fatal)	48 %	21 vs. 39	1.3 %	0.0163
<sup>1</sup> Based on difference in crude events rates occurring over a median follow-up of 3.9 years.				
AMI = acute myocardial infarction; CABG = coronary artery bypass graft; CHD = coronary heart disease; MI = myocardial infarction; PTCA = percutaneous transluminal coronary angioplasty.				

There was no evidence of a difference in the treatment effect by patient's gender, age, or baseline LDL-C level. A favourable trend was observed regarding the mortality rate (82 deaths in the placebo group vs. 61 deaths in the atorvastatin group, p=0.0592).

### Recurrent stroke

In the SPARCL study (Stroke Prevention by Aggressive Reduction in Cholesterol Levels), the effect of atorvastatin 80 mg daily or placebo on stroke was evaluated in 4731 patients who had a stroke or transient ischaemic attack (TIA) within the preceding 6 months and no history of coronary heart disease (CHD). Patients were 60 % male, 21 to 92 years of age (average age 63 years), and had an average baseline LDL of 133 mg/dL (3.4 mmol/L). The mean LDL-C was 73 mg/dL (1.9 mmol/L) during treatment with atorvastatin and 129 mg/dL (3.3 mmol/L) during treatment with placebo. Median follow-up was 4.9 years.

Atorvastatin 80 mg reduced the risk of the primary endpoint of fatal or non-fatal stroke by 15 % (hazard ratio 0.85; 95 % CI, 0.72-1.00; p=0.05 or 0.84; 95 % CI, 0.71-0.99; p=0.03 after adjustment for baseline factors) compared to placebo. All cause mortality was 9.1 % (216/2365) for atorvastatin versus 8.9 % (211/2366) for placebo.

In a *post-hoc* analysis, atorvastatin 80 mg reduced the incidence of ischaemic stroke (218/2365, 9.2 % vs. 274/2366, 11.6 %,  $p=0.01$ ) and increased the incidence of haemorrhagic stroke (55/2365, 2.3 % vs. 33/2366, 1.4 %,  $p=0.02$ ) compared to placebo.

- The risk of haemorrhagic stroke was increased in patients who entered the study with prior haemorrhagic stroke (7/45 for atorvastatin versus 2/48 for placebo; hazard ratio 4.06; 95 % CI, 0.84-19.57), and the risk of ischaemic stroke was similar between groups (3/45 for atorvastatin versus 2/48 for placebo; hazard ratio 1.64; 95 % CI, 0.27-9.82).
- The risk of haemorrhagic stroke was increased in patients who entered the study with prior lacunar infarct (20/708 for atorvastatin versus 4/701 for placebo; hazard ratio 4.99; 95 % CI, 1.71-14.61), but the risk of ischaemic stroke was also decreased in these patients (79/708 for atorvastatin versus 102/701 for placebo; hazard ratio 0.76; 95 % CI, 0.57-1.02). It is possible that the net risk of stroke is increased in patients with prior lacunar infarct who receive atorvastatin 80 mg/day.

All cause mortality was 15.6 % (7/45) for atorvastatin versus 10.4 % (5/48) in the subgroup of patients with prior haemorrhagic stroke. All cause mortality was 10.9 % (77/708) for atorvastatin versus 9.1 % (64/701) for placebo in the subgroup of patients with prior lacunar infarct.

Paediatric population

#### *Heterozygous familial hypercholesterolaemia in paediatric patients aged 6 to 17 years*

An 8-week, open-label study to evaluate pharmacokinetics, pharmacodynamics, and safety and tolerability of atorvastatin was conducted in children and adolescents with genetically confirmed heterozygous familial hypercholesterolemia and baseline LDL-C  $\geq 4$  mmol/L. A total of 39 children and adolescents, 6 to 17 years of age, were enrolled. Cohort A included 15 children, 6 to 12 years of age and at Tanner Stage 1. Cohort B included 24 children, 10 to 17 years of age and at Tanner Stage  $\geq 2$ .

The initial dose of atorvastatin was 5 mg daily of a chewable tablet in Cohort A and 10 mg daily of a tablet formulation in Cohort B. The atorvastatin dose was permitted to be doubled if a subject had not attained target LDL-C of  $<3.35$  mmol/L at Week 4 and if atorvastatin was well tolerated.

Mean values for LDL-C, TC, VLDL-C, and Apo B decreased by Week 2 among all subjects. For subjects whose dose was doubled, additional decreases were observed as early as 2 weeks, at the first assessment, after dose escalation. The mean percent decreases in lipid parameters were similar for both cohorts, regardless of whether subjects remained at their initial dose or doubled their initial dose. At Week 8, on average, the percent change from baseline in LDL-C and TC was approximately 40 % and 30 %, respectively, over the range of exposures.

#### *Heterozygous familial hypercholesterolaemia in paediatric patients aged 10 to 17 years*

In a double-blind, placebo controlled study followed by an open-label phase, 187 boys and postmenarchal girls 10 to 17 years of age (mean age 14.1 years) with heterozygous familial hypercholesterolaemia (FH) or severe hypercholesterolaemia were randomised to atorvastatin ( $n=140$ ) or placebo ( $n=47$ ) for 26 weeks and then all received atorvastatin for 26 weeks. The dosage of atorvastatin (once daily) was 10 mg for the first 4 weeks and up-titrated to 20 mg if the LDL-C level was  $>3.36$  mmol/l. Atorvastatin significantly decreased plasma levels of total-C, LDL-C, triglycerides, and apolipoprotein B during the 26 week double-blind phase. The group compared to 5.91 mmol/l (range: 3.93-9.96 mmol/l) in the placebo group during the 26-week double-blind phase.

An additional paediatric study of atorvastatin versus colestipol in patients with hypercholesterolaemia aged 10 to 18 years demonstrated that atorvastatin (N=25) caused a significant reduction in LDL-C at week 26 ( $p < 0.05$ ) compared with colestipol (N=31).

A compassionate use study in patients with severe hypercholesterolaemia (including homozygous hypercholesterolaemia) included 46 paediatric patients treated with atorvastatin titrated according to response (some subjects received 80 mg atorvastatin per day). The study lasted 3 years: LDL-cholesterol was lowered by 36 %.

The long-term efficacy of atorvastatin therapy in childhood to reduce morbidity and mortality in adulthood has not been established.

The European Medicines Agency has waived the obligation to submit the results of studies with atorvastatin in children aged 0 to less than 6 years in the treatment of heterozygous hypercholesterolaemia and in children aged 0 to less than 18 years in the treatment of homozygous familial hypercholesterolaemia, combined (mixed) hypercholesterolaemia, primary hypercholesterolaemia and in the prevention of cardiovascular events (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Atorvastatin is rapidly absorbed after oral administration; maximum plasma concentrations ( $C_{max}$ ) occur within 1 to 2 hours. Extent of absorption increases in proportion to atorvastatin dose. After oral administration, atorvastatin film-coated tablets are 95 % to 99 % bioavailable compared to the oral solution. The absolute bioavailability of atorvastatin is approximately 12 % and the systemic availability of HMG-CoA reductase inhibitory activity is approximately 30 %. The low systemic availability is attributed to presystemic clearance in gastrointestinal mucosa and/or hepatic first-pass metabolism.

### Distribution

Mean volume of distribution of atorvastatin is approximately 381 litres. Atorvastatin is  $\geq 98$  % bound to plasma proteins.

### Metabolism

Atorvastatin is metabolised by cytochrome P450 3A4 to ortho- and parahydroxylated derivatives and various beta-oxidation products. Apart from other pathways these products are further metabolised via glucuronidation. *In vitro*, inhibition of HMG-CoA reductase by ortho- and parahydroxylated metabolites is equivalent to that of atorvastatin. Approximately 70 % of circulating inhibitory activity for HMG-CoA reductase is attributed to active metabolites.

### Excretion

Atorvastatin is eliminated primarily in bile following hepatic and/or extrahepatic metabolism. However, atorvastatin does not appear to undergo significant enterohepatic recirculation. Mean plasma elimination half-life of atorvastatin in humans is approximately 14 hours. The half-life of inhibitory activity for HMG-CoA reductase is approximately 20 to 30 hours due to the contribution of active metabolites.

## Special populations

*Elderly:* Plasma concentrations of atorvastatin and its active metabolites are higher in healthy elderly subjects than in young adults while the lipid effects were comparable to those seen in younger patient populations.

*Paediatric population:* In an open-label, 8-week study, Tanner Stage 1 (N=15) and Tanner Stage  $\geq 2$  (N=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of film-coated atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.

*Gender:* Concentrations of atorvastatin and its active metabolites in women differ from those in men (women: approx. 20 % higher for  $C_{max}$  and approx. 10 % lower for AUC). These differences were of no clinical significance, resulting in no clinically significant differences in lipid effects among men and women.

*Renal insufficiency:* Renal disease has no influence on the plasma concentrations or lipid effects of atorvastatin and its active metabolites.

*Hepatic insufficiency:* Plasma concentrations of atorvastatin and its active metabolites are markedly increased (approx. 16-fold in  $C_{max}$  and approx. 11-fold in AUC) in patients with chronic alcoholic liver disease (Child-Pugh B).

*SLCO1B1 polymorphism:* Hepatic uptake of all HMG-CoA reductase inhibitors including atorvastatin, involves the OATP1B1 transporter. In patients with SLCO1B1 polymorphism there is a risk of increased exposure of atorvastatin, which may lead to an increased risk of rhabdomyolysis (see section 4.4). Polymorphism in the gene encoding OATP1B1 (SLCO1B1 c.521CC) is associated with a 2.4-fold higher atorvastatin exposure (AUC) than in individuals without this genotype variant (c.521TT). A genetically impaired hepatic uptake of atorvastatin is also possible in these patients. Possible consequences for the efficacy are unknown.

### **5.3 Preclinical safety data**

Atorvastatin was negative for mutagenic and clastogenic potential in a battery of 4 *in vitro* tests and 1 *in vivo* assay. Atorvastatin was not found to be carcinogenic in rats, but high doses in mice (resulting in 6 to 11 fold the  $AUC_{0-24h}$  reached in humans at the highest recommended dose) showed hepatocellular adenomas in males and hepatocellular carcinomas in females.

There is evidence from animal experimental studies that HMG-CoA reductase inhibitors may affect the development of embryos or foetuses. In rats, rabbits and dogs atorvastatin had no effect on fertility and was not teratogenic, however, at maternally toxic doses foetal toxicity was observed in rats and rabbits. The development of the rat offspring was delayed and post-natal survival reduced during exposure of the dams to high doses of atorvastatin. In rats, there is evidence of placental transfer. In rats, plasma concentrations of atorvastatin are similar to those in milk. It is not known whether atorvastatin or its metabolites are excreted in human milk.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### **Tulip 10 mg film-coated tablets**

Core: microcrystalline cellulose (E460), lactose monohydrate, croscarmellose sodium (E468), hydroxypropyl cellulose (E463), polysorbate 80 (E433), heavy magnesium oxide (E530), colloidal anhydrous silica (E551), magnesium stearate (E572).

Coating: hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), macrogol 6000, talc (E553b).

#### **Tulip 20 mg film-coated tablets**

Core: microcrystalline cellulose (E460), lactose monohydrate, croscarmellose sodium (E468), hydroxypropyl cellulose (E463), polysorbate 80 (E433), heavy magnesium oxide (E530), colloidal anhydrous silica (E551), magnesium stearate (E572).

Coating: hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), macrogol 6000, yellow iron oxide (E172), talc (E553b).

#### **Tulip 30 mg film-coated tablets**

Core: microcrystalline cellulose (E460), lactose monohydrate, croscarmellose sodium (E468), hydroxypropyl cellulose (E463), polysorbate 80 (E433), heavy magnesium oxide (E530), colloidal anhydrous silica (E551), magnesium stearate (E572).

Coating: hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), macrogol 6000, yellow iron oxide (E172), talc (E553b).

#### **Tulip 40 mg film-coated tablets**

Core: microcrystalline cellulose (E460), lactose monohydrate, croscarmellose sodium (E468), hydroxypropyl cellulose (E463), polysorbate 80 (E433), heavy magnesium oxide (E530), colloidal anhydrous silica (E551), magnesium stearate (E572).

Coating: hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), macrogol 6000, yellow iron oxide (E172), talc (E553b).

#### **Tulip 80 mg film-coated tablets**

Core: microcrystalline cellulose (E460), lactose monohydrate, croscarmellose sodium (E468), hydroxypropyl cellulose (E463), polysorbate 80 (E433), heavy magnesium oxide (E530), colloidal anhydrous silica (E551), magnesium stearate (E572).

Coating: hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), macrogol 6000, talc (E553b), yellow iron oxide (E172).

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

3 years.

### 6.4 Special precautions for storage

Store in the original package in order to protect from moisture.

#### **6.5 Nature and contents of container**

##### **Tulip 10 mg film-coated tablets**

Blister made of hard and soft aluminium foil, box containing 30, 60 or 90 film-coated tablets in a blister pack (3 x 10, 6 x 10, 9 x 10).

##### **Tulip 20 mg film-coated tablets**

Blister made of hard and soft aluminium foil, box containing 30, 60 or 90 film-coated tablets in a blister pack (3 x 10, 6 x 10, 9 x 10).

##### **Tulip 30 mg film-coated tablets**

Blister made of hard and soft aluminium foil, box containing 30 film-coated tablets in a blister pack (3 x 10).

##### **Tulip 40 mg film-coated tablets**

Blister made of hard and soft aluminium foil, box containing 30, 60 or 90 film-coated tablets in a blister pack (3 x 10, 6 x 10, 9 x 10).

##### **Tulip 80 mg film-coated tablets**

Blister made of hard and soft aluminium foil, box containing 30 film-coated tablets in a blister pack (6 x 5).

#### **6.6 Special precautions for disposal and other handling**

No special requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Lek Pharmaceuticals d.d., Verovškova 57, Ljubljana, Slovenia.

### **8. MARKETING AUTHORISATION NUMBER(S)**

H/01/01575/001-011

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: December 14, 2001

Date of latest renewal: July 28, 2011

### **10. DATE OF REVISION OF THE TEXT**

March 9, 2015